Quantum Mechanics
for
Agnes Scott College
Lifelong Learning Class

Professor Emeritus of Theoretical Physics
John Boccio
Swarthmore College

Table of Contents

I - Basic Class Lectures

1 - Beginning Ideas and Thoughts 001

2 - A Simple Introduction to Quantum Ideas via Experiments 042

3 - Applying Mathematics ideas to Quantum Mechanics 110

4 - Stern-Gerlach Experiments 151

5 - Quantum States/Postulates - Real Meaning of Formalism 173

6 - Some Review, Some New Details and Explicit Time Evolution 261

7 - Entanglement, EPR and Bell 341

8 - Quantum Measurement 452

II - Supplementary Readings connected to Lectures

1 - The Mathematics Of Quantum Mechanics - 1st Installment 619

2 - The Einstein-Bohr-God Debate 663

3 - The Saga of Two Gloves 672

4 - Light and Interference 679

5 - The Gambler's Ruin Problem and Quantum Measurement 695

6 - Mathematics Extra 719

III - Discussion by other Authors

1 - Questions, challenges, and new ideas about Quantum Measurement 723

Decoherence and Quantum Measurement - Last Pass Summary

How Do the Probabilities Arise in Quantum Measurement?

Quantum measurement: a game between observer and nature?

Can Decoherence Solve the Measurement Problem?

5 - The Born Rule - Axiom or Result? 806

6 - The Development of the Interpretation of the Quantum Theory according to Heisenberg 816

7 - Taking Heisenberg's Potentia Seriously 850

8 - A Final Summary — Quantum fluctuations, particles and entanglement: Solving the quantum measurement problem 873

9 - Another Way by Sidney Coleman 910

IV - Advanced material for the Adventurous Student

1 - Polarization 952

2 - Schrödinger Equation 1037

3 - Density Matrix and Decoherence 1072

4 - Two Discussions of Measurement 1139

Start of Part #1

By the end of this class you will understand these sentences:

Everything we call real is made of things that cannot be regarded as real.

Quantum mechanics asks: Is FALSE the SAME as NOT TRUE?

What is quantum mechanics trying to tell us?
“Correlations have physical reality, that which they correlate does not"

Beginning Ideas and Thoughts

Two Big questions. How does the universe work?

What are the fundamental principles involved?

Any discussions/answers must involve Physics, Mathematics AND possibly Philosophy

19th century versus 20th century

Physicists arrived at very different ways of thinking.

Classical Physics
Determinism
Locality
Foundation of physics 1650-1920

Quantum Physics
Randomness
Non-locality
Foundation of all modern physics 1920 -2024

These two paths offer radically different views of reality

For our present purposes: Let us ask: What is quantum mechanics?

Before 1920 physicists thought they understood everything, but world was not what it seemed!

Microscopic world = Quantum Macroscopic world = Classical (but not so sure now!)

Clear they needed a new theory to explain new experiments being done in the Microworld.

Our everyday experience(classical) is very misleading, i.e.,
it leads to ideas that do not work in the microworld as we will see.

We will find that QM => the fundamental building blocks of everything
=> the us with an ability to develop a theory of microworld

Two important new properties arise.

They are adequately summed up by the words

Potentiality and Uncertainty

Must use Experiments —allow development→ Postulates and Tools ==> QM Theory

The Goals of this Class

How QM works
Why it must work this way
How to understand it
What it all means

My assumptions about you

You remember some HS math + you are willing to learn some new mathematics.
Basically, you will be learning a new language!

You are willing to **completely** change your view about how the universe works.

To discuss QM - we first need to agree on some terminology from Philosophy

Epistemology versus **Ontology**

↓ ↓

Study of things we just know versus **Study of what is actually out there**

↓ ↓

Realist **versus** **Instrumentalist**

↓ ↓

Science = reality versus **Theory must agree with data**

↓ ↓

Objects are really there versus **Does not care if objects are really there**

Example to think about ……..

When we study the **state** of a QM System

It will **seem** to depend on what **we know.**

Realist ⟶ will say state is not real because our knowledge can change

Instrumentalist ⟶ will say all OK, it is just our information that changes

The Classical Point of View due to Newton and Maxwell

Fundamental concept = particle trajectory in physical variable(position/time) spaces

System = collection of particles **isolated** from universe

State = Values of set of observables or measurable quantities ⟶ Trajectory = particle state

State = collection of particle states

Extrinsic versus Intrinsic observables

↓ ↓

position/velocity versus mass/charge

Information is known with infinite precision (in principle)

Limited only by instrument precision but **not** by Nature

Outcome of all measurements can be predicted using trajectories

Newton's Laws of Motion ⟶ { Position+velocity+momentum+energy+force
Requires knowledge of position/velocity **at some time**.

—> initial conditions


They believed that if we can determine initial conditions without disturbance.

then could **predict** trajectory ⟶ then know all **future** behavior

Belief: accurate predictions limited only by accuracy of initial conditions + knowledge of forces

These ideas work very well for macroworld for centuries - **We did land on the moon!!**

One fly in the ointment = **chaos** ⟶ Sensitivity to initial conditions ⟶ not predictable



There is no randomness involved in this chaotic motion; it just looks like it!

In QM, as it turns out, we will discover that **true** randomness actually exists

Concept of a Newtonian Clock ==> a **Determinate** Universe

Everything we observe **had** a cause in the past!
Causality is **fundamental** in this classical world
If same cause **,** then same effect! (**QM will say this is WRONG!!**)
It will seem like there is no **free will** (and **maybe** there is none!)

Now we go through the Looking Glass: The Quantum Point of View(#1)

Will observe that "classical particles" **do not exist** in the microword!

⟶ we will **need** a radical revision of the definition of a particle.

We will struggle with understanding **when/why** it seems that a "Particle **not** a Particle"?

We will see: objects seem to behave like a classical particle sometimes **and** seem to behave like a classical wave sometimes —→ the so-called **wave-particle duality** that you have all probably heard about!

We will find, however, that quantum particles are **neither** classical particles **nor** classical waves.

They only **seem** to be in particular circumstances!

WE MUST KEEP AN OPEN MIND

Object properties **may have nothing** to do with either classical idea(waves or particles)!
All such classical ideas may be **totally** wrong!!

If we do find classical physics is WRONG(it disagrees with experiment), **then we should not** try to make quantum things be like the classical things that do not work!

To set the stage for our later discussions - here are some of the strange quantum ideas that will appear in during those discussions:

(1) Quantum properties are not well-defined **until** measured

(2) QM rarely gives a **definite** answer to any question

(3) The quantum state will be a mixture of **all** possibilities

(4) QM will **only** predict possibilities and probabilities

(5) QM will **seem** statistical - i.e., seem to only apply to ensembles of systems.

(6) Random chance will **seem** to control everything

(7) Will find that when measurement gives value - then repeat - get same value
—> **HAS** that value at that moment

(8) Will find that before measurement there is **no** value! **Only** a **potentiality** exists!

(9) Will find that measurement **alters** system - it is unavoidable - effect cannot be zero!

(10) Will wonder if particles exist when their properties are **not** being measured?

(11) Will ask, is what we study in physics, reality or merely our **perceptions** of reality?

There are many more I could list!! **WOW!**

The Decline and Fall of the Trajectory (remember : Trajectory = Classical View)

trajectory = set of (positions, momentum, time) for a particle

where first elements of the set = **initial conditions**

ensemble of "identical" systems = set of "identical" trajectories
= set of "points + errors"

classical world —> errors can be made small as we wish(no theoretical limit) - **ASSUMPTION**

quantum world —> this is **NOT TRUE** <— exists an uncertainty principle = fundamental limit

True randomness in quantum world **will destroy** the old classical deterministic idea!

Question: Can Classical and Quantum Physics Get Together Somewhere?


Since limit of large systems in microworld ⟶ macro world

does QM ⟶ CM in this limit? Is there a "dividing line"?

Bohr proposal - Correspondence Principle

Where is dividing line? The so-called **Heisenberg Cut**

No one knows - it may not exist!!



QM will try to tell us - will have to see what it says!!

QM will tell us:

Do not make any statements that can not be verified.

or put another way……

We can only know properties that can be measured.

Physics(QM), as we will see, is an **experimental** science

We get ideas from experiments and theory predictions **must be verified by experiment**

If particle **is** measured to be at (x_1,t_1) and (x_2,t_2), **then** particle **was** at x_1 at t_1 **and** x_2 at t_2

Since there were no measurements, QM **cannot** talk about particle in **between** two points

i.e., we did not measure particle to be anywhere else, so we cannot say it was in QM!

Clearly, this strange manner of thinking will affect the QM view of Reality

The study of QM is **difficult for many reasons**:

(1) We are **macro objects — here is a quote —**

> **Things on a very small scale behave like nothing you have any direct experience about. They do not behave like waves, they do not behave like particles, they do not behave like clouds, or billiard balls, or weights on springs, or like anything that you have ever seen.**
>
> — Feynman

(2) The microworld can be understood, but it **cannot be seen(with our eyes!)**.
It can only measured with sensitive instruments which our eyes can then read.

(3) Quantum mechanics is inherently **mathematical. Why?**

Because **Mathematics** is the **language** of quantum physics.

It will only make sense when described with mathematics.

We must deal with that fact and we **will**!

Be prepared to **give up** many ideas that you did not possibly think could be wrong!!!!

Let us proceed! ⟶ **Into the Microworld we go.**

Like Alice though the Looking Glass!

Our **first pass** through the quantum ideas and experiments will now be done **without** knowing the theory that follows.

3 important ideas dominate: **Quantization, Uncertainty and Duality**

Quantization: Macroworld **seems** to be **continuous**.
Microworld is **discrete**. **This will be most important!**

Uncertainty: Fundamental **limitation** on measurement.
Some observables cannot be measured at **same** time.
——> Heisenberg Uncertainty Principle

Duality: Usually seen in so-called Wave-Particle Duality, but is a much wider effect than that.

It will be the first **challenge** to our classical intuition.

How Physics Works(or is supposed to work).

Physics **starts** with experimental data, which is then **followed** by a **hypothesis** that explains the data.

Usually we build a hypothesis slowly via **many** experiments.

Full theory can be so complex it requires approximations to make predictions.

Sometimes we use **thought** experiments; but real experiments are **required** before conclusions can be made.

Let us look at an experiment that has been both kinds of things!

—> **The Double-Slit Experiment**

First discuss —— Waves versus Particles Classically

Particle = trajectory => spatially localized + well-defined speed.

Carries energy (localized)

Obeys Newton's laws

Wave => not spatially localized => wavelengths and frequencies.

Carries energy/momentum spread over a wave front.

Exhibits non-particle-like properties;

Maxwell's equations —> **light = EM wave**.

—> Diffraction, Interference and Polarization

Waves versus Particles in the Microworld

The Microworld does not seem to have such a **clean** separation between the two ideas.

It will be very **hard** to classify objects as a particle or a wave or neither.

It will **seem** that all classical ideas have to be tossed away!

Light


Newton **thought** light = particles based on experiment + classical theory;

Young (then Maxwell) **thought** light = wave based on experiment + classical theory

Einstein finally **showed** light = particle based on experiment/theory(photoelectric effect)

If it was a particle —> see sharp shadows, but waves can bend around objects.

direction of wave motion




wave get behind barrier!

particles would exhibit sharp edge

Young said it is a wave. Shadows would be fuzzy!

Used **wave interference** property to confirm this fact.



Ultimately devised a 2-slit experiment to confirm ideas.

—-> Light is a wave **because** it exhibits **interference**.

Maxwell says light = electromagnetic radiation(wave) from his famous equations.


Wave ideas and explanations so **perfect** in classical realm that light as particle idea **fades**.

Walking the Planck

Blackbody radiation - What is it? —> Radiation observed when a body is heated!

$$\lambda = \frac{c}{f}$$

—> Blackbody Spectrum and first theory by Wien —> dashed line



$$\longrightarrow \quad \rho(f,T) = \frac{8\pi k_B T}{c^3} f^2 = \frac{8\pi k_B T}{c} \lambda^{-2}$$

How did Planck explain it? Planck used thermodynamics + intuition(always dangerous!)

Model = radiation from atoms in solid due to charge(in walls of cavity) oscillations

$$\longrightarrow \quad \rho(f,T) = \frac{Af^3}{e^{Bf/T} - 1}$$

Agrees with experiment perfectly - but it is an **empirical** formula - he could not predict A and B

—> A and B must be determined by experiment —> **not** a real **predictive** theory

Low and high frequency limits were perfect. But he could not justify law rigorously.

Mathematical trick now leads to a new physical idea and a puzzle!

During a 2nd attempt at the derivation Planck used a common math trick(a simplifying assumption) during an intermediate step in the derivation.

Usually after doing algebra and taking appropriate limits the simplifying assumption is removed. It was a standard mathematical procedure that was used everywhere(**it works!**).

Now electromagnetic theory says a body emits/absorbs energy **continuously(waves)**;

During his derivation Planck assumed(the trick) the emission/absorption process was discrete(particle), i.e., the radiation did not come out continuously(wavelike) but in discrete chunks(particle-like) governed by a new constant **h**;

He then did the math and then took the appropriate limits at the end(the standard mathematical procedure) to go back to emission being a continuous property.

When energy value was discrete in units (constant(h) x frequency) he derived that

$$\rho(f,T) = \frac{8\pi f^2}{c^3}\frac{hf}{e^{hf/k_BT}-1}$$

The mathematics trick enabled Planck to derive values for A and B as seen above.

He now had a full-fledged theory!

However his result disappears if he removes the trick!!!!!

Experimental agreement —> constant h **must** be 6.63×10^{-34} J-sec

Planck's constant

Because it is not zero ===> energy not continuous (energy is quantized)!

If make h=0, which the math trick always requires in the end, then theory fails to work!

Puzzle: Is getting the correct answer an accident due to using the math trick or is this episode telling us something about world that we would not have guessed?

That is the danger of using intuition…….

Quantized light or "Particles" of light

Einstein grabbed hold of the "energy is not continuous" idea and did this……

Einstein —> energy in EM field not spread out on wavefronts —> light is **not** a wave!!!!

Energy is localized in clumps(**quanta**!) = photon where $E = hf$ f = frequency

Only **very intense** light appears to be a wave! This can be proved!

He said Planck's trick worked in the derivation **because energy IS quantized**.

IT WAS NOT THE TRICK - NATURE ACTUALLY WORKS THAT WAY!


This quantum idea will be fundamental later to order to explain quantum measurement.

Quantized light energy or the photon idea had now explained blackbody radiation.

But any good theory should be able to explain **all** other phenomena also.

Photoelectric Effect - Einstein gets Nobel Prize

= Ejection of electrons from metal when light shines on it



Experimental results:

1. Exists a threshold frequency f_0
2. Ejection independent of light intensity unless $f < f_0$
3. If $f > f_0$ detection is instantaneous
4. Detected electron energy $= h(f - f_0)$

If light = wave, then energy proportional to **intensity** —> if bright enough should eject electrons independent of frequency, **but it does not unless above rules satisfied!!**

Also, waves cannot transmit energy so **quickly**.

Explain **bathtub-rubber duck** analogy.

The photon (quantized light) model <—> Explains every feature of the experiment!

1. $f < f_0$ —> single photon energy too small
   threshold => minimum energy to eject(called the material **work function**)
2. Energy of photon depends on frequency NOT intensity
3. Photon absorption is instantaneous (true for all quantum creation/annihilations)
4. Energy must be larger than work function of metal

Note that photon picture later also corroborated by Compton in x-ray scattering by electrons.


Problem: Light exhibits interference and particles do not interfere or so everyone thought!

Interference and Diffraction


For lots of details read **Light and Interference** see supplementary notes**.**

or Ask questions after class

Single-slit



screen pattern




Interference Pattern



Note the nodes or zeroes -> NO PHOTONS

Simple Theory - see supplementary readings for more details

Wave form is $$y(x,t) = A\sin\left(\frac{2\pi}{\lambda}(x - vt)\right)$$



—> wave transports
energy and momentum

Waves can be added/subtracted = **superposition**


$$y_1(x,t) = A\sin\left(\frac{2\pi}{\lambda_1}(x - vt)\right) \qquad y_2(x,t) = A\sin\left(\frac{2\pi}{\lambda_2}(x - vt)\right)$$


$$y(x,t) = y_1(x,t) + y_2(x,t) = A\sin\left(\frac{2\pi}{\lambda_1}(x - vt)\right) + A\sin\left(\frac{2\pi}{\lambda_2}(x - vt)\right)$$

$$y(x,0) = 2A\underbrace{\cos\left[\pi(\bar{k}_1 - \bar{k}_2)x\right]}_{\text{low frequency}}\underbrace{\sin\left[\pi(\bar{k}_1 + \bar{k}_2)x\right]}_{\text{high frequency}} \longrightarrow$$



$$\bar{k} = 1/\lambda$$






Example:
Water Waves


2-slit interference pattern(light)






The Huygens-Fresnel Principle

Each point on waveform is source of new spherical waves




Theory



Experiment

using water waves



Double slit using this principle.

Slits = new sources —— new waves superpose and interfere

Amount of interference depends on wavelength and distance from source to screen —>

$$A(\theta) = 2A_0 \cos\left(\frac{\pi d \sin\theta}{\lambda}\right)$$

Features
peaks = constructive interference
nodes = destructive interference


This is a **wave property**!!

But Einstein **proved** that light is photons = particles

Wave properties **cannot** explain photoelectric and Compton effects

The amazing experimentalists to the rescue!!

New Technology Lasers(controlled light beams) + CCD cameras(single photon detectors)



Light registers in detectors as single impacts = particle-like property


Light seems to be a particle when it needs to be and a wave when it needs to be - what it **looks like** seems to depend on the **CONTEXT** of the experiment (the question the experiment asked) - **a very important (a crucial) point as we will see**!

Much more about this later.

Now electrons are particles classically and in microworld they exhibit particle-like properties in many experiments.

Do electrons exhibit wave-like properties (in the macroworld) - do they interfere?

If I had a room with two doors in one wall and a bunch of basketballs and I threw the basketballs at the wall then with only one door open I see the pattern (distribution of the basketballs in the next room - other side of wall) as shown below:




Particles through doors(slits)(only 1 open) see either pattern I_u or I_l

Peaks are directly behind doors(slits).

This is what is observed! With enough basketballs, the distribution is **non-zero** everywhere.


If we keep both doors (slits) open at the same time, then we see the pattern left.




This is just the **sum** of the two previous distribution as we would **expect** classically since no basketballs are lost and standard classical particles **do not exhibit** any kind of interference effects.

What happens if we do the same experiment with electrons.


Davisson/Germer 1920




electron beams sent through crystal lattice,
i.e., parallel planes of atoms
replace slits!



and Hitachi 1989





Electrons do not behave as classical waves; they behave like waves with a wavelength given by

$$\lambda = \frac{h}{\sqrt{2mE}} = \frac{h}{p}$$ **deBroglie**

SAME constant h as Planck used in Blackbody radiation


Note that there are regions in the data sets where **no** electrons arrive on the screen! whereas stones(basketballs) would impact everywhere —— **Think about that for a second!**

Now reduce intensity so that we have only 1 electron in system at a time

If waves, then after few seconds should see only a weak interference pattern, but should **always** see entire pattern

Does not happen! Look at pictures. Initially pattern looks random But clearly it is not after long time.

Arrive as particles (individual impacts/flashes - quantized!) randomly but generate interference pattern after long time.



Summary

impacts = particle behavior, but pattern = wave behavior

block 1 slit = pattern like particles through 1 slit

1 electron at a time - still get final pattern -> ask question: How does each separate electron “know” what to do? **A Bad question!**

Is electron behaving as a particle/wave in same experiment or **are we misinterpreting the whole thing by trying to describe what is happening with classical ideas and macro world words!**

Maybe it has nothing to do with particles and waves which are classical ideas and is the result of something totally NEW happening.

The statement:

“Experiment seems to say that objects can be both at same time”

may simply be **last desperate attempt** to maintain classical descriptions
as we proceed to find classical physics fails.

To explain double-slit results we need a completely new theory.
We need to rethink all of our ideas about the physical world.
We need to learn a new language to understand a new world!

Part of our problem is our use of ordinary words - that is the wrong language!

Quantum Mechanics, Ordinary Language and Mathematical Language

It is **not possible** to reduce the quantum universe to everyday ways of thinking (usually called **common sense**).

In fact, in order to understand the ideas and implications of the theory we will have to **adjust** all of our ways of thinking at the most fundamental level.

Imagine, for a moment, that you are attempting to **understand** a new culture.

If you are serious about it, the **first thing** you would do is to learn the language appropriate to that culture so that you can put your new experiences in the proper context.

Understanding the universe of quantum phenomena is **much like** understanding a new culture

where the **appropriate** language is **mathematics**

and the experiences we are attempting to put into context

are **microscopic experiments**.

We have to use a **mathematical language** to describe the quantum world

since ordinary language,

which was developed to explain everyday occurrences (experiments on macroscopic objects),

will turn out to be totally **inadequate**.

There are **no macroworld models or classical analogs** that will ever give us any insight into the workings of the quantum world - so we will not make any!

Since it **makes no sense** to attempt any understanding of the nature of quantum phenomena

without first learning to speak and use the language of the quantum world,

one should **spend some time very early** on in learning the appropriate mathematics,

in particular, the subject of linear vector spaces (we will in this class).

The **adjustment** of our ways of thinking at the fundamental level

that will be needed is not simply a mathematical matter, however.

The **development** of the necessary mathematical language will not come into **conflict**
with our everyday modes of thinking in any major way.

Although, the mathematics of linear vector spaces is very elegant,
you will be able to understand the simple features we need **without much difficulty**
and **without having** your basic view of the world changed at any fundamental level.

You will be **troubled**, however, when
we apply the mathematics to physical systems that develop according to quantum rules.

You will need to **attach physical meaning to the mathematical formalism**
in ways that will **conflict** with your well-developed views
(I will call these classical views) about how the world works.

Dirac was able to join the conceptual structure with the mathematical structure.

He **invented** a mathematical language (I purposely do not use the word notation like others)
that directly embeds the philosophy of quantum mechanics
into the mathematical structure used to do calculations.

The new language **directly exhibits** what is being said about nature in quantum mechanics.

Dirac language **exposes** the internal logic of quantum mechanics
in a way that mere words cannot possibly accomplish.

It **displays** the sense, the physical meaning of the theory

in every equation one writes

without the need for further explanation or any need for always inadequate models.

It is **very important** to understand that the Dirac language

is not simply a new notation for quantum mechanics (as many physicists seem to think).

It is not merely a way of writing ———— A way of writing **expresses** a way of thinking.

Dirac language, as we will see, is a way of thinking.

Dirac language will **allow us** to use the physical ideas of quantum mechanics

to develop the appropriate mathematical language

rather than the other way around.

This **allows** the very mathematical quantum theory

to be more closely connected to experiment than any other physical theory.

Dirac language **expresses** the quantum mechanical way of thinking.

With it one **can proceed** from the philosophy of the subject to its mathematical expression

rather than the other way around.

That is the way one should study quantum mechanics.

One should **proceed from meaning** and Dirac language is perfectly suited to this task.

Meaning for us **does not reside** in mathematical symbols, however.

It **resides** somehow in the thoughts surrounding these symbols.

It is **conveyed** in words, which we will use to **assign** meaning to the symbols.

Dirac language is able to take notions expressed in words

and **replace them** with simple mathematical statements

that we are eventually able to place within a complete and logical structure

that **allows** for a fundamental understanding

of what quantum mechanics means and is saying about nature.

This task is **impossible without mathematics**.

Mathematics is the **true language** of all of physics.

Words alone only suffice for thinking about the physics of everyday objects.

These statements about the importance of understanding the mathematical language

appropriate to the physics under consideration

do not only apply to the quantum world.

It is **true**, I believe, for all areas of physics and other sciences.

One **should always learn** the appropriate language

before studying any field that relies on that language for its understanding.

So be careful.

Beginning now we shall buckle down

and take a serious look at the mathematical and physical structure of quantum mechanics.

But **tread warily**,

lest you slip back into the black abyss of classical thinking and consequent confusion.

With this warning out of the way, we **can now proceed**.

Quantum Thoughts - A Quick Overview

Quantum theory works. — 10000's of experiments - extraordinary accuracy

QM is thought to be not fully understood. — What is meaning of assumptions?
What happens during a measurement?
What is relation of QM to Reality?
(Will need to define Reality first!)
and lots more……

All physicists no matter their **interpretation** (answers to above questions) of QM make the same predictions —> all interpretations must be **equivalent**.

This class <—> Postulates + Time Evolution + Measurement + Interpretations + Reality

Some Contrasts…….

Classical View reality = everyday experiences
everyday experiences = language = reality

Classical View simultaneous measurement always OK (arbitrary accuracy)

Quantum View simultaneous measurement NOT always OK(fundamental limitations)

Einstein relativity(Classical) speed of light = maximum speed for information flow

Quantum Mechanics seemingly instantaneous influences **‘’spooky action at a distance”**

(we will clear this up this kind of nonsense!)

Quantum Mechanics measurement introduces randomness

Identical Systems

Classical - all measure **exactly** the same within experimental error.

Quantum - random results within a fixed set —— only have a **Probability**

Simple Probability Idea we will use…..

Given a set of data

n_h	h
50	150
100	160
200	170
300	180
300	190
200	200
100	210
60	220

$$probability(x) = \frac{\text{number of times value } x \text{ was measured}}{\text{total number of measurements}}$$

$$\text{probability}(190) = \frac{\text{number of times value 190 was measured}}{\text{total number of measurements}} = \frac{3}{13}(23\%)$$

Important to remember: Rules of classical probability only predictive when the number of measurements gets very large.

—> This is an intuitive and comfortable definition.

But, not necessarily correct!

Back to 2-slit experiment

Only one slit(B) open => photons on screen centered directly behind B

Only one slit(A) open => photons on screen centered directly behind A

Both slits open = **no** photons in some spots where photons appeared when only one slit open

Think about that!
More Ways
=
Less Photons!

Experiment also work with electrons.

Experiment works no matter intensity of beam (even 1 particle at a time)

Reverse the thinking. 2 slits open = screen pattern with nodes(zero photons)

Close one slit = more particles make it to node region! Same for electrons!

Makes no classical sense - i..e., it does not happen if we were using marbles!!

A poorly worded question — How does a single particle ‘’know” 1 or 2 slits open?

Quantum particles(are quantized) cannot be split
—> so it cannot be one piece interfering with other piece

If we try to detect which slit —> experiment changed —> pattern is destroyed

Can use wave picture and interference to explain things, but Einstein already showed that photons were particles (actually showed they were quanta) and only look like waves when the beams are very intense - wave idea fails entirely when we have weak beams as we will see.

Clearly we need new way of thinking in Microworld.

Scaling down from macroscopic to microscopic radically alters particle distributions (they must be somewhere) and no interference effects becomes interference effects

Never see two independent things conspiring to cancel each in macroscopic world!

But happens often in microworld, i.e., the nodes in the interference pattern.

How will quantum theory want us to think about all of this so that we have some hope of understanding what is happening?

As we will see…….

For a photon ‘’going between source and screen” (whatever those ordinary words mean)
—> photon state CANNOT be described as

(1) having gone through slit B or
(2) having gone through slit A or

(3) having gone through both slits simultaneously or

(4)having gone through neither slit

This exhausts **all logical possibilities** in the macroscopic world (using **ordinary language**).


So **Ordinary Language Fails**.

In order to give an explanation classically we will only be adding new words/new definitions to old language and not explaining anything!

As we will see **probabilities** will win out over **exact trajectories**.

Let us try to measure the position of an electron.


Measurement Process



— —> Note that we do not observe the microword DIRECTLY!

Where in sequence of instruments (measurements) is the information which makes the position meter dial point to definite value **replacing** the quantum uncertainties of electron with the definiteness of pointer?

i.e.,



Where is discontinuity?

Is there a discontinuity?

The saga of an electron passing through a wall.

Suppose we shoot a beam of electrons at a thick metal wall.

If we were shooting bullets instead, they would bounce off(that is what classical particles do!)

But, with electrons, **some** will appear on the other side of the wall.

How did they get there?

Obvious answer (to a classical mind) is they must have passed through the wall.

Therefore, I should be able to set up a detector inside the wall and detect them as they pass through - I mean, they had to pass through to get to the other side - CORRECT?

Experimental result: With the detector in place checking, no electrons appear on the other side!

No detector - electrons appear; detector in place - no electrons appear!

Seems like electrons just disappear on one side and appear on the other side without passing through the wall (since no measurement shows that if they are detected inside the wall that they get to other side!!)

Here and there and NEVER in between. **That is the way QM works!**

How do photons each **know** what to do? (this will turn out to be a **poor/incorrect question**).

Only satisfying explanation will be **probability**.

Each photon has probability P(x) to arrive at position x on the screen(detector).

Each photon **independently** uses P(x) to choose impact point.

Each photon **independently** generates **part** of interference pattern.

2 slits = interference pattern

If observe to see which slit = information about which slit particle then

photon either went through that slit or it did not and

we “mess up” or ‘’change” particle to a known slit state => pattern changes to 1 slit pattern

Some Last Thoughts Before Developing a Theory

We will find that:

We must resort to probabilities! Probabilistic information = maximum information!

Measurement disturbs our information(probability)!

In microworld we will find that probability is everything.

Until a measurement changes information we will have to say that there is no definite value for a state —> everything is simply set of probabilities before measurement!

Particle is not somewhere unless measured to be there!

In between measurements it is only probabilities!

This statement will have measurable (it can be proven) effects.

Read supplementary lecture notes ——- EinsteinBohrGod

Ask questions after class

Read supplementary lecture notes ——- Tale of Two Gloves
It illustrates QM phenomena via a story

Ask questions after class

Equivalent phenomena in the microword really behave this way

No explanations are given yet - that comes later in the class

As we will see, quantum theory **arrives** at what it deems to be an acceptable interpretation of these puzzles by **insisting** that we stick to practicalities.

It is no good, and indeed very dangerous, **to speculate** about what **seems** to happen in such any situation.

Stick to what actually occurs, and can be recorded and verified, and you will be **correct**.

For example, if you **cannot** actually send an instantaneous message of your own devising, then it is **meaningless** to guess at what might or might not have been secretly going on between the two gloves in the story.

You **might think** that if we do not understand all aspects of what is going on

then we will not be able to do anything useful with this quantum theory stuff.

It **turns out** just the opposite!

Probabilities will be sufficient!

As will **see later** - Einstein took **one side** in this dispute - he was **WRONG**

Classical measurement => gain knowledge pre-existing state

Quantum measurement => impart specific value to state

We will approach QM from real theorist's viewpoint.

You have **never seen** theoretical physicist(me) in action.

I might do things that seem off the wall(no reason).

It is **important** to remember that theorist has **one goal** —> to understand universe.

No assumption is considered too crazy to be included.

Invention of new ways of thinking will be commonplace.

Strange interpretations will be all over the place.

What is correct? .... Only that which **agrees** with measurements!

We proceed **along lines** set out by Paul Dirac in 1920s

modified by much hindsight from intervening 100 years of discovery.

Now on to Lecture 2

Start of Part #2

No theory yet - This story I will now tell will help push us in the correct direction!

A Simple Introduction to Quantum Ideas via Experiments

I am going to tell you a story about a world of electrons that have 2 properties

Color and Hardness

In this imagined Microworld:

Electrons have 2 properties ==> color and hardness

Note that In the Real world:

Electrons also have 2 properties ==> z- and x-spin (orthogonal directions)

and

Photons also have 2 properties => horizontal and vertical polarization

The Strange World of Color and Hardness

Known experimental facts:

Color has **two** possible values = green or magenta.


This means that **nothing else** has ever been **measured** for electron color.


Similarly, for hardness property.

Electrons are either hard or soft (**only** possibilities).

Nothing else ever **measured**.

Color/Hardness Boxes (have 3 apertures)







Electrons enter boxes through **left** aperture and exit through one of other two apertures.

These boxes represent the same physics as real world devices but without complications.

[Examples : Stern-Gerlach devices and Polarizers as we will see later.]






Boxes **separate** electrons in physical space (put them onto **different paths**)
according to the value of color or hardness as shown.

Thus, in either case, one can distinguish the electrons **after** they pass through a box
by their final position(path) in real physical space, i.e., we have **separate** beams at end.

AN EXPERIMENTAL PROPERTY

Measurements with hardness or color boxes are **repeatable**.



The 1st color box determines the electron color and we choose the magenta beam.

Then, without tampering(must define this concept) between measurements the magenta beam is subsequently (immediately=> $\Delta t=0$) sent into left aperture of another color box.

The electron will (with probability = 1) emerge from 2nd box through magenta aperture.

This **same** property of repeatability holds for green electrons and color boxes

and for hard or soft electrons with hardness boxes.


It is **simply** a statement of the way the real microworld and its measuring devices work in the laboratory.

So our representation of the real world should behave that way also.

Suppose we suspect the possibility that color and hardness properties are related in some way.

We must investigate => check for **correlations** (or **relationships**) between the measured values of hardness and color properties.

The boxes allow for checking whether such correlations exist.



No such correlations exist,

i.e., of any large collection of, say, magenta electrons,

all fed into left aperture of hardness box, **precisely** 1/2 emerge through hard aperture,

and **precisely** 1/2 emerge through soft aperture - one cannot predict which goes where!


Same result occurs for green electrons and similar results hold if we send hard or soft electrons into color box - all results are exactly 50% —> **there are no correlations**.

Color (hardness) of electron apparently gives no information about hardness (color).

This usually means that no correlations seem to exist.

Now, suppose we set up a sequence of **three** boxes.


Color box, then hardness box and finally color box.



Suppose an electron(1 of 100) that emerges from magenta aperture of 1st color box

then goes into left aperture of hardness box

(no tampering => no other measurements or no time interval).



Then suppose an electron that emerges(50 out of 100 do) from hardness box through soft aperture (1/2 of input 100 magenta electrons)

and are sent into left aperture of last color box

(again no measurements or time intervals allowed).

Presumably, the electron(1 out of 50) that enters that last box

is known to be **BOTH** magenta **and** soft,

which were results of two previous measurements just made on it!

If so, then we expect the electrons(50) to emerge from magenta aperture(probability = 1),

confirming the result of 1st measurement.

Any **reputable** classical physicist would say at this point

that electrons entering last color box are known to be **magenta/soft**.

They have properties and we are just measuring again what they already have!!!!

Repeating, in the classical world, particles have **objective reality** - have **real properties**


and we just **measure to find out** values of properties.

We just used 2 experiments to determine these properties!

PROBLEM: **this is not what happens in the real microworld in this experiment!**

Precisely half of electrons entering last box emerge from magenta aperture

and precisely half emerge from green aperture!!



Therein lies the fundamental puzzle of the quantum world.


Any theory we develop must account for this result.

In fact, if 1st two measurements give magenta/soft, magenta/hard, green/soft or green/hard

which represents all possible cases, then,

when any of these beams sent into last box,

precisely half emerge from each aperture!!



It seems as if presence of hardness box between two color boxes

constitutes some sort of color measurement or color tampering and vice versa.


Hardness box **seems to be changing** half of magenta electrons into green electrons.

As we will see, this will turn out to be an **incorrect way** of thinking about the results of the experiment!

Hardness box must be the blame since if it were not there,

then the last box would only see magenta electrons

(a **different** experiment that corresponds to repeatability)



This all seems so weird (non-classical)

—> we must question (challenge) all features of experiment before we accept story as truth.

Perhaps hardness box is poorly built (not enough $$$$ from Congress in our research grant).

Maybe measuring hardness correctly disrupts color —> bad design.

Two fundamental questions arise:

1. Can hardness boxes be built that will measure hardness without disrupting color?

2. In case of poorly built apparatus, it seems as if half of electrons change color….

   What is it that determines which electrons have color changed and which do not change?

Question 2 can be answered by checking correlations —> **none exist**.

Those electrons that have color changed by hardness box

and those electrons whose color is not changed by hardness box

do not differ from one another in any **measurable** way.

So 2nd question has no answer that we can figure out from measurements.

If we believe, as I do, that **I can only know properties that I can measure**,

then it means there is NO answer to this question,

i.e., there is no property of electrons that determines which electrons get color changed.


This is completely counter to classical notion of cause/effect and objective reality!!!

Question 1: no matter how we build hardness box (remember a device qualifies as a hardness box if it can separate electrons by hardness value) **all disrupt color measurements** - all change half of entering electrons.

Any hardness (color) measurement seems to **randomize** next color (hardness) measurement, i.e., make it 50% green/ 50% magenta.

What about building a color AND hardness box?

This would need 5 apertures as shown.



Has to consist of hardness boxes and color boxes or their equivalents

because we must be measuring hardness and color in some way.

But as we have already seen, whichever box the electrons pass through **last**

provides reliable information ONLY about that measured quantity

and the other quantity is **randomized** (i.e., half/half).


No one has succeeded in building device

which can simultaneously measure both color and hardness.

Seems fundamentally beyond our means no matter how clever we are.

We will find the reason later!

Probing Deeper



We now build a more complicated experimental setup involving two paths as shown.

We will see this device often later - it is called a Mach-Zehnder Interferometer.

The mirrors are just reflectors which do not affect color/hardness values.


That means it has been experimentally checked to be true.

In this device, the hard and soft electrons follow **different paths in physical space**

and eventually the electrons are recombined into single beam again in black box at end.

All the black box does is recombine two beams into one beam by direction changes (like light beams with mirrors)

Again, this is done without changing hardness or color values (checked experimentally).



So if we start

with a mixed (hard + soft) beam of electrons,

entering the first hardness box


then we end up

with a mixed (hard + soft) beam (unchanged) at the end.

The effectiveness of device has been checked separately for both hard and soft electrons and it works.

Start with hard —> hard at end and they **went** on hard path!

Start with soft —> soft at end and they **went** on the soft path!



This is very important so I repeat:

If hard or soft electrons are sent in separately,

they simply travel along different paths and end up in same place

with their hardness property **unchanged**.


They go on their separate paths and emerged unchanged at end

Now some experiments can be done with apparatus.

All of these experiments have **actually** been done in real laboratories with equivalent setups.

Listen carefully now to see where your classical mind is misleading you.

magenta




Send magenta electrons into 1st hardness box.

At end (after recombination of beams)

add a final hardness box

and thus we are measuring hardness at that point

i.e., after the “magenta” electrons have **“passed”** through the apparatus.



magenta

Analysis:

For entering magenta electrons,


50% take h route and 50% take s route

at the end(recombined beam) 50% are hard electrons

(remember nothing bothers hardness along routes)

and 50% are soft electrons. —-> Experimentally verified!

Change experiment



Send hard electrons into hardness box.

At end add color box (**replace** final hardness box with a color box) and **measure color**.

Analysis:

All hard electrons follow h route.


When we measure color of hard electron it is 50-50 green-magenta.

Similarly for soft electrons.

Therefore end up with 50-50 green/magenta coming out of color box.

These experimental outcomes are what would **expect** from earlier results and **do happen** exactly as described above in real world experiments. So **no** problems yet!!!

Now again send magenta electrons into hardness box, but at the end add a color box and measure COLOR.



magenta

What do you expect? The true classical physicist would give the following argument.

Since, for magenta electrons,

50% are hard and 50% are soft

(what happens to magenta electrons when sent into hardness box —> are randomized)

and each kind of electron takes the appropriate h and s routes,

at end, 50% of h (those on h route) electrons or 25% of total are magenta

and 50% of s (those on s route) electrons or 25% of total are magenta.

Thus, for 100% magenta sent in,

classical reasoning says that only 50% are magenta at end. A prediction!

Seems like a valid conclusion(prediction) since hardness boxes(1st box) **supposedly** randomize color.

Problem:

This last part of story, which a classical mind desperately wants to accept as correct, is **false**.

When we actually do experiment, **all(100%)** of electrons at end are magenta!!!

Very odd!

Hard to imagine (**classically**) what can possibly be going on in this system.


Of course, maybe it is just that our **classically oriented minds**

cannot imagine what is happening and need to be **retrained**!!!

Let us **try another experiment** in hopes of understanding what is going on.

Rig up small, movable, electron-stopping wall that can be inserted in and out of route s.

When the wall **out**, we just have the earlier apparatus.



wall not inserted

wall inserted

When wall is **in**, **only** those electrons moving along h route

get through apparatus to the beam recombination device.

The electrons on the s route are blocked.

Analysis: First, there are 50% less electrons reaching beam recombination device.

When wall out all(100%) of electrons that get to beam recombination device

are magenta at end (earlier experimental result).

That means (**or seems to mean**) that all electrons that take s route

(**or we think are taking s route**) end up magenta

However, we did not check this!!!

and all that take h route (**or we think are taking h route**) end up magenta

However, we did not check this!!!

at beam recombination device when **no wall is inserted**.

This means that with wall inserted, we should end up with 50%

(1/2 lost impacting inserted wall) electrons at beam recombination device.

Based on the earlier experiment results, they should **all** be magenta

since an inserted wall on soft path **cannot possibly affect** electrons on h path

and they **all** came out magenta in the previous experiment!

This is an assumption called **locality**, which means the following:

We can separate the hard and soft paths by a very large distance

and therefore make sure that nothing we do on soft path(like inserting wall)

can affect electrons on hard path during time they are on hard path.

This is possible because the maximum speed of information transfer is

speed of light from the study of relativity.

What is actual result?

Again we are wrong in our classical way we are analyzing things. The result is **50/50 again**.

h route beam(**if there really is one**) seems to end up randomized(50-50 green-magenta)

and 50% of 50% is 25% 25 magenta and 25 green at end!

If we insert a wall in h route, **same thing happens**. Still end up only with 25% magenta!

Clearly, we have **real** trouble with our classical way of reasoning!

Seems if check (**do measurement**)

to **see which path** electrons are passing through device

(i.e., if we check to see whether the magenta electron
is passing through apparatus as a hard or a soft electron)

we get 25% magenta (50% change to green) at end
and if we DO NOT check we get 100% magenta (0% change to green).

Our classical minds are **spinning** at this point!!

We now turn to a quantum system **containing only one particle at any instant of time**
instead of using beams with many particles at any instant of time.

This was not possible during the early formative day of quantum mechanics.

It only became possible in the 1980s.

Now consider a **single magenta electron** passing through apparatus when wall is out.

Can it have taken h route?

No, since all such electrons are hard and hard electrons are 50-50 green-magenta
(and need 100% magenta).

If on hard path, then they are hard - that is a measurement.

Can it have taken s route?

No, since all such electrons are soft and soft electrons are 50-50 green-magenta
(and need 100% magenta).

If on soft path, then are soft - it is measurement.

Can it somehow have taken **BOTH** routes at same time?

If **look** (**measure**), then half the time find electron on h route

and half the time find electron on s route,

but never find two electrons in apparatus,

or two halves of single, split electron,

one-half on each route, or anything like that.

No experimental way in which electron seems to be taking **both** routes simultaneously.

Therefore, as physicists, must **rule out** this possibility.

This eventually leads to the hypothesis for the meaning of the the word "quantum" as we will see.

Can it have taken **neither** route (got there some other way)?

Certainly not.

If we put walls in **both** routes, then NOTHING gets through at all.

Thus, the results **had** to have something to do with the box and the paths.

Summarize the dilemma for classical view.

Electrons passing through apparatus, based on measurements

do not take route h

and do not take route s

and do not take both routes at same time

and do not take neither of routes,

i.e., electrons have **zero probability** for doing any of these things (**from experiment**).

Problem is that those 4 possibilities are simply **all** of the logical possibilities

we have any notion of how to entertain using the **everyday language** of classical physics!

What can these electrons be doing?

They **must** be doing something which has simply never been thought of before in classical physics.

Electrons (in the microworld) seem to have modes of being, or modes of moving,

available to them which are unlike anything

we can think about using **words** derived from everyday ideas and classical physics.

The name of new mode (a name for something do not understand at the moment) is

SUPERPOSITION.

In the end all we can say about initially magenta electron, which is passing through our apparatus with the wall out,

Prepare yourself for this!

it is **NOT** on h path and **NOT** on s path and **NOT** on both paths and **NOT** on neither,

it is in a **SUPERPOSITION** of being on h and being on s.

What this last statement means, other than none of above, we do not know at this time;

it requires a theory to fill in the gaps and we will do that.

That is what one does in this (or any) quantum theory class.

Final and most confusing experiment.

New experimental box = **do-nothing box** - It has just 2 apertures.

An electron goes into one aperture and emerges from other aperture

with **ALL measurable properties** (color, hardness, energy, momentum, whatever)

UNCHANGED. It is Experimentally verified!

Box does not measure anything.

If we measure the properties of electrons before entering box and then after passing through box,

no measured properties are changed.

Also the time it takes for electron to get through box

is **identical** to time it would have taken if box were not there.


So nothing mysterious **seems** to be delaying it/messing around with it while in box.

Such a **do-nothing box** can actually be constructed in the laboratory.

Experimentally if we build such a do-nothing box, then,

when inserted into **either** one of two paths (h or s),

it will **change all** electrons that were **magenta** at end into **green** electrons at **end**.



box not inserted box inserted

When the box is removed, they all go **back** to all being magenta.

So inserting box into only **one** path changes the color of electrons passing through the entire apparatus.

What is going on here?

Do-nothing boxes do not change any measurable property of electrons

that pass through it

AND of course,

do-nothing boxes do not change any measurable property of electrons

that DO NOT pass through.

That would not make any **sense** at all.

So once again our only explanation will go like.....

It is not possible for the electron to have passed through the do-nothing box

since we already said that cannot change anything.

It is not possible for the electron to have passed outside the box

since the box certainly does not have a chance to bother anything

that does not even pass through it

(even though it would not do anything anyway).

It is not possible that the electron passes

both simultaneously inside and outside of the box or neither as before.

Only answer is that the electron passes through our apparatus

in a **superposition** of passing through do-nothing box

and not passing through do-nothing box

and this must **cause** the color change somehow.

This theory has got to be really neat when we finally figure out what it is we are saying about the world.

Now onto mathematics to learn/ understand the appropriate language of physics, develop a theory and then we will be able figure out what is going on.

A pattern exists……

Mathematics language <—> Relationships

Relationships <—> Correlations

Correlations <—> Quantum Mechanics


Quantum Mechanics <—> Mathematics language

The reality of this pattern will become clear as we proceed through the class.

The Mathematics Of Quantum Mechanics - Part 1

We must learn some mathematics BECAUSE it is language of QM.
We will not be able to explain and use the ideas of QM using WORDS!!


First, some High School level Stuff…

Vector =  directed line segment


if in a plane—-> 2 numbers (a,b) = number of dimensions

or Vector —> 3 numbers (a,b,c)= number of dimensions in ordinary space


Multiplication by a scalar(a number)




Given two vectors



one can add/subtract them with the geometric rules shown below



Sum



Difference


call the “tail to tip” rule

Now we generalize —> extend idea to a general linear combination(sum) of vectors

$$\vec{C} = \alpha\vec{A} + \beta\vec{B}$$ it is still a vector!

using $\alpha = 2$

$\beta = 0.5$



adding parts => total

I am drawing pictures in the real 3-dimensional world here!

Here, I note that all coefficients α, β are real numbers at this time.

These ideas and methods work perfectly for all macroworld vectors(high school math).

But we must generalize further in order to deal with microworld vectors(this class).

Let us see how……..

The Space of Physics

vector = 3 numbers (3 = dimension of our world)


Location in space (relative to an origin) = 3 numbers

use axis labels $(x = x_1, y = y_1, z = z_1)$

—-> can define a position or radius vector

$$\vec{r} = (x, y, z) = (x_1, x_2, x_3) = (a, b, c)$$

as shown in diagram




Now to add more mathematics to increase the descriptive power:


First - define 3 special vectors (called a **basis**) that will be used to construct all other vectors - several different label sets….

$$\hat{e}_1, \hat{e}_2, \hat{e}_3 \text{ or } \hat{i}, \hat{j}, \hat{k} \text{ or } \hat{x}, \hat{y}, \hat{z}$$

They are unit vectors (length=1) and define a Right-Handed coordinate system as shown.



By using vector addition rules we have (could prove this using tip-to-tail rules)



$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k} = x_1\hat{e}_1 + x_2\hat{e}_2 + x_3\hat{e}_3$$

$$= x\hat{x} + y\hat{y} + z\hat{z}$$

In words, x_1=x=component of $\vec{r}$ in 1-direction(x-direction), etc

$x_1\hat{e}_1$ **is a vector in 1-direction of length x_1**

$\vec{r}$ **is vector sum of three vectors** $x_1\hat{e}_1$, $x_2\hat{e}_2$, $x_3\hat{e}_3$ **draw it yourself**

$\hat{e}_1, \hat{e}_2, \hat{e}_3$ or $\hat{i}, \hat{j}, \hat{k}$ or $\hat{x}, \hat{y}, \hat{z}$ **called a basis set**

Important fact —> any vector can be written in terms of a basis

vector length $r = |\vec{r}| = \sqrt{x^2 + y^2 + z^2}$

using Pythagorean theorem

Go to 2 dimensions (for drawing **simplicity)**
Everything I do generalizes to higher dimensions easily.

$$\hat{e}_x = \text{ unit(length } = 1) \text{ vector in } x - \text{direction}$$

$$\hat{e}_y = \text{ unit(length } = 1) \text{ vector in } y - \text{direction}$$


multiply by a scalar


$$A_x\hat{e}_x = \text{ vector of length } A_x \text{ in the } x - \text{direction}$$

$$A_y\hat{e}_y = \text{ vector of length } A_y \text{ in the } y - \text{direction}$$

$$\vec{A} = A_x\hat{e}_x + A_y\hat{e}_y$$

by addition rule for vectors as shown






Now from the diagram

$$A_x = A\cos\alpha \quad , \quad A_y = A\sin\alpha$$

trigonometry —> **components**



For two vectors we now have

$$\vec{A} = A_x\hat{e}_x + A_y\hat{e}_y \quad , \quad \vec{B} = B_x\hat{e}_x + B_y\hat{e}_y$$

$$\vec{A} + \vec{B} = (A_x + B_x)\,\hat{e}_x + (A_y + B_y)\,\hat{e}_y$$

$$\vec{A} - \vec{B} = (A_x - B_x)\,\hat{e}_x + (A_y - B_y)\,\hat{e}_y$$

thus the old tip-to-tail idea is now **eliminated** in favor of adding and subtracting components; tip-to-tail idea cannot be extended to more complicated situations such as in microword but the new idea **can**!!

More Generalization - Further thinking about vectors

old $\vec{V} = v_1\hat{e}_1 + v_2\hat{e}_2$ or $|V\rangle = (v_1, v_2)$

just a change of "notation" —> removes unit vectors

$$\longrightarrow \quad |V\rangle = (v_1, v_2) \quad \longrightarrow \quad |A\rangle = (A_x, A_y)$$

Start of Dirac language

new symbol $|\ \rangle$ —> Dirac **Ket** vector - it is the **first** element in a new language

Your only job is to learn the new terminology(language);
I will do all the algebraic manipulations in class.

Now using new language to replace old equations……

addition

$$\text{If } |V_1\rangle = (7, -2) \text{ and } |V_2\rangle = (-5, 3)$$

$$\text{then the sum } |V\rangle = |V_1\rangle + |V_2\rangle = (7 + (-5), (-2) + 3) = (2, 1)$$

we are still adding components in same way as before!

difference

$$\text{If } |V_1\rangle = (7, -2) \text{ and } |V_2\rangle = (-5, 3)$$

$$\text{then the difference } |V\rangle = |V_1\rangle - |V_2\rangle = (7 - (-5), (-2) - 3) = (12, -5)$$

we are still subtracting components in same way as before!

length

$$\text{If } |V\rangle = (v_1, v_2) \text{ then length } V = \sqrt{v_1^2 + v_2^2}$$

Now we have the language elements that will allow us to extend ideas so they are useful in the microword

We now define a **new operation** and the **generalized vector length**

$$\text{If } |V\rangle = (v_1, v_2) \quad , \quad |U\rangle = (u_1, u_2)\,,$$

$$\text{then } \langle V\,|\,U\rangle = v_1 u_1 + v_2 u_2 \qquad \textbf{definition}$$

Here we are using vectors with real-valued components —> called **Euclidean vectors**

the symbol $\langle \cdots | \cdots \rangle$

—> Dirac **bra**-c-**ket** or **braket** symbol —> another element in Dirac language

Then we have

$$\langle V\,|\,V\rangle = v_1 v_1 + v_2 v_2 = v_1^2 + v_2^2 = (\text{length})^2$$

$$\text{length } = \sqrt{\langle V\,|\,V\rangle} = \sqrt{v_1^2 + v_2^2}$$

In the world we are in at this moment with real-valued component Euclidean vectors all vector lengths are non-negative numbers, i.e.,

$$\langle V\,|\,V\rangle \geq 0$$

One of the reasons for developing these new ideas is that the formalism now easily extends to more dimensions……

3 dimensions

$$|V\rangle = (v_1, v_2, v_3) \quad , \quad |U\rangle = (u_1, u_2, u_3)$$

$$\langle V \,|\, V\rangle = v_1 v_1 + v_2 v_2 + v_3 v_3 \qquad \langle U \,|\, U\rangle = u_1 u_1 + u_2 u_2 + u_3 u_3$$

$$\langle V \,|\, U\rangle = v_1 u_1 + v_2 u_2 + v_3 u_3$$

Definition:

summation symbol

$$\sum_{i=n_1}^{n_2}$$

$$\sum_{j=1}^{5} A_j = A_1 + A_2 + A_3 + A_4 + A_5$$

$$\sum_{i=3}^{7} A_i B^i = A_3 B^3 + A_4 B^4 + A_5 B^5 + A_6 B^6 + A_7 B^7$$

and so on ……

then we can write

$$\langle V \,|\, U\rangle = v_1 u_1 + v_2 u_2 + v_3 u_3 = \sum_{k=1}^{3} v_k u_k$$

$$\text{length} = \sqrt{\langle V \,|\, V\rangle} = \sqrt{v_1 v_1 + v_2 v_2 + v_3 v_3} = \sqrt{\sum_{k=1}^{3} v_k^2}$$

More than 3 dimensions (we are not in Kansas anymore, Toto)

I can no longer easily visualize(draw on paper) so I stop trying!!

$$|V\rangle = (v_1, v_2, ..., v_n) \quad , \quad |U\rangle = (u_1, u_2, ..., u_n)$$

$$\langle V \,|\, U\rangle = v_1u_1 + v_2u_2 + \cdots + v_nu_n = \sum_{k=1}^{n} v_k u_k$$ **extended definition**

$$\text{length } = \sqrt{\langle V \,|\, V\rangle} = \sqrt{v_1v_1 + v_2v_2 + \cdots + v_nv_n} = \sqrt{\sum_{k=1}^{n} v_k^2}$$

Dirac Braket = scalar or inner product(or “dot” product)

Now the next step is to generalize the idea of "components"

Up to now "components" = "real" numbers and vectors were "Euclidean vectors"

Generalization —-> complex components i.e., components are complex numbers

Digression on simple properties of complex numbers

$i^2 = -1 \ , \ i^3 = -i \ , \ i^4 = +1 \ , \ i^5 = i \ , \ i^6 = -1 \ , \quad \text{and so on}$ **definition** of i

$z = a + bi$ **definition** of complex number a = **real** part b= **imaginary** part

$$(7 + 4i) + (-2 + 9i) = 5 + 13i$$

definition of addition(or subtraction)

add real parts and imaginary parts separately

$$(7 + 4i)(-2 + 9i) = (7)(-2) + (7)(9i) + (4i)(-2) + (4i)(9i)$$
$$= -14 + 63i - 8i - 36 = -50 + 55i$$

using $i^2 = -1$

definition of multiplication

definition complex conjugate $z^* = a - bi$

definition absolute value

$$|z|^2 = z^* z = (a - bi)(a + bi) = a^2 + b^2 \qquad |z| = \sqrt{a^2 + b^2}$$

Note that if z is real(imaginary part = 0) then z* = z

In this generalized world the **braket** also has an extended definition

$$\langle V \,|\, U \rangle = v_1^* u_1 + v_2^* u_2 + \cdots + v_n^* v_n = \sum_{k=1}^{n} v_k^* u_k$$

If components are real(Euclidean vectors), the $v_i^* = v_i$ and we have the same formula as earlier

$$\langle V \,|\, U \rangle = v_1 u_1 + v_2 u_2 + \cdots + v_n u_n = \sum_{k=1}^{n} v_k u_k$$

now $$\text{length} = \sqrt{\langle V \,|\, V \rangle} = \sqrt{\sum_{k=1}^{n} v_k^* v_k} = \sqrt{\sum_{k=1}^{n} |v_k|^2}$$

Vectors in QM = Euclidean vectors with complex components - they live in a mathematical place called a "Hilbert space"

Now to extend all these idea to include Basis Vectors

just 2 dimensions for simplicity

$$|1\rangle = (1,0) \leftrightarrow \hat{e}_x \quad , \quad |2\rangle = (0,1) \leftrightarrow \hat{e}_y$$

new representation for old unit basis vectors

$$\hat{e}_1 = \hat{e}_x \quad \hat{e}_2 = \hat{e}_y \quad \hat{e}_3 = \hat{e}_z$$

3 dimensions

$$|1\rangle = (1,0,0) \quad , \quad |2\rangle = (0,1,0) \quad , \quad |3\rangle = (0,0,1)$$

called n-tuples

Clearly length = 1 (unit vectors) and they are perpendicular or orthogonal(by definition) and as we will prove shortly

All the same……with changed mathematics language

Some examples:

$$|V\rangle = v_1\hat{e}_1 + v_2\hat{e}_2 + v_3\hat{e}_3$$

$$|V\rangle = (v_1, v_2, v_3)$$

$$|V\rangle = v_1\,|1\rangle + v_2\,|2\rangle + v_3\,|3\rangle$$

$$|1\rangle = (1, 0, 0)$$

$$\langle 1\,|\,V\rangle = v_1\,\langle 1\,|\,1\rangle + v_2\,\langle 1\,|\,2\rangle + v_3\,\langle 1\,|\,3\rangle$$

$$= v_1(1) + v_2(0) + v_3(0) = v_1$$

Definition: Vector Space = collection of vectors such that

if add any two vectors together => another vector in collection;

and have **a scalar product(bracket)** defined


—> Hilbert space in microworld

Can write **any** vector in terms of **basis <—> orthonormal set**, i.e.,

$$|v_1, v_2\rangle = v_1(1,0) + v_2(0,1) = (v_1, 0) + (0, v_2) = (v_1, v_2)$$

$$|v_1, v_2\rangle = v_1(1,0) + v_2(0,1) = v_1 |1\rangle + v_2 |2\rangle$$

$$\vec{v} = (v_1, v_2) = v_1 \hat{e}_1 + v_2 \hat{e}_2 \quad \textbf{“old way”}$$



no longer a valid picture when components are complex; just suggestive!!

v_1 = component of vector in 1-direction, etc

Similarly for 3 dimensions

$$|U\rangle = (u_1, u_2, u_3) = u_1(1,0,0) + u_2(0,1,0) + u_3(0,0,1) = u_1 |1\rangle + u_2 |2\rangle + u_3 |3\rangle$$

and so on

Using earlier rule for evaluating the scalar product (sum of component products)

$$\langle 1 \,|\, 1 \rangle = (1,0) \cdot (1,0) = (1)(1) + (0)(0) = 1 \qquad \langle 1 \,|\, 2 \rangle = (1,0) \cdot (0,1) = (1)(0) + (0)(1) = 0$$

$$\langle 2 \,|\, 1 \rangle = (0,1) \cdot (1,0) = (0)(1) + (1)(0) = 0 \qquad \langle 2 \,|\, 2 \rangle = (0,1) \cdot (0,1) = (0)(0) + (1)(1) = 1$$

$$|V\rangle = v_1 \,|1\rangle + v_2 \,|2\rangle$$

=1 **=0** **=0** **=1**

$$\langle 1 \,|\, V \rangle = v_1 \,\langle 1 \,|\, 1 \rangle + v_2 \,\langle 1 \,|\, 2 \rangle \qquad \langle 2 \,|\, V \rangle = v_1 \,\langle 2 \,|\, 1 \rangle + v_2 \,\langle 2 \,|\, 2 \rangle$$

$$\longrightarrow \quad \langle 1 \,|\, V \rangle = v_1 \qquad \langle 2 \,|\, V \rangle = v_2$$

in new language

—> The component of vector in particular direction(along a basis vector) is given by the scalar product(bracket) of vector with corresponding basis vector. New Definition of component!

Remember component = projection on the axis in old real world!

But now, in Hilbert space, it is a “braket” - no more pictures!!

Thus, we can always write

$$|V\rangle = \underset{v_1}{\langle 1 \,|\, V \rangle} \,|1\rangle + \underset{v_2}{\langle 2 \,|\, V \rangle} \,|2\rangle$$

general definition in new language

Orthogonality

First way - remember High School - using old language

$$\vec{V} = (7, 4) \qquad \hat{x} = (1, 0) \quad , \quad \hat{y} = (0, 1)$$

$$\longrightarrow \quad \vec{V} = 7\hat{x} + 4\hat{y} \qquad \textbf{using old notation}$$

Define old “dot” product:


$$\vec{A} \cdot \vec{B} = A_x B_x + A_y B_y \ = \langle A \,|\, B \rangle$$

components

$$7 = v_x = \hat{x} \cdot \vec{V} \quad , \quad 4 = v_y = \hat{y} \cdot \vec{V}$$

$$\sqrt{v_x^2 + v_y^2} = \sqrt{65} \quad \text{length}$$

Notice that every equation is identical to those in Dirac language. We only went to new language because these older ideas cannot be extended into microworld!



Now we can always write (in peculiar way - kind of stuff mathematicians and theoretical physicists do all the time!)

$$7 = v_x = \hat{x} \cdot \vec{V} = length(\hat{x}) \times length(\vec{V}) \times \frac{v_x}{length(\vec{V})}$$


length =1

whole mess just multiplies v_x by 1

Can you see?

What are you really doing when you carry out algebra? You are either multiplying by 1 or adding 0 — otherwise you will change the expression you are working on!

Now $$\frac{v_x}{length(\vec{V})} = \text{ cosine of angle between } \vec{V} \text{ and } \hat{x}$$ **—> definition of cosine**



so $$\vec{A} \cdot \vec{B} = length(\vec{A}) \times length(\vec{B}) \times \cos(angle\, between\, \vec{A}\, and\, \vec{B})$$

generalize $$\vec{A} \cdot \vec{B} = AB\cos\theta$$ **0 if perpendicular**

Important special cases:

$$\hat{x} \cdot \hat{x} = length(\hat{x}) \times length(\hat{x}) \cos(0^\circ) = 1 \quad \hat{x} \cdot \hat{y} = length(\hat{x}) \times length(\hat{y}) \cos(90^\circ) = 0$$

—> Basis vectors are orthonormal! As I have been assuming!

= orthogonal + normalized to 1(length=1) —> orthonormal

Generalize further (add new notation) by defining in a vector space

$$\langle A \,|\, B \rangle = AB \cos\left(\theta_{AB}\right)$$ orthonormality clear!

no simple geometric picture possible (as with real components); “angle” is not geometric angle anymore, but just a some kind of parameter now!

To easily explore further we need the **definition** of a new object - the ~~**Kronecker Delta.**~~

$$\langle n \,|\, m \rangle = \delta_{nm} = \begin{cases} 1 & n = m \\ 0 & n \neq m \end{cases}$$

for basis vectors $|n\rangle, |m\rangle$

definition just says that basis vector are unit length + orthogonal or orthonormal

then this implies that

$$\langle 1 \,|\, 1 \rangle = \delta_{11} = 1$$ **—-> length =1**

$$\langle 1 \,|\, 2 \rangle = \delta_{12} = 0$$ **—-> orthogonal**

$$\langle 2 \,|\, 1 \rangle = \delta_{21} = 0$$ **—-> orthogonal**

$$\langle 2 \,|\, 2 \rangle = \delta_{22} = 1$$ **—-> length =1**

Using Kronecker delta to do algebra -> very powerful tool that I will use to do manipulations and you will use if you make the effort to learn how to do the algebra.

Examples

$$\sum_{k=1}^{3} A_k \delta_{k2} = A_1\delta_{12} + A_2\delta_{22} + A_3\delta_{32} = A_1(0) + A_2(1) + A_3(0) = A_2$$

$$\sum_{k=1}^{n} A_k \delta_{km} = A_m \quad m \leq n$$

so appearance of kronecker delta inside a summation eliminates one summation —> a very powerful algebraic tool !!

In order to complete the mathematical tools needed to describe the microworld we need one last mathematical object —> completely new for some of you guys —> a **Matrix.**

Definition

m × n matrix is **m × n array (m rows and n columns)** of numbers(**matrix elements**) with a well-defined set of associated mathematical rules.

For example, see sample matrices shown below

$\begin{pmatrix} 2 \\ 5 \end{pmatrix}$ 2 elements

2 × 1 matrix (= column vector)

$\begin{pmatrix} 2 & 5 \end{pmatrix}$ 2 elements

1 x 2 matrix (= row vector)

$\begin{pmatrix} 2 & 5 \\ -3 & 10 \end{pmatrix}$ 4 elements

2 × 2 matrix

$$\begin{pmatrix} 2 & 5 & -1 \\ -3 & 10 & 5 \end{pmatrix}$$

2 × 3 matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad , \quad \mathbf{B} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$$

now using row-column labels

diagonal elements

off-diagonal elements

Consider two matrices

$$\mathbf{A} = \begin{pmatrix} 2 & 5 \\ -3 & 10 \end{pmatrix} \quad , \quad \mathbf{B} = \begin{pmatrix} -8 & 15 \\ 7 & 4 \end{pmatrix}$$

—> two 2 × 2 matrices (note **boldface**)

addition => add elements

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} (2) + (-8) & (5) + (15) \\ (-3) + (7) & (10) + (4) \end{pmatrix} = \begin{pmatrix} -6 & 20 \\ 4 & 14 \end{pmatrix}$$

matrix multiplication

$$\mathbf{AB} = \begin{pmatrix} 2 & 5 \\ -3 & 10 \end{pmatrix} \begin{pmatrix} -8 & 15 \\ 7 & 4 \end{pmatrix} = \begin{pmatrix} (2)(-8) + (5)(7) & (2)(15) + (5)(4) \\ (-3)(-8) + (10)(7) & (-3)(15) + (10)(4) \end{pmatrix} = \begin{pmatrix} 19 & 50 \\ 94 & -5 \end{pmatrix}$$

I just followed the **general rule**

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad , \quad \mathbf{B} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$$

using row-column labels

$$\mathbf{AB} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{pmatrix}$$

These following are definitions so that you know what I mean when I refer to some matrix property.

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

Special cases:

definition

$$\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = identity\ matrix \quad \textbf{—>} \quad \mathbf{IA} = \mathbf{AI} = \mathbf{A}$$

effect = no change

$$\det \mathbf{A} = a_{11}a_{22} - a_{12}a_{21} = determinant\ of\ \mathbf{A}$$

definitions only for 2 x 2 matrices

definition

$$\mathbf{A^{-1}} = \frac{1}{\det \mathbf{A}} \begin{pmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{pmatrix} = inverse\ matrix \quad \textbf{—>} \quad \mathbf{AA^{-1}} = \mathbf{A^{-1}A} = \mathbf{I}$$

effect

Example:

$$A = \begin{pmatrix} 1 & 4i \\ -4i & 7 \end{pmatrix} \qquad A^{-1} = \frac{1}{2 - 12i} \begin{pmatrix} 2 & -4i \\ -3 & 1 \end{pmatrix} \quad \textbf{—>} \quad AA^{-1} = I$$

$$\mathbf{A^T} = \begin{pmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \end{pmatrix} = transpose\ matrix \qquad \mathbf{A^\dagger} = \begin{pmatrix} a_{11}^* & a_{21}^* \\ a_{12}^* & a_{22}^* \end{pmatrix} = Hermitian\ conjugate\ matrix$$

definition

definition

If $\mathbf{A^\dagger = A}$ matrix = Hermitian

$$A = \begin{pmatrix} 1 & 4i \\ -4i & 7 \end{pmatrix}$$

If $\mathbf{A^\dagger = A^{-1}}$ matrix = Unitary

$$A = \frac{1}{2} \begin{pmatrix} 1+i & 1-i \\ 1-i & 1+i \end{pmatrix}$$

Finally $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ **—>** $Tr(\mathbf{A}) = a_{11} + a_{22}$ **= sum over diagonal elements**

Trace

Now start to pull it all together for QM.

or we now broaden the Dirac language for use in QM

lots of jokes…
Dirac was oblivious!

$$|V\rangle = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \text{“}ket\text{”}\ vector$$

column vector

$$\langle V| = \begin{pmatrix} v_1^* & v_2^* \end{pmatrix} = \text{“}bra\text{”}\ vector$$

new definition

row vector

Every Ket vector has a corresponding Bra vector in the Hilbert space.

Then real mathematical meaning of the “braket” is

$$\langle V\,|\,U\rangle = \begin{pmatrix} v_1^* & v_2^* \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = v_1^* u_1 + v_2^* u_2 = \text{“}braket\text{”}$$

—> rule is just matrix multiplication

From now on we assume that all ket/bra vectors are unit vectors(required for QM to be valid in the microworld as we will see)

Finally we need to think about Operators

An **operator** is a mathematical object which acts on any vector in our vector space and **results in another vector in same vector space**.

Its operation and behavior is similar to a **function with numbers** y = f(x)

insert number in function —> new number

Example

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad , \quad |V\rangle = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$\mathbf{A}\,|V\rangle = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} a_{11}v_1 + a_{12}v_2 \\ a_{21}v_1 + a_{22}v_2 \end{pmatrix} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = |U\rangle$$

just another vector in the space!!

Therefore

$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ = an operator **—> matrices are operators in QM**

why I needed to introduce the idea!

Operators are very important in QM because

observables or measurable quantities are represented in QM by Hermitian operators(matrices) where $\mathbf{A}^\dagger = \mathbf{A}$

and

transformations or physical changes of vectors are represented in QM by Unitary operators (matrices) where $\mathbf{A}^\dagger = \mathbf{A}^{-1}$

Properties of operators - $\hat{\mathbf{O}}$ (note boldface + hat)

Matrix representation (I show you this so you know what it is when I refer to it)

you will not need to do such manipulations yourself!!

Definition: Matrix element

$o_{nm} = \langle n| \hat{\mathbf{O}} |m\rangle$ = (nm) matrix element of $\hat{\mathbf{O}}$ (nm) = (row,column)

—-> matrix representing $\hat{\mathbf{O}}$ is **Using the $|1\rangle, |2\rangle$ basis**

$$\mathbf{O} = \begin{pmatrix} \langle 1| \hat{\mathbf{O}} |1\rangle & \langle 1| \hat{\mathbf{O}} |2\rangle \\ \langle 2| \hat{\mathbf{O}} |1\rangle & \langle 2| \hat{\mathbf{O}} |2\rangle \end{pmatrix} = \begin{pmatrix} o_{11} & o_{12} \\ o_{21} & o_{22} \end{pmatrix}$$

Now assume that $\mathbf{O}|1\rangle = |q\rangle \quad , \quad \mathbf{O}|2\rangle = |r\rangle$ that is what operators do!

then $o_{11} = \langle 1| \hat{\mathbf{O}} |1\rangle = \langle 1 | q\rangle$

$o_{12} = \langle 1| \hat{\mathbf{O}} |2\rangle = \langle 1 | r\rangle$

$o_{21} = \langle 2| \hat{\mathbf{O}} |1\rangle = \langle 2 | q\rangle$

$o_{22} = \langle 2| \hat{\mathbf{O}} |2\rangle = \langle 2 | r\rangle$

—-> Matrix elements are just brackets = numbers!!

Most important special case:

If $\hat{\mathbf{A}}\,|V\rangle = a\,|V\rangle$ (i.e., we get same vector back) then

$|V\rangle$ called an **eigenvector** of operator $\hat{\mathbf{A}}$ with **eigenvalue** a .

For Hermitian operators, used for observables in QM, the set of eigenvectors **always** forms an orthonormal basis -> called a **complete set** and the eigenvalues are all **real**. Very useful!

If basis set that is used to calculate matrix representing an operator

is the set of eigenvectors of the operator, i.e., if

$$\hat{\mathbf{O}}\,|1\rangle = o_1\,|1\rangle \quad , \quad \hat{\mathbf{O}}\,|2\rangle = o_2\,|2\rangle \qquad \{o_1, o_2\} \qquad \text{= eigenvalues}$$

then

$$o_{11} = \langle 1|\,\hat{\mathbf{O}}\,|1\rangle = o_1\,\langle 1\,|\,1\rangle = o_1 \quad o_{21} = \langle 2|\,\hat{\mathbf{O}}\,|1\rangle = o_1\,\langle 2\,|\,1\rangle = 0$$

$$o_{12} = \langle 1|\,\hat{\mathbf{O}}\,|2\rangle = o_2\,\langle 1\,|\,2\rangle = 0 \quad o_{22} = \langle 2|\,\hat{\mathbf{O}}\,|2\rangle = o_2\,\langle 2\,|\,2\rangle = o_2$$

$$\longrightarrow \mathbf{O} = \begin{pmatrix} \langle 1|\,\hat{\mathbf{O}}\,|1\rangle & \langle 1|\,\hat{\mathbf{O}}\,|2\rangle \\ \langle 2|\,\hat{\mathbf{O}}\,|1\rangle & \langle 2|\,\hat{\mathbf{O}}\,|2\rangle \end{pmatrix} = \begin{pmatrix} o_1 & 0 \\ 0 & o_2 \end{pmatrix}$$

Matrix representation with operator's eigenvectors is **diagonal** matrix

(nonzero elements (eigenvalues) on diagonal).

For general basis vectors(not eigenvectors),

matrix representation is generally **not** diagonal.

An **alternate(and useful) method** for defining operators uses the "ket" and "bra" vectors.

Consider the quantity

using our rules we get

$$\mathcal{P}_{fg} = |f\rangle\langle g| \longrightarrow \mathcal{P}_{fg}|V\rangle = (|f\rangle\langle g|)|V\rangle = \langle g\,|\,V\rangle\,|f\rangle = number \times |f\rangle$$

Thus, the new object is an **operator**! ⟶ Called the **"ket-bra"**

If g = f, then we get

$\mathcal{P}_f = |f\rangle\langle f|$ **= projection** operator i.e.,

$$\mathcal{P}_f|V\rangle = (|f\rangle\langle f|)|V\rangle = \langle f\,|\,V\rangle\,|f\rangle$$

"projects" $|V\rangle$ onto $|f\rangle$ i.e., final vector now in that direction

so $\langle f\,|\,V\rangle$ **gives amount of** $|V\rangle$ **in direction of** $|f\rangle$

the component

Notice we used the rule: $(\langle f|)(|f\rangle) = \langle f\,|\,f\rangle$ **to simplify notion** $||\Rightarrow|$

return to old braket symbol

Now consider set of orthonormal basis vectors: $\{|n\rangle\},\ n = 1, 2, 3, ..., N$

orthonormal **—>** $\langle k\,|\,n\rangle = \delta_{kn}$

We have(doing some algebra for practice):

$$(|k\rangle\langle k|)(|k\rangle\langle k|) = |k\rangle(\langle k\,|\,k\rangle)\langle k| = |k\rangle(1)\langle k| = |k\rangle\langle k|$$

Another example:

$$(|k\rangle\langle k|)(a|1\rangle + b|2)) = |k\rangle(a\langle k\,|\,1\rangle + b\langle k|2\rangle) = (number\rangle \times |k\rangle$$

Now for some useful properties for later derivations and examples of use of the language.

RESULTS are important - ALGEBRA is for those learning how to do it themselves!

Consider operator $\sum_{k=1}^{n} |k\rangle \langle k|$ **—> sum of allprojection operators**

Remember from earlier that for any Ket vector, we can write (example in 2 dimensions)

$$|V\rangle = \langle 1 \,|\, V\rangle \, |1\rangle + \langle 2 \,|\, V\rangle \, |2\rangle$$

or in general

$$|V\rangle = \sum_{k=1}^{2} \langle k \,|\, V\rangle \, |k\rangle$$

so in n dimensions we have(2 terms in sum —-> n terms)

$$|V\rangle = \sum_{k=1}^{n} \langle k \,|\, V\rangle \, |k\rangle$$

clearly this is very powerful notation

We have

no change!!

$$\left(\sum_{k=1}^{n} |k\rangle \langle k| \right) |V\rangle = \sum_{k=1}^{n} \langle k \,|\, V\rangle \, |k\rangle = |V\rangle \qquad \longrightarrow \qquad \sum_{k=1}^{n} |k\rangle \langle k| = \hat{\mathbf{I}}$$

the identity operator

Result will be very important later for doing algebra!

Technical term: It is called a **representation of the identity**

Expectation Value — Very(almost most) important object in QM !!

$$\langle \hat{\mathbf{O}} \rangle = \langle V | \, \hat{\mathbf{O}} \, | V \rangle$$

Will turn out(see later) to be the average value of an observable from a set of measurements on an ensemble of identical systems all in the state $|V\rangle$. Let us see how.

First, we need an **alternative representation** of an operator

(also some simple algebra practice for those interested to do)

Only the result is important (will be used) for our later discussions…….

$\hat{\mathbf{B}} \, |b_k\rangle = b_k \, |b_k\rangle \quad k = 1, 2, 3, ..., n$ **eigenvector/eigenvalues equation**

$\longrightarrow$ $|b_k\rangle$ = eigenvectors of $\hat{\mathbf{B}}$ $\longrightarrow$ b_k = eigenvalues of $\hat{\mathbf{B}}$

if operator is Hermitian $\longrightarrow$ the set $|b_k\rangle$ are an orthonormal basis so that

$$\langle b_i \, | \, b_j \rangle = \delta_{ij} \qquad \text{and} \qquad \sum_{k=1}^{n} |b_k\rangle \, \langle b_k| = \hat{\mathbf{I}}$$

Then

$$\hat{\mathbf{B}} = \hat{\mathbf{B}}\hat{\mathbf{I}} = \hat{\mathbf{B}} \sum_{k=1}^{n} |b_k\rangle \langle b_k|$$

identity operator does not change anything

$$= \sum_{k=1}^{n} \hat{\mathbf{B}} |b_k\rangle \langle b_k| = \sum_{k=1}^{n} b_k |b_k\rangle \langle b_k|$$

operator moves through summation and operates on Ket vector

or

$$\hat{\mathbf{B}} = \sum_{k=1}^{n} b_k |b_k\rangle \langle b_k|$$

—> operator written in terms of eigenvalues/eigenvectors or projection operators

Called the “spectral representation” of the operator

Now further algebra gives

(if interested see full mathematics supplementary lecture notes for details)

$$\hat{\mathbf{B}}^2 = \sum_{k=1}^{n} b_k^2 |b_k\rangle \langle b_k| \qquad \hat{\mathbf{B}}^3 = \sum_{k=1}^{n} b_k^3 |b_k\rangle \langle b_k| \qquad \hat{\mathbf{B}}^n = \sum_{k=1}^{n} b_k^n |b_k\rangle \langle b_k|$$

Definition: Power series representation of a function:

Sometimes a function can be written as a “power series” as shown below:

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots = \sum_{k=0}^{\infty} a_k x^k$$

Examples(these are the real definitions of the functions):

$$e^{\alpha x} = \sum_{k=0}^{\infty} \frac{1}{k!} \alpha^k x^k$$

$$\sin \alpha x = \sum_{k=0}^{\infty} \frac{1}{(2k+1)!} (-1)^k \alpha^{2k+1} x^{2k+1} \qquad \cos \alpha x = \sum_{k=0}^{\infty} \frac{1}{(2k)!} (-1)^k \alpha^{2k} x^{2k}$$

Now consider

$$e^{\alpha x} = \sum_{k=0}^{\infty} \frac{1}{k!} \alpha^k x^k \xrightarrow{x \to \hat{\mathbf{O}}} e^{\alpha \hat{\mathbf{O}}} = \sum_{k=0}^{\infty} \frac{1}{k!} \alpha^k \hat{\mathbf{O}}^k$$

get a function of an operator now

In special case where $\hat{\mathbf{O}}^2 = \hat{\mathbf{I}}$ (algebra is in mathematics supplement)

$$e^{i\alpha \hat{\mathbf{O}}} = \cos \alpha \hat{\mathbf{I}} + i \hat{\mathbf{O}} \sin \alpha$$

This is one of the most powerful relations in QM

(original derivation was for ordinary functions by Euler in 16th century

$$e^{i\theta} = \cos \theta + i \sin \theta \qquad \rightarrow e^{i\pi/2} = i \;\; , \;\; e^{i\pi} = -1$$

—> real definition of i

In general, if we need an operator for an arbitrary function $f(x)$ we do the following:

if $\hat{\mathbf{B}}\,|b_k\rangle = b_k\,|b_k\rangle \quad k = 1, 2, 3, ..., n$ **then** $f(\hat{\mathbf{B}}) = \sum_{k=1}^{n} f(b_k)\,|b_k\rangle\,\langle b_k|$

Another Important Definition

$$\text{commutator } = [\hat{\mathbf{A}}, \hat{\mathbf{B}}] = \hat{\mathbf{A}}\hat{\mathbf{B}} - \hat{\mathbf{B}}\hat{\mathbf{A}}$$

ordinary numbers “commute” but operators may not “commute”

This is mathematical property behind Heisenberg uncertainty principle as we will see later.

Reminder: Useful properties to remember:

$$|V\rangle = v_1\,|1\rangle + v_2\,|2\rangle \quad , \quad |U\rangle = u_1\,|1\rangle + u_2\,|2\rangle$$

$$\langle V\,|\,U\rangle = v_1^* u_1 + v_2^* u_2 \qquad \langle U\,|\,V\rangle = u_1^* v_1 + u_2^* v_2 = \langle V\,|\,U\rangle^*$$

$$|V\rangle = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \qquad \langle V| = \begin{pmatrix} v_1^* & v_2^* \end{pmatrix} = |V\rangle^\dagger$$

Suppose in the middle of a dream, I have an interesting thought….(this is example of how a theoretical physicist works…….)

$\langle \hat{\mathbf{B}} \rangle = \langle V | \hat{\mathbf{B}} | V \rangle$ = expectation value

$\hat{\mathbf{B}}$ is Hermitian —>

$\hat{\mathbf{B}}$ eigenvectors are complete orthonormal set = basis —> can make expansions —> so write

$$|V\rangle = \sum_{k=1}^{n} d_k |b_k\rangle \quad , \quad d_k = \langle b_k | V \rangle$$ definition of components

and

$$\langle V| = \sum_{k=1}^{n} d_k^* \langle b_k|$$

We can always do this with any basis set!!

More algebra gives(see supplement):

$$\langle V | \hat{\mathbf{B}} | V \rangle \quad = \quad \langle \hat{\mathbf{B}} \rangle \quad = \sum_{k=1}^{n} b_k |d_k|^2$$

expectation value of the operator in the state |V> using eigenvector of operator **B**

But, in general, if expectation value = average value, then we must have

$$\langle \hat{\mathbf{B}} \rangle = \sum_{allowed\ values} (\text{allowed value of B}) \times (\text{Probability of that value})$$

which is just the **definition** of the average value!!

Now we just derived

$$\langle \hat{\mathbf{B}} \rangle = \sum_{k=1}^{n} b_k |d_k|^2$$

Comparing(see arrows) last two equations - we might guess that in our future theory

allowed measurement values = eigenvalues

and

probability of a value = square of corresponding vector component value

Now we can write

$$|d_k|^2 = d_k^* d_k = (\langle b_k \,|\, V\rangle)^* \,\langle b_k \,|\, V\rangle = |\,\langle b_k \,|\, V\rangle\,|^2 = \langle V \,|\, b_k\rangle\,\langle b_k \,|\, V\rangle$$

This assumption would then say that

$|\,\langle b_k \,|\, V\rangle\,|^2$ = probability of observing value b_k when in state $|V\rangle$

It also says

$$\langle V \,|\, b_k\rangle\,\langle b_k \,|\, V\rangle = |\,\langle b_k \,|\, V\rangle\,|^2 = |\text{component}|^2$$

= probability of observing value b_k in state $|V\rangle$

= expectation value of the projection operator $|\,b_k\rangle\,\langle b_k\,|$

These ideas will become postulates - that is what random thinking sometimes produces.

That is the kind of dream a theoretical physicist might have!

That is all mathematics we will need for now(we will add a couple of things later) to proceed with developing QM.

Lots of repetition coming…….as we use this new language!

As we worked through the mathematics

we have laid the ground work for the postulates of QM

(in some cases we have actually stated the postulate already).

We will not need any more mathematics than I have shown you already, although we will add an extension here and there.

You will get better at using the mathematics as we will use it in class and you get used to the language.

If you cannot follow all the details of the mathematics, then just follow along with the ideas that I will present based on the mathematics.

That will be sufficient to understand QM.

Applying Mathematics ideas to Quantum Mechanics

We now expand our understanding of QM theory and also explain the color/hardness experiments

Quantum state = set of information **from** measurements.

It will be represented by a KET $|....\rangle$ $|banana\rangle\ ,|747\rangle\ ,|you\rangle$

where ……. = labels indicating what we know(must have measured) about the state

important aspect

Thus

electron - no measurements yet	$\|undetermined\rangle = \|?\rangle$
electron into COLOR box emerges magenta aperture	$\|magenta\rangle = \|m\rangle$
electron into COLOR box emerges green aperture	$\|green\rangle = \|g\rangle$
electron into HARDNESS box emerges hard aperture	$\|hard\rangle = \|h\rangle$
electron into HARDNESS box emerges soft aperture	$\|soft\rangle = \|s\rangle$

Clearly the labels tell us everything we "know" about the state

As we will see later — **Postulate #1 of quantum theory will be — it is all in the labels!**

Continuing, a measuring device is represented by an OPERATOR

Definition: Operators take Kets into Kets $\hat{O}\left|x\right\rangle = \left|y\right\rangle$

Some operator properties from experimental results:

$$\mathcal{P}_g\left|hard\right\rangle = \left(\left|g\right\rangle\left\langle g\right|\right)\left|h\right\rangle \;\; = \left\langle g\,|\,h\right\rangle\left|g\right\rangle \rightarrow \left|g\right\rangle$$ All state always **renormalized** to 1

green projection operator

$$\mathcal{P}_m\left|h\right\rangle = \left(\left|m\right\rangle\left\langle m\right|\right)\left|h\right\rangle \;\; = \left\langle m\,|\,h\right\rangle\left|m\right\rangle \rightarrow \left|m\right\rangle$$

magenta projection operator

Last step:

All state always normalized to 1

$\left\langle g\,|\,g\right\rangle = 1$ by definition

State $\left|Q\right\rangle = \left\langle g\,|\,h\right\rangle\left|g\right\rangle$

is not normalized to 1

—> always renormalize to 1 at **each** stage

Projection operators **seem** to represent measurement(pick out states) - we will see!

You will note that I do not tell you WHY I make certain choices, I only tell you HOW the my particular choices work.

That is the difference between Physics and Philosophy.

Physics NEVER answers WHY questions, we leave those for Philosophers, who have never once answered a “why” question!!

Now send a magenta electron into a hardness box

it comes out either hard or soft (50%-50%) (out the appropriate aperture).

A magenta electron was described earlier as a **SUPERPOSITION** of hard and soft electrons,

although we did not, at that time, specify what was meant by the term “superposition”.

Now we represent a **superposition** mathematically by **addition of kets** (simply vector addition) as follows:

$$|m\rangle = \frac{1}{\sqrt{2}}|h\rangle - \frac{1}{\sqrt{2}}|s\rangle$$

$$|g\rangle = \frac{1}{\sqrt{2}}|h\rangle + \frac{1}{\sqrt{2}}|s\rangle$$

$$|h\rangle = \frac{1}{\sqrt{2}}|g\rangle + \frac{1}{\sqrt{2}}|m\rangle$$

$$|s\rangle = \frac{1}{\sqrt{2}}|g\rangle - \frac{1}{\sqrt{2}}|m\rangle$$

This will be Postulate #2 of quantum theory.

The reasons for the particular numerical coefficients will become clear shortly.

This is the way theory works:

Make assumptions(postulates),

i.e., meaning of term “superposition” above.

Develop measurable predictions from the assumptions.

Then test experimentally to check the validity of the postulates.

More about Probability

Suppose we have a box with N numbered balls

set of numbers **on** balls number of balls **with** given number

$$\{\nu_i\} = \nu_1, \nu_2, \nu_3, \nu_4, ... \quad , \quad \{n_i\} = n_1, n_2, n_3, n_4, ...$$



$$\sum_k n_k = n_1 + n_2 + n_3 + n_4 + ... = N$$

total number

probability that a random picked ball has kth number ν_k is $p_k = \frac{n_k}{N}$

$n_k = 0$ —> $p_k = 0$ **impossible**

$n_k = N$ —> $p_k = 1$ **certainty**

$$0 \leq p_k \leq 1 \text{ for all } k \quad , \quad \sum_k p_k = 1$$

above are all common sense ideas

In QM

$$probability(q) = \frac{number\, of\, times\, value\, q\, was\, already\, measured}{total\, number\, of\, measurements\, so\, far}$$

identical systems —> create ensemble —> allows frequency model representing probabilities

Using the standard definition of the average one can also show $\langle f(\nu) \rangle = \sum_k p_k f(\nu_k)$ for any function.

Continuing with QM discussion

Although QM is a probabilistic theory, the fundamental quantity will **not** be probability but instead it is a mathematical object called the **probability amplitude,** which we will define below.

Now if the electron to be measured was in an arbitrary state $|\psi\rangle$,

then probability amplitude that one will measure color = magenta

if one sends this electron into a color box is given by

a new mathematical object (amplitude) **represented** by the Dirac braket symbol

$$\langle property\,to\,be\,measured \,|\, state\,being\,measured \rangle = \langle m \,|\, \psi \rangle$$

—-> complex number = component of ket $|\psi\rangle$ in $|m\rangle$ direction

amount

$$A = probability\,amplitude = \langle m \,|\, \psi \rangle$$ **The Dirac Bracket!!!!**

$$P = probability = |\,\langle m \,|\, \psi \rangle\,|^2 \;\; = |A|^2$$

= probability to measure magenta if electron in state $|\psi\rangle$

This will be Postulate #3 of quantum theory

Now, we use this idea and see what it actually says, i.e., what it implies……

Send an electron into a color box

and look at the beam emerging from magenta aperture

= magenta electrons - all in state $|m\rangle$ if box is working properly

Experiments can be used to determine the numerical value of amplitudes:

Probability that magenta electron emerges from magenta aperture of color box

$$P = |\langle m \mid m \rangle|^2 = 1 \longrightarrow \langle m \,|\, m \rangle = 1$$

Probability that magenta electron emerges from green aperture of color box

$$P = |\langle m \mid g \rangle|^2 = 0 \longrightarrow \langle m \,|\, g \rangle = 0$$

Probability that hard electron emerges from hard aperture of hardness box

$$P = |\langle h \mid h \rangle|^2 = 1 \longrightarrow \langle h \,|\, h \rangle = 1$$

Probability that hard electron emerges from soft aperture of hardness box

$$P = |\langle h \mid s \rangle|^2 = 0 \longrightarrow \langle h \,|\, s \rangle = 0$$

Similarly, we have

$$\langle g \,|\, g \rangle = 1 \qquad \langle s \,|\, s \rangle = 1 \qquad \langle g \,|\, m \rangle = 0 \qquad \langle s \,|\, h \rangle = 0$$

What do these numerical results mean?

Clearly,

kets $\{|m\rangle, |g\rangle\}$ or kets $\{|h\rangle, |s\rangle\}$ from earlier discussions

are just two different orthonormal bases for all quantum states in world of color/hardness

$\langle m \,|\, m\rangle = 1$ $\langle g \,|\, g\rangle = 1$ $\langle m \,|\, g\rangle = 0$

normalized to 1 orthogonal

—> orthonormal set

Using one of these basis sets to describe world —> using “color” or “hardness” language

Since they are a basis ⟶ why we can write any arbitrary quantum state

as $|\psi\rangle = a\,|m\rangle + b\,|g\rangle$

or as $|\psi\rangle = c\,|h\rangle + d\,|s\rangle$

Remember I wrote earlier

$$|m\rangle = \frac{1}{\sqrt{2}}\,|h\rangle - \frac{1}{\sqrt{2}}\,|s\rangle \qquad |g\rangle = \frac{1}{\sqrt{2}}\,|h\rangle + \frac{1}{\sqrt{2}}\,|s\rangle$$

$$|h\rangle = \frac{1}{\sqrt{2}}\,|g\rangle + \frac{1}{\sqrt{2}}\,|m\rangle \qquad |s\rangle = \frac{1}{\sqrt{2}}\,|g\rangle - \frac{1}{\sqrt{2}}\,|m\rangle$$

Mysterious property called **SUPERPOSITION** is, as we stated, **just vector addition!**

Does this formalism work?

Does it allow us to say correct things about quantum experiments?

YES! Let me show you.

Consider hard electron sent into color box,

then, using rules we have developed,

the probability that it will be (**emerge as or be measured as**)

a magenta electron is given by(**this is how the algebra works**)

$$|\langle magenta \mid hard\rangle|^2 = \left|\langle magenta| \left[\frac{1}{\sqrt{2}}\,|green\rangle + \frac{1}{\sqrt{2}}\,|magenta\rangle\right]\right|^2$$

$$= \left|\frac{1}{\sqrt{2}}\,\langle magenta \mid green\rangle + \frac{1}{\sqrt{2}}\,\langle magenta \mid magenta\rangle\right|^2$$

$$= \left|\frac{1}{\sqrt{2}}(0) + \frac{1}{\sqrt{2}}(1)\right|^2 = \frac{1}{2}$$ **which is the correct probability!!**

and probability that it will emerge as green electron given by

$$|\langle green \mid hard\rangle|^2 = \left|\langle green| \left[\frac{1}{\sqrt{2}}\,|green\rangle + \frac{1}{\sqrt{2}}\,|magenta\rangle\right]\right|^2$$

$$= \left|\frac{1}{\sqrt{2}}\,\langle green \mid green\rangle + \frac{1}{\sqrt{2}}\,\langle green \mid magenta\rangle\right|^2$$

$$= \left|\frac{1}{\sqrt{2}}(1) + \frac{1}{\sqrt{2}}(0)\right|^2 = \frac{1}{2}$$ **which is also correct!!**

Thus, a hard electron has 50-50 chance

of coming out of magenta and green apertures

which agrees with the results of the earlier experiments.

That is why I earlier chose particular numerical coefficients (component values) $\pm\frac{1}{\sqrt{2}}$

So our formalism seems to agree so far with all of the earlier experiments.

For example, look at repeatability experiment.

First, hard electron $|h\rangle$ sent into color box.

Probability will emerge from magenta aperture is

$$|\langle m \mid h\rangle|^2 = \frac{1}{2}$$

Then, electron emerging from magenta aperture sent into another color box.

However, it is **now** a magenta electron and is represented by ket |m⟩.

Probability that it will emerge from magenta aperture of second color box is

$$|\langle m \mid m \rangle|^2 = 1$$ **= repeatability property.**

During experiment,

hard electron(initially in superposition of magenta/green properties), i.e.,

$$|h\rangle = \frac{1}{\sqrt{2}}|g\rangle + \frac{1}{\sqrt{2}}|m\rangle$$

appears as electron in state |m⟩

when it emerges from magenta aperture of first color box.

The so-called **Copenhagen** interpretation of quantum theory says

that during 1st measurement (first color box), the state of electron(hard)

is "**collapsed" or "reduced"** from a superposition of possibilities to a definite

value(magenta) corresponding to value just measured, i.e., to aperture we looked at!

This interpretation seems to say that the measurement **caused** the collapse to occur.


This will be postulate #4 of our 1st version of quantum theory.

Eventually, we will eliminate the need for this "collapse" postulate completely.

We will find that it is already built into the other postulates we define!

Two-Path Experiment now explained using the New Formalism

Added time and position values to diagram.



Need to use states or kets which correspond

to electrons having color or hardness

AND positions.

This is OK because these are what are called "**compatible** properties",

i.e., we will see that it is valid in QM to

say a magenta electron is located

at a particular position

(since it can be created in lab).

Thus, electrons can have color (or hardness) and simultaneously be at definite position.

However, as saw earlier,

an electron CANNOT be hard AND magenta simultaneously.


These are incompatible properties, which we will see is related to the Uncertainty Principle.

These states take the form (just **expand the labels** because we **KNOW** more stuff):

$$|color, x, y\rangle = |color\rangle\,|x, y\rangle$$

$$|hardness, x, y\rangle = |hardness\rangle\,|x, y\rangle$$

so the labels still specify all that we "know"



At time t_1,

when particle is about to enter apparatus,

(having just left a color box via

magenta aperture)

the state is

$$|color = magenta, x = x_1, y = y_1\rangle = |m\rangle\,|x_1, y_1\rangle \qquad \text{at } t_1$$

$$= \left(\frac{1}{\sqrt{2}}\,|h\rangle - \frac{1}{\sqrt{2}}\,|s\rangle\right)|x_1, y_1\rangle = \frac{1}{\sqrt{2}}\,|h\rangle\,|x_1, y_1\rangle - \frac{1}{\sqrt{2}}\,|s\rangle\,|x_1, y_1\rangle$$

(everything just multiplied out)

It is a superposition of a hard electron at (x_1,y_1) and soft electron at (x_1,y_1)!

A magenta electron is always such a superposition at any given position!

Here is how QM will tell us to calculate what happens next.

Consider: If state at time t_1 were just the hard part

$$\frac{1}{\sqrt{2}}\,|h\rangle\,|x_1, y_1\rangle$$

and if a hardness box behaves properly,


then the state at time t_2 would be

$$\frac{1}{\sqrt{2}}\,|h\rangle\,|x_2, y_2\rangle$$

i.e., the electron would have just emerged

through hard aperture

and be on what we will call the **hard path**.



Similarly, if state at time t_1 were just soft part $\frac{1}{\sqrt{2}}\,|s\rangle\,|x_1, y_1\rangle$

and if hardness box behaves properly, then state at time t_2 would be $\frac{1}{\sqrt{2}}\,|s\rangle\,|x_3, y_1\rangle$

i.e., electron would have just emerged through soft aperture

and be on what we will call the **soft path.**

This is what we earlier said the apparatus does to hard/soft electrons!


IMPORTANT: we can only say the electron has two compatible measurable properties at the same time if the respective state separates exactly like this one did.

However, as we said, the state at t_1

is neither just a soft part nor just a hard part,

but the superposition

$$|m\rangle\,|x_1, y_1\rangle = \frac{1}{\sqrt{2}}\,|h\rangle\,|x_1, y_1\rangle - \frac{1}{\sqrt{2}}\,|s\rangle\,|x_1, y_1\rangle$$



when using color box

-> USE color language to describe world —> start in color language(basis)

when using hardness box

-> USE hardness language to describe world —> change to hardness language(basis)

This is VERY Important

Physics will only make sense if you are using the correct language (appropriate to next measurement) when discussing what is happening!

Earlier discussion therefore says

that state at time t_2 must be (if hardness box behaves properly)

$$\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_2,y_2\right\rangle - \frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_3,y_1\right\rangle$$

|1> **|2>**

i.e., if something happens to state |1> (hard part) and something happens to state |2> (soft part) when sent through separately

then when both are present at same time in a superposition **in words**

SAME things still happen to each part

[Remember we are not measuring anything yet!!]

So we have at time t_2 $\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_2,y_2\right\rangle - \frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_3,y_1\right\rangle$

Now expand

$$\frac{1}{\sqrt{2}}\left(\frac{1}{\sqrt{2}}\left|g\right\rangle + \frac{1}{\sqrt{2}}\left|m\right\rangle\right)\left|x_2,y_2\right\rangle - \frac{1}{\sqrt{2}}\left(\frac{1}{\sqrt{2}}\left|g\right\rangle - \frac{1}{\sqrt{2}}\left|m\right\rangle\right)\left|x_3,y_1\right\rangle$$

$$\longrightarrow \quad \frac{1}{2}\left|g\right\rangle\left(\left|x_2,y_2\right\rangle - \left|x_3,y_1\right\rangle\right) + \frac{1}{2}\left|m\right\rangle\left(\left|x_2,y_2\right\rangle + \left|x_3,y_1\right\rangle\right) \qquad \text{at time } t_2$$

This correspond to color electrons at superpositions of two different positions !!!!

i.e., a green electron can be found at two positions if we look(observe/measure)

i.e., there is a nonzero probability to be at two positions

The state

$$\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_2, y_2\right\rangle - \frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_3, y_1\right\rangle$$

is an **entanglement** between hardness/color properties

and position (coordinate space) properties of electron

(note color and position properties

were separated to start with, i.e., $\left|color = magenta, x = x_1, y = y_1\right\rangle = \left|m\right\rangle\left|x_1, y_1\right\rangle$

i.e., **had definite values**, namely, magenta and (x_1,y_1)).

This new state involves **nonseparable correlations(entanglement)** between hardness/color and coordinate-space properties of electron.

There are no definite hardness or color properties of electron in this state.

Neither do any of its coordinate-space properties (position, momentum, etc)

have any definite values — the hardness box did that!!!

This is a **superposition** of states,

where one part of electron state indicates

probability amplitude for being on hard path

and other of electron state indicates

probability amplitude for being on soft path.

IMPORTANT: Note that we are NOT saying that electron is actually on either path.

REMEMBER: No measurements have been made yet!!!!!

These results are the reason that

we earlier had to make statements about not hard, not soft, not both and not neither.

Those ideas are not only false, they are meaningless in this state!

We have entangled the properties of two **classical states** that **cannot exist** at same time for single electron!

That is the idea of the state that is called an **entangled state**.


These states will get our thoughts/ideas all entangled later on

when we discuss the Einstein paradox and Bell inequality!!

Continuing on.....

last state of electron above (at t_2)

then leads to state of electron at t_3



$$\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_3,y_3\right\rangle-\frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_4,y_2\right\rangle$$

and then state at time t_4 is

$$\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_5,y_4\right\rangle-\frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_5,y_4\right\rangle \quad=\left(\frac{1}{\sqrt{2}}\left|h\right\rangle-\frac{1}{\sqrt{2}}\left|s\right\rangle\right)\left|x_5,y_4\right\rangle$$

$$=\left|m\right\rangle\left|x_5,y_4\right\rangle$$

pull out common factor

$$\frac{1}{\sqrt{2}}|h\rangle|x_5,y_4\rangle - \frac{1}{\sqrt{2}}|s\rangle|x_5,y_4\rangle = \left(\frac{1}{\sqrt{2}}|h\rangle - \frac{1}{\sqrt{2}}|s\rangle\right)|x_5,y_4\rangle = |m\rangle|x_5,y_4\rangle$$

We see

that color state and coordinate-space state have become **separate** again!

Position of electron **once again** has definite value and color **once again** has definite value!!

Remember that we have not bothered(measured) the electron during this experiment.

So fact that hard electron fed into total device will come out hard(follow hard path),

and that soft electron fed into total device will come out soft(follow soft path),

together with our quantum assumptions,

means that a magenta electron fed into total device

will come out (as EXPERIMENT says) magenta!

That is way quantum mechanics works! Again, the formalism agrees with experiment!

What if we **change the experiment** in middle by measuring position of the electron at, say, t_3?

For example, let us insert wall on soft path.

This Implies that if any electron gets through device, it **must** have traveled on hard path.

This is a measurement of position.

Same goes for if inserting wall on hard path.

Then the superposition goes away.

According to the suggested postulates,

during a measurement a collapse **would** occur

and state just after measurement would be either

$$|h\rangle\,|x_3, y_3\rangle \quad or \quad |s\rangle\,|x_4, y_2\rangle$$

block soft path — block hard path

each with probability 1/2 (square component assumption)

Then state at t_4 would be either

$$|h\rangle\,|x_5, y_4\rangle \quad or \quad |s\rangle\,|x_5, y_4\rangle$$

block soft path — block hard path

After this point in time,

only one part of state continues to evolve in time,

while other remains unchanged (due to the wall).

Must emphasize that we collapsed the particle to hard or soft

by measuring its position in an entangled state.

That is way entangled or nonseparable states work

collapsing one property (position in this case)

collapses all other properties (hardness in this case) simultaneously.

The state vector

$$\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_5, y_4\right\rangle - \frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_3, y_1\right\rangle$$

is **nonseparable**

between hardness/color properties and coordinate-space properties.

This means it is **not associated** with any definite values

for hardness or color or position or momentum.

Note that since we have inserted a wall on only one path(soft),

the remaining electrons in device have **definite hardness**,

which are 50/50 magenta/green.

This agrees with experiments described earlier.

Again, the formalism works!

Now I must say something about how to build a **do-nothing box**.

If electron is in a state $|A\rangle$ and we measure color,

then the probability of measuring magenta is determined as shown below.

Write $|A\rangle$ as superposition of color states

(since color is thing want to measure, we must use color language).

$$|A\rangle = \alpha\,|magenta\rangle + \beta\,|green\rangle$$

Can always be done since color kets are a basis(anything measurable provides a basis).

This says that

$$\langle magenta \mid A\rangle = \alpha\,\langle magenta \mid magenta\rangle + \beta\,\langle magenta \mid green\rangle = \alpha$$

$$\langle green \mid A\rangle = \alpha\,\langle green \mid magenta\rangle + \beta\,\langle green \mid green\rangle = \beta$$

The probability that the measurement indicates the electron is magenta is

$$|\langle magenta \mid A\rangle|^2 = |\alpha|^2$$

Now consider the state

$$-|A\rangle = -\alpha\,|magenta\rangle - \beta\,|green\rangle$$

If we measure the color on this state,

we get same probabilities for every possible outcome as with state |A⟩, i.e.,

$$|\langle magenta \mid A\rangle|^2 = |\alpha|^2 = |-\langle magenta \mid A\rangle|^2$$

So if a state vector |A⟩ changes to state vector –|A⟩

there are NO OBSERVABLE CHANGES

there are NO PROBABILITIES CHANGES

NOTHING HAPPENS to that state as far as quantum theory is concerned!!!.

So any box which changes state of any incoming electron

into –1 times incoming state will be **do-nothing box**,

since it will not change any measurable values,

i.e., will not change any probabilities

of values of any of observables of any electron which passes through it.

Obviously, it will also not effect any electron which passes outside of it.

But the effects of such a box on an electron

which is in a superposition of passing through it and outside of it

may be **quite a different matter**.


Suppose a do-nothing box is inserted in soft path of two-path device at (x_3,y_1).

Then, if initial input electron magenta(as earlier), state at t_2 will be

$$\frac{1}{\sqrt{2}}\,|h\rangle\,|x_2,y_2\rangle - \frac{1}{\sqrt{2}}\,|s\rangle\,|x_3,y_1\rangle \qquad \text{(same as earlier)}$$

and state at t_3 (after passage through box) will be not be

$$\frac{1}{\sqrt{2}}\,|h\rangle\,|x_3,y_3\rangle - \frac{1}{\sqrt{2}}\,|s\rangle\,|x_4,y_2\rangle \qquad \text{(earlier result)}$$



but will be

$$\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_3,y_3\right\rangle+\frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_4,y_2\right\rangle$$

where sign of 2nd term has been changed by the do-nothing box in the soft path.

If we now follow new state to t_4 as before we find

$$\begin{aligned}\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_5,y_4\right\rangle+\frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_5,y_4\right\rangle &= \left(\frac{1}{\sqrt{2}}\left|h\right\rangle+\frac{1}{\sqrt{2}}\left|s\right\rangle\right)\left|x_5,y_4\right\rangle\\ &= \left|g\right\rangle\left|x_5,y_4\right\rangle\end{aligned}$$

instead of

$$\begin{aligned}\frac{1}{\sqrt{2}}\left|h\right\rangle\left|x_5,y_4\right\rangle-\frac{1}{\sqrt{2}}\left|s\right\rangle\left|x_5,y_4\right\rangle &= \left(\frac{1}{\sqrt{2}}\left|h\right\rangle-\frac{1}{\sqrt{2}}\left|s\right\rangle\right)\left|x_5,y_4\right\rangle\\ &= \left|m\right\rangle\left|x_5,y_4\right\rangle\end{aligned}$$

So a do-nothing box has changed color of all electrons from magenta to green

(as the experiment said) even though it has no measurable effect

on electrons that passed though it !!!!.

That is way quantum mechanics works!! Hopefully you think it is strange!!

The new mathematical formalism seems to works very well since the theory it provides agrees with experiment, **which is only test required!**

Short digression: Some further mathematical thoughts based on what we have learned and to clean up some loose ends.

Define the operator

$$\hat{G} = |g\rangle \langle g|$$

Properties

$$\hat{G}\,|g\rangle = |g\rangle \langle g \mid g\rangle = |g\rangle \quad , \quad \hat{G}\,|m\rangle = |g\rangle \langle g \mid m\rangle = 0$$

renormalization

Expectation value

$$\langle g|\,\hat{G}\,|g\rangle = |\langle g \mid g\rangle|^2 = 1 \qquad \text{in green state}$$

$$\langle m|\,\hat{G}\,|m\rangle = |\langle g \mid m\rangle|^2 = 0 \qquad \text{in magenta state}$$

These results make sense(physics) if we interpret $\hat{G} = |g\rangle \langle g|$

as operator corresponding to measurement of green property of electrons,

i.e., it represents an observer looking at the output of the green aperture of the color box.

1st result then says, using an earlier result we derived $prob(b_k) = |\langle b_k \mid \psi \rangle|^2$

$$\langle g| \hat{G} |g\rangle = |\langle g \mid g \rangle|^2 = 1 \qquad \text{in green state}$$

that probability that color of green electron is green is 1 as expected

and

2nd result says if measure probability color of green electron is magenta get 0 as expected.

$$\langle m| \hat{G} |m\rangle = |\langle g \mid m \rangle|^2 = 0 \qquad \text{in magenta state}$$

Things make sense!

Pushing these strange ideas further.

If we assume, as earlier, that hard electron is superposition of green and magenta electrons,

$$|hard\rangle = |h\rangle = \frac{1}{\sqrt{2}} |g\rangle + \frac{1}{\sqrt{2}} |m\rangle$$

then the expectation value of $\hat{G}$ in hard state is

$$\langle g|\,\hat{\mathbf{G}}\,|h\rangle = \left(\frac{1}{\sqrt{2}}\langle g| + \frac{1}{\sqrt{2}}\langle m|\right)\hat{\mathbf{G}}\left(\frac{1}{\sqrt{2}}|g\rangle + \frac{1}{\sqrt{2}}|m\rangle\right)$$

$$= \frac{1}{2}\langle g|\,\hat{G}\,|g\rangle + \frac{1}{2}\langle g|\,\hat{G}\,|m\rangle + \frac{1}{2}\langle m|\,\hat{G}\,|g\rangle + \frac{1}{2}\langle m|\,\hat{G}\,|m\rangle$$

$$= \frac{1}{2}\langle g|\,\hat{G}\,|g\rangle = \frac{1}{2}$$ i.e., equal parts 0(m) and 1(g) as expected!!!!

Another way of saying this is, using earlier result (for those who like more math)

$$\left\langle \hat{B} \right\rangle = \sum_k b_k prob(b_k) = \sum_k b_k \left|\langle b_k \mid \psi\rangle\right|^2$$

$$\langle h|\,\hat{G}\,|h\rangle = \sum (\text{eigenvalue g })(\text{probability of g in } |h\rangle)$$

$$= (1)\left|\langle eigenvalue = 1 \mid h\rangle\right|^2 + (0)\left|\langle eigenvalue = 0 \mid h\rangle\right|^2$$

$$= (1)\left|\langle g \mid h\rangle\right|^2 + (0)\left|\langle m \mid h\rangle\right|^2 = (1)\frac{1}{2} + (0)\frac{1}{2} = \frac{1}{2}$$

again makes sense,

i.e., if have beam of hard electrons,

then we measure electron to be green 1/2 of time as observed earlier!

Clearly, this formalism and associated ideas is both neat and very powerful
and certainly seems to have the potential to describe earlier observations.

We will see shortly that this formalism, based on set of postulates,
can completely represent quantum systems and quantum measurements.

Now let us summarize some of ideas have been discussing.

More times we think about it, better we will understand it.

Sometimes theory is just **guessed** (educated guesses based on experiments)
as a set of postulates.

Let us take this approach at this time.



“The mathematician plays a game in which he himself invents the rules while the physicist plays a game in which the rules are provided by nature, but as time goes on it becomes increasingly evident that the rules which the mathematician finds interesting are the same as those which nature has chosen”

— Paul A.M. Dirac

Quantum Mechanics Postulates (using the color/hardness world)

5 parts(ASSUMPTIONS/AXIOMS) to QM algorithm.

(A) 1st Postulate -**Physical States** - **All** physical systems represented by ket vectors normalized to 1.

Called **"state vectors"** → $|\psi\rangle$ where $\langle\psi \mid \psi\rangle = 1$.

Literally means ALL.

A green or hard electron is represented by a ket.

An atom is represented by a ket.

A banana is represented by a ket.

A car is represented by a ket.

YOU are represented by a ket (albeit a very complex one).

Ket labels consist of everything we "know" i.e., have measured about the system represented by the Ket!

(B) 2nd Postulate - **Measurable Properties = observables**

Remember: for linear operators: $(\hat{A}+\hat{B})\,|\psi\rangle = \hat{A}\,|\psi\rangle + \hat{B}\,|\psi\rangle$

If $\hat{A}\,|\psi\rangle = \alpha\,|\psi\rangle$ then $|\psi\rangle$ is an eigenvector of $\hat{A}$ and α is the corresponding eigenvalue.

If $\hat{A}$ is an observable, then the system represented by the state $|\psi\rangle$ **has** the value α of that observable in that state.

That is, if you perform measurement corresponding to $\hat{A}$ on system in state represented by $|\psi\rangle$, then with **certainty** (probability = 1) you measure value α .

Since eigenvalues of operators representing observables are supposed to be measurable numbers, they must also be real numbers.

This means we can only choose a certain kind of operator to represent observables, namely, HERMITIAN operators that are guaranteed to always have real eigenvalues => added bonus for quantum theory is

Eigenvectors of HERMITIAN operator are always a complete, orthonormal set.

i.e., they comprise a set of mutually orthonormal vectors - basis - **see later that there are important connections to measurement here!**

Example: back to color and hardness.....

Operators $\hat{H}$ and $\hat{C}$ representing observables hardness/color are Hermitian

-> use either set of eigenvectors as basis set

for quantum theory of color and hardness.

Corresponds to choosing a language!!!!

One such basis is then $|hard\rangle = |h\rangle \quad , \quad |soft\rangle = |s\rangle$

where(by definition)

$$\hat{H}\,|h\rangle = |h\rangle \rightarrow eigenvalue = 1 \quad (by\ convention)$$
$$\hat{H}\,|s\rangle = -\,|s\rangle \rightarrow eigenvalue = -1 \quad (by\ convention)$$


± 1 values are arbitrary

or

|h⟩ is state with measured value of hardness = 1

|s⟩ is state with measured value of hardness = –1

on some meter!



Since the basis is an orthonormal set -> satisfies

$$\langle h \mid h\rangle = 1 = \langle s \mid s\rangle$$
$$\langle h \mid s\rangle = 0 = \langle s \mid h\rangle$$

Showed earlier, any operator can be written in terms of its eigenvalues and projection operators (spectral decomposition)

$$\hat{H} = (+1)\,|h\rangle\langle h| + (-1)\,|s\rangle\langle s| = |h\rangle\langle h| - |s\rangle\langle s|$$

Using $\hat{H} = (+1)\,|h\rangle\langle h| + (-1)\,|s\rangle\langle s| = |h\rangle\langle h| - |s\rangle\langle s|$

$$\hat{H}\,|h\rangle = (|h\rangle\langle h| - |s\rangle\langle s|)\,|h\rangle = |h\rangle\langle h \mid h\rangle - |s\rangle\langle s \mid h\rangle = |h\rangle\,(1) - |s\rangle\,(0) = |h\rangle$$
$$\hat{H}\,|s\rangle = (|h\rangle\langle h| - |s\rangle\langle s|)\,|s\rangle = |h\rangle\langle h \mid s\rangle - |s\rangle\langle s \mid s\rangle = |h\rangle\,(0) - |s\rangle\,(1) = -|s\rangle$$

as expected

Eigenvector/eigenvalue equations say that hardness operator and hence hardness box acting on state vector

does not change states of definite hardness,

namely, |h⟩ and |s⟩

(overall minus sign does not change any measurable properties of the |s⟩ state)

as required.

Now can write matrices representing these objects(using earlier definitions/rules)

in hardness basis (called **matrix representation**)

$$|h\rangle = \begin{pmatrix} \langle h \mid h\rangle \\ \langle s \mid h\rangle \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad , \quad |s\rangle = \begin{pmatrix} \langle h \mid s\rangle \\ \langle s \mid s\rangle \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$[H] = \begin{pmatrix} \langle h|\,\hat{H}\,|h\rangle & \langle h|\,\hat{H}\,|s\rangle \\ \langle s|\,\hat{H}\,|h\rangle & \langle s|\,\hat{H}\,|s\rangle \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

(Hermitian!)

where we have used

$$\langle h|\,\hat{H}\,|h\rangle = \langle h|\,(|h\rangle\langle h| - |s\rangle\langle s|)\,|h\rangle = \langle h \mid h\rangle\langle h \mid h\rangle - \langle h \mid s\rangle\langle s \mid h\rangle = 1 + 0 = 1$$

(=1, =1, =0, =0)

Similarly, another basis(equivalent) is $|magenta\rangle = |m\rangle \quad , \quad |green\rangle = |g\rangle$

$$\hat{C}\,|g\rangle = |g\rangle \rightarrow eigenvalue = 1 \quad (by\ convention)$$
$$\hat{C}\,|m\rangle = -\,|m\rangle \rightarrow eigenvalue = -1 \quad (by\ convention)$$

|g⟩ is state with measured value of color = 1

and |m⟩ is state with measured value of color = –1.

Operator $\hat{C}$ represents entire color box.

Operator can be written in terms of its eigenvalues and projection operators —> we have

$$\hat{C} = |g\rangle\langle g| - |m\rangle\langle m|$$

They form a basis (orthonormal set) -> satisfies

$$\langle g \mid g\rangle = 1 = \langle m \mid m\rangle$$
$$\langle g \mid m\rangle = 0 = \langle m \mid g\rangle$$

Eigenvector/eigenvalue equations say that color operator

and hence color box acting on state vector does not change states of definite color,

namely, |g⟩ and |m⟩ as required.

Can write matrices representing these objects in any basis

(called matrix representation in that basis)

In color language we have

$$|g\rangle = \begin{pmatrix} \langle g \mid g\rangle \\ \langle g \mid m\rangle \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad , \quad |m\rangle = \begin{pmatrix} \langle m \mid g\rangle \\ \langle m \mid m\rangle \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$[C] = \begin{pmatrix} \langle g|\hat{C}|g\rangle & \langle g|\hat{C}|m\rangle \\ \langle m|\hat{C}|g\rangle & \langle m|\hat{C}|m\rangle \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

where we have used $\langle g|\hat{C}|m\rangle = -\langle g \mid m\rangle = 0$

Now write |g⟩ and |m⟩ vectors in terms of |h⟩,|s⟩ vectors.

Always possible since |h⟩,|s⟩ is basis and |g⟩,|m⟩ just other vectors in same space(world).

Just principle of SUPERPOSITION mentioned earlier, Write

$$|g\rangle = a\,|h\rangle + b\,|s\rangle$$
$$|m\rangle = c\,|h\rangle + d\,|s\rangle$$

Normalization: states must be normalized to 1(assume a, b, c, d are real for simplicity)

$$\langle g \mid g\rangle = 1 = (a\,\langle h| + b\,\langle s|)\,(a\,|h\rangle + b\,|s\rangle) = a^2 + b^2$$
$$\langle m \mid m\rangle = 1 = (c\,\langle h| + d\,\langle s|)\,(c\,|h\rangle + d\,|s\rangle) = c^2 + d^2$$

Orthogonality: the states must be orthogonal (they are a basis)

$$\langle g \mid m\rangle = 0 = (a\,\langle h| + b\,\langle s|)\,(c\,|h\rangle + d\,|s\rangle) = ac + bd$$

One possible solution to equations (the color states) is

$$a = b = c = -d = \frac{1}{\sqrt{2}}$$

$$|g\rangle = \frac{1}{\sqrt{2}}|h\rangle + \frac{1}{\sqrt{2}}|s\rangle \quad , \quad |m\rangle = \frac{1}{\sqrt{2}}|h\rangle - \frac{1}{\sqrt{2}}|s\rangle$$

Similarly, can express |h⟩,|s⟩ vectors in terms |g⟩, |m⟩ basis to get

$$|h\rangle = \frac{1}{\sqrt{2}}|g\rangle + \frac{1}{\sqrt{2}}|m\rangle \quad , \quad |s\rangle = \frac{1}{\sqrt{2}}|g\rangle - \frac{1}{\sqrt{2}}|m\rangle$$

Sums/differences of vectors = **superpositions** of physical states.

States of definite color = **superpositions** of hardness states.

States of definite hardness = **superpositions** of color states.

Hardness/color operators are **incompatible** observables

in the sense that states of definite hardness (eigenvectors of hardness operator)

apparently have no definite color value (not eigenvectors of color operator)

and vice versa.

Since color and hardness are **incompatible —-> operators do not commute**.

Can see by determining matrix for $\hat{H}$ in color basis and then computing commutator. We have

$$[H] = \begin{pmatrix} \langle g|\hat{H}|g\rangle & \langle g|\hat{H}|m\rangle \\ \langle m|\hat{H}|g\rangle & \langle m|\hat{H}|m\rangle \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

where we have used

to do algebra with operators one must write them in same basis

$$\langle g|\hat{H}|m\rangle = \left(\frac{1}{\sqrt{2}}\langle h| + \frac{1}{\sqrt{2}}\langle s|\right)\hat{H}\left(\frac{1}{\sqrt{2}}|h\rangle - \frac{1}{\sqrt{2}}|s\rangle\right)$$

$$= \frac{1}{2}\left(\langle h|\hat{H}|h\rangle - \langle h|\hat{H}|s\rangle + \langle s|\hat{H}|h\rangle - \langle s|\hat{H}|s\rangle\right)$$

$$= \frac{1}{2}\left(\langle h\mid h\rangle + \langle h\mid s\rangle + \langle s\mid h\rangle + \langle s\mid s\rangle\right) = \frac{1}{2}(1+0+0+1) = 1$$ and so on

Then

$$\left[\hat{C},\hat{H}\right] = \hat{C}\hat{H} - \hat{H}\hat{C} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$= 2\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \neq 0$$

These descriptions of color/hardness = 2-dimensional vector space.

Now for the **final part** of the 2nd postulate:

if system is in state $|\psi\rangle$ and measure observable $\hat{B}$ (and $|\psi\rangle$ **NOT** an eigenstate of $\hat{B}$),

then the **ONLY possible results** of measurement are the eigenvalues of $\hat{B}$

that is, the set $\{b_k\}$

Read that statement again! It is a truly amazing statement!!!!!

Nothing else will ever be measured but the eigenvalues!!!!!

(C) 3rd Postulate: Dynamics of state vectors

There exist **"deterministic"** laws (same as classical world rules)

about how a state vector of any system changes with time.

Every state vector representing real physical system must have length = 1

—> changes of state vectors dictated by dynamical laws

(called **Schrodinger** equation or the **time-development** equation)

are changes of **"direction"** (and never of **"length"**) of ket vectors.

We define a "time evolution or development" operator $\hat{U}$

that governs how state vector changes in time by the relationship

$$|A, t + \Delta t\rangle = \hat{U}(\Delta t)\,|A, t\rangle$$

—> state vector at time t + Δt given

by time evolution operator $\hat{U}$ operating on state vector at time t.

In general, ket labels(which contain whatever we know(have measured) about state)

are the **only thing that changes**

Time evolution operator = unitary operator(because length does not change) $\hat{U}$ which means

if $\hat{U}\hat{U}^{-1} = \hat{I}$ or $\hat{U}^{-1}$ is the inverse of $\hat{U}$ then the Hermitian conjugate $\hat{U}^{\dagger} = \hat{U}^{-1}$

Time evolution operator is related to energy operator $\hat{U}(t) = e^{i\hat{E}t/\hbar}$ more later

(D) 4th Postulate - Connection with Experiment/Measurements

Have said so far :

particular physical state whose state vector is an eigenvector, with eigenvalue α ,

of the operator associated with a measurable property

will **"have"** value α for that property

and that a measurement of that property, carried out on the system

which happens to be in that state,

will produce result α with certainty(probability = 1)

and that if system is **not** in an eigenvector of the observable being measured,

then one can **only** measure one of its corresponding **eigenvalues**.

Need much more than that to deal with real-world experiments.

What if we measure a certain property of physical system

at a moment when the state vector of system

does not happen to be an eigenvector of that measurement property operator?

(this is the case most of the time).

i.e., what if we measure the color of a hard electron(remember it is a superposition of being green and being magenta)?

What happens then?

All our earlier assumptions/postulates are no help here. We need a **new** assumption/postulate.

Suppose we have system in state $|\psi\rangle$,

and carry out measurement of value of property (observable) B

associated with operator $\hat{B}$.

Assume eigenvectors of $\hat{B}$ are states $|b_i\rangle$ which means that

$$\hat{B}\,|b_i\rangle = b_i\,|b_i\rangle \quad , \quad i = 1, 2, 3, 4, ...$$

where the b_i are the corresponding eigenvalues.

Quantum theory —> outcome of measurement is strictly a matter of "probability".

Quantum theory **stipulates** that the probability that

outcome of measurement of $\hat{B}$ on state $|\psi\rangle$ (not an eigenvector)

will yield result b_i

(remember only possible results are eigenvalues of $\hat{B}$ no matter what state we are in),

is equal to

$$|\langle b_i \mid \psi\rangle|^2$$

or absolute-square of **corresponding** ket vector component!

This postulate means the following:

(a) Probability as so defined is always ≤ 1 (must be to make sense),

which results from requirement that allowable states have length = 1.

That was reason for earlier imposing the normalization requirement.

(b) If $|\psi\rangle = |b_i\rangle$ (= eigenvector),

then probability to measure b_i is

$$\text{probability } = |\langle b_i \mid b_i\rangle|^2 = 1$$

and for any other eigenvalue b_k $k \neq i$

$$\text{probability } = |\langle b_k \mid b_i\rangle|^2 = 0 \quad , \quad k \neq i$$

(c) Probability that green electron is found to be soft during hardness measurement = 1/2.

State being measured is

$$|g\rangle = \frac{1}{\sqrt{2}}|h\rangle + \frac{1}{\sqrt{2}}|s\rangle$$

$$prob(soft|green) = |\langle s \mid g\rangle|^2 = \left|\langle s|\left(\frac{1}{\sqrt{2}}|h\rangle + \frac{1}{\sqrt{2}}|s\rangle\right)\right|^2$$

$$= \left|\frac{1}{\sqrt{2}}(\langle s \mid h\rangle + \langle s \mid s\rangle\right|^2 = \left|\frac{1}{\sqrt{2}}(0+1)\right|^2 = \frac{1}{2}$$

(d) Similarly, probability that green electron is hard during hardness measurement = 1/2.

Probability that hard electron green during color measurement = 1/2.

Probability that hard electron is magenta during a color measurement = 1/2 and so on.

The new formalism(4 Postulates) correctly predicts

all experimental results for hardness and color measurements(experiments).

It is important to realize that we cannot say anything definite about color if system in hardness state and vice versa.

Can then only make probability statements.

BEFORE we measure the color of electron, the electron DOES NOT have a color, according to quantum theory!

Our information about electron color is only set of probabilities before measurement!!!!

But **all of your experience** says

that objects **have** values for measured quantities

before they are measured,

i.e., your experience tells you

that electron **has** color even if we do not measure it.

That is your view (standard classical view) about what is real and what is not real.

Quantum theory says you are **wrong** in both cases!!

If you assume otherwise, then QM would not work...**but it does!!!**

Eventually we will devise real experiments to show that quantum theory is correct

and your classical views about reality are incorrect!! Think about what I just said!

Must also **devise theory** to show

why it seems to be this way for electrons

but does not seem to be true for macroscopic objects.

Finally, let me state the last and most controversial postulate.

(**E) Collapse - Measurements are always repeatable. Seems like innocuous statement!**

Once a measurement is carried out and a result obtained for some observable,

the state of system must be such as to guarantee (probability = 1)

that if **same** measurement is repeated,

the exact **same** result is obtained.

Since systems evolve in time,

this can only be true if the 2nd measurement follows the 1st **instantaneously**

or within such a small time that the system **does not** have a chance to evolve.

What does this mean about state vector of a measured (after measurement) system?

One view - something **must happen** to state vector when measurement occurs.

If measurement of observable $\hat{O}$ carried out on system S,

and if outcome of measurement is value o_q (one of its eigenvalues)

then, whatever state vector of S was before measurement of $\hat{O}$

The **only way to guarantee** (probability = 1)

that another measurement of $\hat{O}$ will give the same result

is that state vector of system S after the measurement

necessarily must now be the eigenvector of $\hat{O}$ with eigenvalue o_q .

This must be true according to Postulate #2 and **all postulates must be consistent**.

Thus, in **this view**,

the effect of measuring any observable

must necessarily be to **"change"** state vector of measured system,

to "**COLLAPSE"** it, i.e.,

to make it **"JUMP"** from whatever state it was in prior to measurement

into an eigenvector of the just measured observable operator.

This is called **collapse of state vector or reduction of state vector.**

It says that state vector changes(**discontinuously**)

during measurement **from** representing a range of possibilities

(superposition of all possible states)

to **definite** state or only one possible outcome.

Which particular eigenvector it gets changed into

is determined by the outcome of measurement

and cannot be known until then!!!!!!.

It cannot be predicted!!!!

Remember we only can predict probabilities!

Since the outcome(earlier assumption) is matter of probability,

it is **at this point and at no other point** in the discussion,

that an element of "**pure chance**" enters into time evolution of the state vector

and **determinism goes out window**.

The postulates are correct!!

The time evolution of system is **continuous and deterministic** between measurements,

and **discontinuous and probabilistic**(random) during measurements.

The last(5th) postulate is **very controversial.**

We will discuss this controversy in detail later.

Those are principles(postulates) of quantum theory.

They are **most precise** mechanism

for predicting outcomes of experiments on physical systems ever devised.

No exceptions to them have ever been discovered (108 years).

NOBODY expects any.

Now let us use these principles to **study QM**.

We will **see** how quantum theory makes predictions

and **how** the strange results of various experiments are **accounted for** by the theory.

So now we enter a strange world

where quantum systems behave in rather mysterious and non-intuitive ways.

Remember that any behavior we predict and observe

will just be a **consequence** of the 5 postulates just stated.

That is how theoretical physics works!

Stern-Gerlach Experiments

Now we repeat everything using electrons and Stern-Gerlach devices (real world devices)

—> we reinforce knowledge learned and expand our understanding.

Discussion involves the results of a classic experiment first done in 1922.

Stern-Gerlach (S-G) experiment studies how electrons behave in magnetic fields.

Illustrates how electrons behave in another context(beyond color/hardness)

and it thereby further develops the idea of quantum states

and the assumptions of quantum theory

and allows for later discussion of modern quantum experiments

and of ideas which seem so paradoxical to classical physicists.

Remember

Seemingly paradoxical nature of light/electrons has been called **wave-particle duality**

i.e., they could appear as wave in one situation

and a stream of particles in another.

To understand this,

we will need to discard the classical descriptions of waves or particles

(maybe there are no such things quantum mechanically)

and develop new set of rules and concepts to cover the strange microworld.

There will be no wave-particle duality in QM in the classical sense of the words!

We will find that:

Electrons can exhibit wave-like properties during measurements

if that is how we set up experiment

or if **we are asking** questions that require “wave-type” answers.

Electrons can exhibit particle-like properties during measurements

if that is how we set up experiment

or if **we are asking** questions that require “particle-type” answers..

The **context** of experiment will **determine** the experimental results.

The quantum theory of microworld is **contextual** in nature as we shall see.

Quantum mechanics will correctly answer whatever question you ask and the answer will be in context of the question!

The Classic Stern-Gerlach Experiment


QM -> electrons are fundamental particles, have - charge and ~1/2000 proton mass.

Found isolated or inside atoms where held to +charged nucleus by electric forces.

If atoms subjected to large electric field, then one can ionize(remove) electron from atoms.

Old style CRT tubes are large glass containers with a vacuum inside

Electron gun -> uniform beam of electrons



Electrons behave like tiny bar magnets(they have a magnetic moment) when sent into a magnetic field,

i.e., send some tiny bar magnets into non-uniform

magnetic field (stronger at top than bottom of field region)

then field both deflects path of magnets


and aligns(N-S axis) magnets with fields.

Assume that S-G magnet exerts similar magnetic force


on electrons (which have magnetic moment due to spin)

when they are passing between the poles

and that force will deflect the electron's path



non-uniform magnetic field (red arrows) generated between magnets



Hypothetical experiment to see how much deflection takes place when we pass an electron beam between the poles of a magnet.

Detect deflected electrons **outside** field region;

Can detect **single** electrons.

expecting to see spread —> #4

but see only 2 spots—> #5
i.e., only TWO experimental values

Run experiment - observe two things:

(1) No pattern determining which way electrons are deflected;

Either up or down - apparently at random.

(2) Each electron deflected upward or downward, by **fixed** amount (final deflection angle).

2nd result was very surprising to classical physicists:

i.e., amount of deflection is same for each electron.

If electron were truly acting like a tiny magnet (classical picture),

then expect magnets to be pointing in random directions when they enter S-G field.

Consequently, amount of deflection,

which depends on initial orientation of electron's magnet

would be slightly different for each.

End result = range(in space) of detected deflection angles **not** just two fixed deflections.

Classical picture —> spread but experiment sees only 2 spots !

Can interpret the results as follows:

Assume electrons have an internal property (UP/DOWN)

which determines which way they are oriented.

As they emerge from electron gun,

up and down types produced at random(equal numbers)

—> get two equal-sized sets of sorted electrons.

Define electrons deflected up as UP state electrons = $|U\rangle$

and those deflected down as DOWN state electrons = $|D\rangle$.


NOTE: Any assumption that state of electron is determined by an internal property will lead to problems later.

Assume |U⟩/|D⟩ state describes an electron in this context(UP/DOWN language):


As they are moving along top/bottom path through S-G magnet.

some internal state property (U/D) of electron determines which path they follow,

and observing the path of electron is **only** way we can measure that state property.



Can check beam properties with meter



Presumably electron gun producing electron beam for the experiment

is designed to have electrons randomly emerge in either |U⟩ or |D⟩ state.

These electrons then pass through the poles of an S-G magnet

and are sorted by being deflected **according** to the state(ket) labels.


Now we ask if the experimental results

are genuinely measuring

some state property(U/D) of electrons,

or if magnet is simply randomly deflecting

them one way or other.




Answer this question by modification of experiment as shown above

Electron detector removed/replaced

by further pair of S-G magnets

arranged so that electrons passing out of first magnet

pass through only one of two additional magnets



Results are conclusive.

(corresponds to earlier repeatability measurements)

Electrons that emerge along

UP channel of first magnet

then pass through topmost second magnet

and **all** emerge from that magnet's UP channel.

None are deflected downward

(there would be some down if each magnet was randomly deflecting electrons).

Similarly,

DOWN electrons emerging from first magnet

all subsequently deflected down by second magnet.

Second magnets(can even add more)

confirm sorting of first magnet.

Results give the **impression**

that S-G magnets are measuring some **state property** of the electrons.

Turning the Experiments Around

Up to now, we have been using S-G magnets oriented vertically

so they are deflecting electrons **upward or downward**.

The magnets can be turned through 90○

so that they now deflect electrons **right or left**.

In fact, S-G magnets can be oriented at any angle,

but **only need** (UP,DOWN) and (LEFT, RIGHT) for our present discussion.

Results of experiment with S-G magnet turned horizontally

are exactly the same as previous experiments

but now in reference to new orientation of magnets.

Half of electrons deflected to right,

and half to left.

No obvious pattern that predicts which electron will go which way.

Same arguments as earlier suggest

that there are two possible states in this context for the electron $|R\rangle$ and $|L\rangle$,

and the magnet sorts them (RIGHT/LEFT language).

Thus, electrons have second state property (R/L)

that determines which way they are deflected by horizontal magnets.

Adding two further magnets, also arranged horizontally,

to check out the electrons in either deflected beam confirms this as before.

Results are not surprising. |R⟩ electrons from first magnet

are deflected only to right by second magnet

and |L⟩ electrons are deflected to left again by second magnet.

The similarities to Hardness and Color are striking! **By Design on my part!!**

For the physicist, the next step is

to see if |U⟩ and |D⟩ states are linked (**correlated**) to |R⟩ and |L⟩ states in any way.

—> “**determining**” whether state properties connected(correlated)?


Easy to check by constructing experiment

that uses an (UP,DOWN) S-G magnet

with two (LEFT,RIGHT) magnets


so that electrons in UP and DOWN channels

of first magnet are tested to see if either |L⟩ or |R⟩.



symbols mean |U/D property>|R/L property>

In figure used symbol |?⟩ -> not sure state electron in, i.e., property has not been measured yet!



Results of experiment are very interesting.

$|D\rangle$ electron passing into (LEFT, RIGHT) magnet

comes out via either channel;

and also the same for $|U\rangle$ electron (same as happened in color/hardness boxes).

Now if true,

it appears that we are dealing with 4 different combinations of electron states

that are determined by the 2 state properties.

An electron in state $|U\rangle$ could also be in either state $|L\rangle$ or state $|R\rangle$, etc

So the possible combinations are

$$|U\rangle\,|R\rangle \quad |U\rangle\,|L\rangle \quad |D\rangle\,|R\rangle \quad |D\rangle\,|L\rangle$$

Suppose we say: Electron gun is producing equal numbers of each type.

Then, combination of 2 differently oriented (perpendicular) magnets


then would sort out these electrons as shown above. —> equal numbers in each group!

When electrons arrive at first magnet,

no way of knowing either

(UP,DOWN) or (LEFT,RIGHT) state,

hence |?⟩|?⟩.



First magnet sorts into |U⟩ or |D⟩,


but tells us nothing about |R⟩ or |L⟩ state.

Final pair of magnets completes sorting

-> four piles of distinct state combinations, with roughly equal numbers of in each pile.

Things Get More Puzzling

Let us make an extension to the experiment by adding two more magnets to check results



we are measuring electrons before they go into final magnet

Results of experiment are truly remarkable(as they were in color-hardness experiments).

Electrons from beam labelled $|U\rangle|R\rangle$

we have just done two measurements

(thought to contain electrons only in $|U\rangle$ state)

classical view

now pass through last magnet

and emerge from **either** the UP- or DOWN-channel!

Seems some |D⟩ state electrons got mixed with beam that we thought was pure |U⟩.

But that cannot be the explanation since we find no extra electrons in beam.

Results show each of emerging beams contains roughly half of electrons.

Better explanation

(LEFT,RIGHT) magnet changed

the state of some of electrons passing through.

All electrons arriving at magnet are in |U⟩ state,

but perhaps after passing through (LEFT,RIGHT) magnet,

a few flipped into |D⟩ state.


Get Answer from another new experiment(only experiment can give answers!):

Experiment starts with pure beam of |U⟩ electrons

(select UP channel of (UP,DOWN) S-G magnet)

passing through (LEFT,RIGHT) magnet,

producing two beams(half of electrons each).



Now move another (UP,DOWN) magnet very close

-> both beams now pass through final magnet (we are **not** measuring electrons before they go into final magnet).

Not difficult - deflections small -> big distance is needed to separate the beams by measurable amounts.

Before we discuss what actually happens, let us summarize our thinking:

1. Going through (UP,DOWN) magnet splits 1 beam into 2 beams.

2. Implies electron in 2 possible states and magnet is sorting them.

3. Called these states $|U\rangle$ and $|D\rangle$.

4. Similarly, horizontal magnet sorts electrons into $|R\rangle$ and $|L\rangle$ states.

5. —> four combinations of electron states: $|U\rangle$ and $|R\rangle$, $|U\rangle$ and $|L\rangle$, and so on.

6. Passing beam of electrons that is only $|U\rangle|R\rangle$
(came from UP channel and RIGHT channel in that order)
into another (UP,DOWN) magnet divides beam into 2 again.

7. Conclusion from experiment is that
passing $|U\rangle$ beam through (LEFT,RIGHT) magnet
flips (UP,DOWN) state of some of electrons.

If so - which electrons get switched? Is there any determining property?

Remember hardness/color experiments. Sound familiar!

Based on (1-7),

we predict that allowing both beams from (LEFT,RIGHT) magnet

to pass through single (UP,DOWN) magnet produces

same result as having an (UP,DOWN) magnet on each beam.

Should get two beams emerging from single (UP,DOWN) magnet


as magnet has flipped state of some of electrons.

Of course we get an unexpected result!

Using 1 magnet to catch both emerging beams

produces just single beam of pure |U⟩

electrons (and for lower beam also).



We have not measured any effects of R/L magnet so it has no effect!!!!!

Conclusion is clear.

If beams from (LEFT,RIGHT) magnet passed into separate (UP,DOWN) magnets,

then $|U\rangle/|D\rangle$ state of electrons is modified.

However, if both beams from (LEFT,RIGHT) magnet pass through same (UP,DOWN),

then no state flip.

Original state of electrons that entered (LEFT,RIGHT) magnet preserved.

Remember hard/soft electrons in 2-path experiment (all stayed magenta).

Very puzzling until explained.

Up to now, everything we have said about electron states

and the way electrons are deflected (sorted) by S-G magnets

could be a simple extension to classical ideas about electrons.

Now, in this experiment

we are starting to see that the states have a quantum nature

—> behave in rather different (non-classical) way.

Let us try to retain some common sense

We speculate that the flipping of an electron's state

is a process that needs to travel a certain distance(time) in order to happen.


Moving (UP,DOWN) S-G magnet closer

—-> has not given enough opportunity for flip to happen.

Can kill idea and any similar lines of thought by making simple modification to experiment.

Insert small metal plate sufficient to block LEFT channel of (L,R) magnet

—> stop electrons in that channel.



Have not moved magnet any further away,

so all |D⟩|R⟩ electrons will presumably,

if guess about a distance

being needed is correct,

stay in |D⟩ state

and come out of second magnet

along bottom channel.

Wrong again!

Modification adds another puzzle.

Blocking LEFT channel restores flipping (UP,DOWN) state.

As experiment doesn't alter distance travelled by electrons in RIGHT channel,

have eliminated argument based on flipping needing certain distance to work.


Can turn flipping on or off

by blocking one of paths and doing nothing to distance.


(Blocking RIGHT channel gives same type of result.)

Same as happened with color/hardness experiments


Summarize results of experiments below.



(a) Blocking RIGHT channel produces mixture of |U⟩/|D⟩ states in electrons that pass through LEFT channel.



(b) Blocking LEFT channel produces mixture of |U⟩/|D⟩ states in electrons that pass through RIGHT channel.



(c) Having both LEFT and RIGHT channels open produces only |D⟩ state electrons.

What Does It All Mean?

Started with idea that electrons possess certain state property

that determines path through S-G magnet.

Some electrons start in |U⟩ state and some in |D⟩ state,

and when electrons are formed into beam,

|U⟩ and |D⟩ electrons randomly distributed, so can't tell which type coming next.

Crucially,

we have been **assuming** that the state of electron is

fully determined **before** it enters any magnet in its path.

This is the assumption behind the classical idea of a state

i.e., that a measurement simply reveals what is already there.

Results of these experiments completely undermine this idea.

1. Passing beam |D⟩ electrons through (LEFT,RIGHT) magnet separates them into |D⟩|L⟩ and |D⟩|R⟩ states(equal numbers)

2. Passing |D⟩ |L⟩ and |D⟩ |R⟩ electrons into separate (UP,DOWN) magnets produces both |U⟩ and |D⟩ electrons at each magnet. |D⟩ state does not always survive passing through (LEFT,RIGHT) magnet.

3. Passing |D⟩ |L⟩ and |D⟩ |R⟩ electrons into same (UP,DOWN) magnet produces pure |D⟩ beam. |D⟩ state is now preserved.

4. Undermines thought expressed in (1) that we can specify (UP,DOWN) and (LEFT,RIGHT) states at same time. No |D⟩ |L⟩ and |D⟩ |R⟩ states exist, just |U ⟩/|D⟩ OR |R⟩/|L⟩ states.

5. **Suggestion:** distance travelled by electrons on passage through magnet causes effects contradicted by experimental results produced by blocking one of beams.

6. Blocking left- or right-hand beam through (LEFT,RIGHT) magnet separately before reaching same single (UP,DOWN) magnet as in point (3) results in some electrons going up and some going down.

7. —> Nature of electron's state depends on **context of experiment - the question asked!!!**.

Another point makes things even stranger.

If we block LEFT channel,

then electrons passing along RIGHT channel into (UP,DOWN) magnet

emerge either |U⟩ or |D⟩.

However, if they passed along RIGHT channel,

how can they have known that LEFT channel closed? **(poor question)**

Another way:

if suddenly open up LEFT channel,

add more electrons passing into (UP,DOWN) magnet

those that would have gone through RIGHT channel anyway

and those that were blocked in LEFT channel.

Suddenly **all** electrons are now in |D⟩ state.

Remember magenta electrons coming out all magenta!

No results depend on intensity of beam.

If one electron present in apparatus at a time, all experiments => same results.

Disposes of idea that electrons are interacting with each other.

No way that electron passing through one channel

could be influenced by other channel being blocked,

unless there is another electron in that channel at same time to mediate influence.

Clearly, as experiment gives same result with low-intensity beam, **that idea can't work either.**

Results can made a coherent whole as follows.

Consider what information we can obtain from **each** experiment.

When one channel through (LEFT,RIGHT) magnet blocked,

clear that any electron emerging from experiment

must have passed through open channel.

With both channels open,

cannot tell which path the electrons followed through (LEFT,RIGHT) magnet.

Cannot just watch them go past.

Any method used to **determine** which path electrons take **= blocking the path**.

Similarities to old experiments.

Again context of whole experiment proves crucial.

Evidently, knowing electron either $|L\rangle$ or $|R\rangle$ state

prevents us from saying it is in $|U\rangle$ or $|D\rangle$ state.

Having 1 path blocked after 2nd magnet

-> electron entering (UP,DOWN) magnet

clearly either |L⟩ or |R⟩ state


-> lose any idea of being |U⟩ or |D⟩.

Both paths open

-> no information from experiment

tells us about |L⟩/|R⟩ state of electrons.

Then, can retain some information about |U⟩/|D⟩ state.



Interpretation is not required by results of experiments discussed so far.

Look at other quantum experiments to see consistency of approach.

Earlier color/hardness showed getting different results

depended on not being able to tell which path electrons were using.

Here can tell if it is |U⟩/|D⟩ **as long as cannot tell** if is |L⟩/|R⟩.

Results are showing us something important about nature of quantum state.

Quantum States/Postulates - Real Meaning of Formalism

Now we set up the formal axiomatic structure of quantum theory

and along way review most of stuff that we have already talked about.

Hopefully, you understand QM after this pass!

Must redevelop theoretical machinery of quantum physics.

Must build on experimental insight of present discussions

=> valid statements about nature of quantum objects

(no matter how confusing that nature seems to be).

Where Are We Now?

Experiment —> reality slightly out of focus,

until specific experiment forces picture to sharpen into one possibility or another.

Is being **"out of focus"** only an issue of the theory

and not a reflection of what is actually happening?

Maybe future experiments will expose physical variables (not included so far)

-> advanced theory that resolves what appears to be paradoxical

-> called so-called Hidden Variable theories

Maybe current experiments are saying something **genuine** about nature of reality.

Problem then is our **expectations** about reality.

Our view of reality

-> Experience centered on macroworld (live in).

Rocks follow single path.

Common sense understanding colors view of reality, irrespective of scale

No guarantee that such applies outside the macroworld.

Experiments with photons/electrons —> they are **not** tiny rocks.

Consequently, our challenge:

how a rock (made up of microworld objects) can behave in “common sense” way,

given the underlying strangeness of the quantum world.

Need to completely specify scheme for describing our experimental results in consistent way

—> some measure of predictability.

Need to develop quantum theory from 1st principles that can apply everywhere.

Covered ideas earlier and now expand on those discussions

—> all aspects of full theory will be clearly delineated.

Describing Quantum Systems

A classical state cannot describe experiments in the quantum world.

A classical state = a list —> values of various physical properties,

this cannot apply to a photon in an interference experiment

which is “apparently” traveling on 2 paths at same time. (apparently = “seems like”)

Any physical property determining the direction of electron through (S-G) experiment

is influenced by exact details of how experiment was set up(**the context**),


—> runs contrary to classical idea that experiments

just reveal what is already there.

How can we construct a quantum state to replace normal classical description of system.

It is important that certain basic characteristics are designed in from the beginning.


1. Inherent **randomness** must be represented,
i.e., description of photon arriving at half-silvered mirror
must allow photon to have equal chance of being transmitted
or reflected, without stating that it will definitely do one or other.



2. **Contextuality** of quantum behavior must be incorporated.
Experimental results are influenced by overall setup of devices.
If photon detector is placed beyond the half-silvered mirror, then
photon will either reflect or transmit. If no detector present, then
results -> that photon “explores” both possibilities.

3. Quantum systems exist in a mixed state that combines classical states in impossible ways as we will see(have seen, (e.g., simultaneously reflecting and transmitting at half-silvered mirror), i.e., that combine classical states which are **incompatible** with each other!

Figure summarizes what needs to be achieved in quantum description.



LHS = system(electron),

in quantum state = $|\phi\rangle$ (a ket).

Electron interacts with measuring apparatus, M,

-> one of several possibilities can occur,

each with different probability.

Example:

$|U\rangle$ state electron interacting with (LEFT,RIGHT) S-G magnet

-> emerging in $|L\rangle$ or $|R\rangle$ state.

We take most basic direct approach: **Will enable us to learn important things!**

Assume the initial state takes the following form:

$$|\phi\rangle = p_1 |A\rangle + p_2 |B\rangle + p_3 |C\rangle + p_4 |D\rangle + ...$$

where numbers p_1,.....,p_4,.... which represent(do not know how yet) somehow the

probability that electron would end up in each state |A⟩, |B⟩, etc.

Not saying that numbers are probabilities; just related to probabilities(to be determined).

Attractive formulation —> already catches some of flavor of quantum behavior.

Seems to say that quantum state |ϕ⟩ is made up of all possibilities |A⟩, |B⟩, etc,

which may subsequently come about.


Example: S-G experiment, write

$$|U\rangle = p_1 |L\rangle + p_2 |R\rangle$$

to represent initial state of electron

when approaching L-R magnet

(next measurement)



(shown in figure)

After electron passed through magnet,

no longer appropriate

to describe it by state |U⟩:

since now either state |L⟩ or |R⟩,

initial description **seems** to have

“collapsed” (after measurement)

into one(we observed) of two alternatives

that it is “composed” of.

But REMEMBER other state is still somewhere

This way of expressing quantum states is similar to way probabilities are combined.

Imagine trying to calculate the average number of words per page in a book.

One way: count up number of words and divide by number of pages.

Equivalent way:

group pages into sets (each page in set had same number of words).

Average then becomes

$$\text{average number of words} = \frac{(\text{number pages}(700\text{ words}))\text{x}700+(\text{number pages}(600\text{ words}))\text{x}600+..}{\text{total number of pages}}$$

$$\text{average number of words} = \frac{(\text{number pages}(700\text{ words}))\text{x}700}{\text{total number of pages}} + \frac{(\text{number pages}(600\text{ words}))\text{x}600}{\text{total number of pages}} + ..$$

$$\text{average number of words} = (\text{Probability of } 700 \text{ words})\text{x}700 + (\text{Probability of } 600 \text{ words})\text{x}600 + ..$$

Looks like formula that is used above when different possibilities are being considered.

Given an event E_1 with probability P_1, event E_2 with probability P_2

and event E_3 with probability P_3, the rules of probability state

$$\text{Probability}(E_1 \; or \; E_2 \; or \; E_3) = P_1 + P_2 + P_3$$

If events correspond to measuring different values of physical property

(e.g., V_1, V_2, and V_3),

then average value of property after many trials is

$$\text{average value of } V = (V_1 \times P_1) + (V_2 \times P_2) + (V_3 \times P_3)$$

That looks like the assumed form of the quantum mechanical state above.

If two ideas are exactly the same, then terms p_1, p_2, p_3, etc in

$$|\phi\rangle = p_1 |A\rangle + p_2 |B\rangle + p_3 |C\rangle + p_4 |D\rangle + ...$$

would have to be **actual** probabilities, but there is a problem with this idea.

Again, we let experiment show us what is wrong!

Specific Example: Mach-Zehnder Interferometer

Have assumption(above) about

how to represent a quantum state.

Does it work in an experimental situation.


Choose a Mach-Zehnder interferometer.

Experiment **seems to need**

wave/particle descriptions of light

-> good test

of quantum state assumption.



Assume intensity of laser beam turned down(make it a real QM experiment)

—-> only one photon crossing setup at a time.

At 1st half-silvered mirror,

the single photon is **either** reflected **or** transmitted.

Our assumption (state = sum of things it could be) —> quantum state of arriving photon is

$$|\phi\rangle = a\,|T\rangle + b\,|R\rangle$$

where |T〉 represents transmitted state

and |R〉 reflected one as shown in figure.

Numbers a and b are related to probabilities

that photon transmitted or reflected, respectively.

Numbers are determined by construction of mirror.

a always related to the reflected part property

and


b always related to the transmitted part property

Part of state |ϕ〉 designated |R〉

contains information about photons

reflected at half-silvered mirror……..

Unless we have measurement device on either arm of experiment, cannot say photon has either been transmitted or reflected - MUST MEASURE TO KNOW!!



Consequently,

should not imply photon is going in a particular direction.

Alternative idea:


say that reflected photon state moves along upper arm,

but states do not move - they are just math!

Settle on saying(clearly awkward):

reflected photon property/information

meets fully silvered mirror at A



This illustrates is why we should only talk about actual measurements!!

Consequently, quantum state must change at mirror to |G⟩.

At next half-silvered mirror, state either one of transmission through or reflection up:

$$|G\rangle = b\,|L\rangle + a\,|K\rangle$$

Used SAME FACTORS, a (transmission) and b (reflection), as 1st half-silvered mirror.

Meanwhile the other property of the photon is in state |T⟩.

This property reaches mirror position B, where reflected and changes into state |H⟩.

However, assumption says must write |H⟩ in terms of |K⟩ and |L⟩

—> the possible outcomes when |H⟩ interacts with half-silvered mirror.


Figure shows - be careful when constructing representation,

from point of view of |H⟩, |L⟩ is transmitted state and |K⟩ is reflected one.

So, have to write

$$|H\rangle = b\,|K\rangle + a\,|L\rangle$$

same a and b in all three cases



Have |H⟩ and |G⟩ in terms of |K⟩ and |L⟩

—> can go back to original state |ϕ⟩

and write in terms of |K⟩ and |L⟩.

$$|\phi\rangle = a\,|T\rangle + b\,|R\rangle = a\,|H\rangle + b\,|G\rangle$$

$$= a\,(b\,|K\rangle + a\,|L\rangle) + b\,(b\,|L\rangle + a\,|K\rangle)$$

$$= ab\,|K\rangle + ab\,|K\rangle + a^2\,|L\rangle + b^2\,|L\rangle$$

$$= 2ab\,|K\rangle + \left(a^2 + b^2\right)|L\rangle$$

Have produced representation of initial photon state |ϕ⟩


$$|\phi\rangle = 2ab\,|K\rangle + (a^2 + b^2)\,|L\rangle$$

in terms of two **possible final outcomes** |K⟩ and |L⟩.

Numbers 2ab and $(a^2 + b^2)$ must represent (in some way according to assumption)

probabilities that photon will be detected at X and Y, respectively,

i.e., that will end up with state |K⟩ or state |L⟩ for single photon.



Now, the actual experimental results are as follows:

If distances along two paths in detector are equal,

then all photons passing through device are picked up at detector X;

none reach Y.

Consequently, final state of photon cannot include |L⟩.

If our formalism works, then we must have $(a^2 + b^2) = 0$ in that case —> an immediate problem.

If a and b are probabilities, then they must be positive numbers.

Since the square of positive number is positive(also the square of negative number),

no way we can obtain $(a^2 + b^2) = 0$.

Consequently, terms a and b cannot be probabilities.

Of course, could have a = b = 0

—> no physical sense,

i.e., says that half-silvered mirror does not work as observed in real experiments.

Should we abandon this approach for representing quantum systems?

Do not be too hasty.

Mathematicians looking at equation $(a^2 + b^2) = 0$ are not bothered at all.

—> realize immediately that a and b are just complex numbers.

Our assumption led to consistent representation of photon's state

on far side of Mach-Zehnder experiment —>

$$|\phi\rangle = \left(a^2 + b^2\right)|L\rangle + 2ab\,|K\rangle$$

with proviso that $|L\rangle$ state is never observed

if lengths of two arms in experiment equal.

Consequently, need $(a^2 + b^2) = 0$ —> puzzle.

But can work if using imaginary numbers.

If $a = bi$ so that $a^2 = -b^2$, then $(a^2 + b^2) = 0$.

A price to pay……..

If numbers multiplying states are complex numbers, what can they mean?

They cannot directly represent a probability,

which is a real, non-negative number.

We started with idea

that numbers used to multiply states

were related to probability

that a state emerges as result of a measurement.

Our hope —> that numbers might be probability is now

dashed by applying idea to Mach-Zehnder experiment.

Must use complex numbers

if going to represent all possible experimental situations.

Move is required by experimental results! **That is the way Physics is supposed to work!**

That is way theoretical physicist makes progress.

So the numbers cannot be probabilities

—> numbers are called probability amplitudes(as we saw earlier).

$$|\phi\rangle = a_1 |A\rangle + a_2 |B\rangle + a_3 |C\rangle + a_4 |D\rangle + ...$$

where a_1, a_2, a_3, etc are probability amplitudes for states |A⟩, |B⟩, |C⟩, etc.

How are probability amplitudes related to probabilities?

Probability obtained from an amplitude must have **all** factors i removed.

Earlier we earlier learned a procedure

that removes all factors of i from complex number:

—> multiplying number by its conjugate.

Possible interpretation(a guess at this point):

to convert probability amplitudes into probabilities,

multiply amplitude by its complex conjugate.

—> **RULE 1**: If $|\phi\rangle = a_1 |A\rangle + a_2 |B\rangle + a_3 |C\rangle + a_4 |D\rangle + ...$ then

$Prob\left(|\phi\rangle \rightarrow |A\rangle\right) = a_1^* a_1 = |a_1|^2$, $Prob\left(|\phi\rangle \rightarrow |B\rangle\right) = b_1^* b_1 = |b_1|^2$, etc and so on

—> a fundamental rule(postulate) of quantum theory as we saw earlier.

Rules(postulates) can't be proven mathematically.

Mathematics says what probability amplitudes mean

—> it is then job of physics to use this fact.

Only way of doing it is to relate

the mathematics to experimental results.

Thus, we assume rule, use it to do calculations

and then check and see if all predictions are correct.

If all works out, then the rule gets accepted.

In this case, relationship between amplitudes and probabilities

is a cornerstone of quantum theory;

the success of whole theory relies on it being correct.

Quantum theory has been around for over 100 years now and it works,

so we can regard this rule as being confirmed by experiments.

States in Stern-Gerlach(S-G) Experiment

Now we apply these ideas to the S-G experiments we discussed earlier.

If send |U⟩ state electron through (LEFT,RIGHT) magnet,

can emerge from either channel with equal probability.

Similarly, send |D⟩ state electron into (LEFT,RIGHT) magnet,

also emerges from either channel with equal probability.

Using the assumption,

we write two quantum states |U⟩ and |D⟩ in following form (remember color/hardness)

$$|U\rangle = a\,|R\rangle + b\,|L\rangle \qquad\qquad |D\rangle = c\,|R\rangle + d\,|L\rangle$$

where a, b, c, and d are probability amplitudes.

What are the values of these numbers?

Some clues to help us out.

First, a must be different from c,

and/or b must be different from d

since |U⟩ and |D⟩ are different states

although both are combinations of |L⟩ and |R⟩.

Second, if probability of emerging from either channel is same,

then Rule 1 (amplitude absolute squared = probability)

tells us solution that agrees with experiment is

$$aa^* = bb^* = cc^* = dd^* = \frac{1}{2}$$

—> correct result is:

$$|U\rangle = \frac{1}{\sqrt{2}}|R\rangle + \frac{1}{\sqrt{2}}|L\rangle \qquad |D\rangle = \frac{1}{\sqrt{2}}|R\rangle - \frac{1}{\sqrt{2}}|L\rangle$$

Remember color/hardness states.

Could switch +/– signs

—> no change in physical content (remember Do Nothing box).

Are they the correct combinations?

Can’t prove that yet.

In derivation, we have also used following:

RULE 2: NORMALIZATION - If

$$|\phi\rangle = a_1 |A\rangle + a_2 |B\rangle + a_3 |C\rangle + a_4 |D\rangle + ...$$

then from earlier definitions

$$\langle \phi \,|\, \phi \rangle = |a_1|^2 + |a_2|^2 + |a_3|^2 + |a_4|^2 + ... = 1 \qquad \textbf{(vector length)}^2$$

Works for |U⟩ state:

$$|a|^2 + |b|^2 = \frac{1}{2} + \frac{1}{2} = 1$$

Rule 2

—> total probability = sum of probability for each possibility in final state = 1.

i.e., **Something has to happen!**

If sum probabilities < 1,

then it means

there was a probability that something would happen

that was not included in list of possibilities

(states in linear combination),

which violates Rule 1.

Rule 2 puts constraint on amplitudes

—> values must say state has been **normalized** to 1,

or total probability =1

General Stern-Gerlach States

S-G experiments:

We used only positions of magnets at 90° with respect to one another - (U/D) versus (R/L);

It is clearly possible to have any orientation.

Consider beam of |U⟩ state electrons arriving at S-G magnet

with axis tilted at angle θ to vertical. (U/D) = 0° and (R/L) = 90°

Electrons emerge from magnet

along one of the two paths as before.

Call states |1⟩ and |2⟩.

In this case, different number of electrons pass down each channel,

indicating that amplitudes are not same:

In general, have (a not necessarily = to b)

$$|U\rangle = a\,|1\rangle + b\,|2\rangle \quad \text{with} \quad aa^* + bb^* = |a|^2 + |b|^2 = 1$$

A more advanced QM course is able to show(we just quote result) that:

$$|U\rangle = \cos\left(\frac{\theta}{2}\right)|1\rangle + \sin\left(\frac{\theta}{2}\right)|2\rangle \qquad |D\rangle = \sin\left(\frac{\theta}{2}\right)|1\rangle - \cos\left(\frac{\theta}{2}\right)|2\rangle$$

where θ measures angle between axis of magnet and vertical.

These states are consistent with earlier results.

i.e., send |U⟩ and |D⟩ states into (LEFT,RIGHT) magnet, θ = 90° —> θ/2 = 45°.

Now, $\sin(45^\circ) = \cos(45^\circ) = \frac{1}{\sqrt{2}}$ —> get states from earlier.

Note $\sin^2\left(\frac{\theta}{2}\right) + \cos^2\left(\frac{\theta}{2}\right) = 1$ —> Rule 2 satisfied for any θ.

Some Further Thoughts before continuing……

Summarizing: about quantum states:

Mathematical representation of initial quantum state |φ⟩,

—> an expansion (sum) over possible final quantum states as

$|\phi\rangle = a_1 |1\rangle + a_2 |2\rangle + a_3 |3\rangle + a_4 |4\rangle + \ldots = \sum_n a_n |n\rangle$

Amplitudes are collection of complex numbers

related to probability that initial state |φ⟩

will "change" into one of final states |n⟩

as result of measurement.

Rule 1 gives relationship between amplitudes and probabilities.

List of possible final states(possible measurement results)

called **basis** —> **HOME space** of expansion

So now know

how to represent amplitudes and their meaning;

What about basis states?

How can we write down $|n\rangle$ in mathematical terms?

Is there some equation or formula for $|n\rangle$?

Up to now have simply written states such as $|\phi\rangle$ in terms of basis states,

and these in turn

have been written

as a combination of a further basis.

For example we wrote $|U\rangle$ as combination of $|L\rangle$ and $|R\rangle$.

In turn $|L\rangle$ written as combination of $|U\rangle$ and $|D\rangle$ as can be done for $|R\rangle$.

System seems to lead to regression

of writing one thing in terms of another

without actually getting anywhere.

Not entirely true, however.

Remember the structure of a quantum state is a reflection

of the **contextuality** of quantum physics(from earlier).

State |U⟩ can be written as

$$|U\rangle = \frac{1}{\sqrt{2}}|R\rangle + \frac{1}{\sqrt{2}}|L\rangle$$

in context of (LEFT,RIGHT) magnet,

i.e., if U electron **about to enter** a (LEFT,RIGHT) magnet

or state |U⟩ can be written as

$$|U\rangle = \cos\left(\frac{\theta}{2}\right)|1\rangle + \sin\left(\frac{\theta}{2}\right)|2\rangle$$

in context of magnet at angle θ

i.e., if U electron **about to enter** a magnet oriented at angle θ

Each |n⟩ state in basis

represents possible result of measurement

in that context!.

We are still missing

some way of extracting

quantitative information about physics property (observable)

from state |n⟩ (to come later)

REMEMBER:
Always choose the appropriate language (basis states) before discussing an experiment or experimental results will not be understandable!

What Are Quantum States?

According to picture built up so far,

the quantum state of a system

contains a series of complex numbers related

to probability that system will collapse into new state

when a measurement takes place.

Each new state represents a possible result of measurement,

i.e., a path or quantitative value of physical variable.

However, this simple description of a quantum state

hides a number of difficulties.

If we make a measurement on a single electron,

then then result = a distinct value of physical property being measured.

But a single measurement cannot confirm a probability of finding electron with that value.

Must make same measurement many times

—> how often each specific value comes up

—> some practical difficulties,

i.e., how do we ensure a single electron is in exactly same state

every time we make a measurement?

Better off using collection of electrons(ensemble),

if can put all in same initial state,

and perform only one measurement on each.

But, if this is best way of carrying out measurement

then we must ask

what does the quantum state of a **single** system actually represent?

Given a collection of electrons

in same state prior measurement,

does quantum state describe each electron in collection

or does state only meaningfully refer to collection as a whole?

This is more than just debate over terminology:

it raises important questions about the nature of probability.

So let us expand our earlier discussion of probability.

Probabilities occur in many different ways.

Sometimes when we deal with probabilities,

there exists a physical aspect of system that reflects this,

i.e., throwing fair dice

—> each face coming up 1/6 of the time

because there are 6 faces to choose from.

However,

if have collection of balls in bag

and half are red and other half are white,

then probability of drawing red ball out of bag (no looking) is 1/2.

Here the probability is not direct reflection of property of each ball.

Probability, in this case, only exists when the balls placed in a collection.

i.e., the probability state describes only the collection and not the individual balls within it.

If quantum state = collection of systems,

different probabilities might —>

systems not quite identical —>

amplitudes do not represent genuine unpredictability inherent to system.

i.e., they may simply express existence of ignorance of situation at a deeper level.

Maybe hidden variables exist(as said earlier) and if know values, exact predictions possible.

Collection of systems have various possible values of hidden variables, just don't know.

Probability differences —> how many of each type was in collection.

However, if quantum state refers to single system,

probabilities might reflect physical nature of system

—> opens up new way of looking at reality.

If quantum state of system

represented by set of probability amplitudes,

then we are describing the state

in terms of what it can become as result of measurement.

After measurement,

one of possibilities has taken place

so system is in new state.

New state also best described

in terms of what it can become after next measurement.

Therefore continually describing systems

in terms of what they becomes or change into

never what they are.

Perhaps there is nothing more to describing

what something is then saying what it can become or what it can do.

Somehow current state of quantum system

has future implicit within it.

Once measurement taken place,

one of implicit possibilities becomes explicit.

This is a very abstract picture!

But, classical state of system is also abstract.

It is represented by a series of quantities

gotten from the physical properties of system.

Seems more real

since speed, position, mass, etc., of object

are familiar terms.

Meaning of quantum state is a philosophical question.

Provided different quantum descriptions allows calculations

that correctly predict outcome of experiment,

there is no experimental way

in which different ways of thinking about quantum state can be distinguished.

Majority of physicists take very pragmatic view.

Quantum theory works: allows calculations, do experiments, and have a career.

Deeper questions about meaning rather fruitless

(not accessible to experimental resolution).

This, however, is not me!

There is a distinct element of weirdness about all this that I want to understand.

Probability amplitudes seem detached from reality.

The normal puzzlement or the unease one feels when learning an unfamiliar subject

—> a genuine problem in QM:

How can everyday world of experience (seems deterministic)

result from underlying quantum reality

described by probability amplitudes?

More Discussion on Amplitudes

Let us look at amplitudes in detail.

—> ways amplitudes are combined.

—> so we can make significant step toward understanding the way QM works.

Amplitudes are key in QM

—> Link between theory/experiment.

Theorist uses QM to calculate amplitudes for given situation

and then experimenters set up situation and make measurements.

Experiments costly and complicated.

Normally same particle passed through sequence of measuring systems,

each extracts information about situation.

Each measuring device also has some effect on state of system.

Set of amplitudes for effect of measuring device depend on outcome of previous measurement.

Thus, need to trace amplitudes through sequence of situations.

Need know how to combine/convert amplitudes.

Basic rules for combining amplitudes

is implied by experimental results.

Rules are new postulates.

Physics is an experimental science!

RULE 3: State changes(transitions) governed by amplitudes.

Successive transitions

—> amplitudes **multiplied** together.

Two **alternatives** possible

—> probabilities **add**

if the alternatives can be distinguished in experiment.

When two alternatives cannot be distinguished,

amplitudes add and then

probability = absolute square of **total** amplitude.

Best way to understand this rule

is to see how it works **in practice**.

S-G experiments - initial state |ϕ⟩ $|\phi\rangle = a\,|U\rangle + b\,|D\rangle$

Have chosen |U⟩ and |D⟩ as basis for state expansion

since 1st part of experiment = (UP,DOWN) magnet

—> (UP,DOWN) states —> HOME space

for 1st part of experiment (possible results of measurement).

choose correct language or go to HOME space

Then, allow each beam from 1st magnet

to pass into other magnets arranged at angle θ,

so emerging states are |L′⟩ and |R′⟩ (only L/R if θ = 90)

—> expand |U⟩ and |D⟩ states in new basis |L′⟩ and |R′⟩:

$$|U\rangle = m\,|L'\rangle + n\,|R'\rangle \quad , \quad |D\rangle = p\,|L'\rangle + r\,|R'\rangle$$

At 1st magnet, $|\phi\rangle = a\,|U\rangle + b\,|D\rangle$


beam of electrons in state |ϕ⟩

divides into UP beam

containing fraction $|a|^2$ of original beam,

and DOWN beam

containing fraction $|b|^2$ of original beam.



At 2nd magnet,


$|U\rangle$ states collapse into either

$|L'\rangle$ with probability $|m|^2$

or $|R'\rangle$ with probability $|n|^2$.

Same happens to $|D\rangle$ electrons

(see figure)



Probability that

electron starting in state $|\phi\rangle$

ends up in state $|L'\rangle$

having gone

through magnet 1 is

$$Prob\left(|\phi\rangle \underset{via\ magnet\ 1}{\rightarrow} |L'\rangle\right) = |a|^2 \times |m|^2$$

Result obtained by considering

fraction of original beam that makes it through each stage.

Note, get exactly same result

if constructed amplitude governing state change from $|\phi\rangle$ to $|L'\rangle$

successive = multiply

$$amplitude\left(|\phi\rangle \underset{via\ magnet\ 1}{\rightarrow} |L'\rangle\right) = amplitude\left(|\phi\rangle \rightarrow |U\rangle\right) \times amplitude\left(|U\rangle \rightarrow |L'\rangle\right) = a \times m$$

then calculate probability by complex squaring amplitude, i.e.,

$$|a \times m|^2 = (a\,m) \times (a^*\,m^*) = aa^* \times mm^* = |a|^2 \times |m|^2$$

End result is just what Rule 3 said:

when one transition follows another,

amplitudes multiply.

So, 1st part of rule works.

Now a different question:

what would be


probability of electron

ending up in state $|L'\rangle$

if not worried about

which (L′,R′) magnet it went through?

Look back at the last figure.



$$Prob\left(|\phi\rangle \underset{via\ magnet\ 1}{\rightarrow} |L'\rangle\right) = |a|^2 \times |m|^2$$

$$Prob\left(|\phi\rangle \underset{via\ magnet\ 2}{\rightarrow} |L'\rangle\right) = |b|^2 \times |p|^2$$

So overall probability is $Prob\left(|\phi\rangle \rightarrow |L'\rangle\right) = |a|^2 \times |m|^2 + |b|^2 \times |p|^2$

Application of standard rule in probability calculations:

when have one event OR another,

probabilities add.

Rule 3 states, when two alternatives are possible,

probabilities add if alternatives can be distinguished in experiment.


Crucial part here is phrase **"can be distinguished in experiment"**.

Information about which alternative a particular system follows has to be available, EVEN IF we don't choose to use the information in the experiment.

That just leaves us with the final part of Rule 3,

which applies in situations where we **cannot** tell which alternative is happening.

Example:



Modify last experiment.

2nd magnet is pulled forward so both beams pass through.

Probability that electron state $|\phi\rangle$ ends up in state $|L'\rangle$

remembering that possibilities are now indistinguishable.

Rule 3 —> trace amplitude through each possible path individually

$$amplitude\left(|\phi\rangle \underset{top\ path}{\rightarrow} |L'\rangle\right) = a \times m \qquad amplitude\left(|\phi\rangle \underset{tbottom\ path}{\rightarrow} |L'\rangle\right) = b \times p$$

then add amplitudes together

$$amplitude\left(|\phi\rangle \underset{cannot\ tell\ which\ path}{\rightarrow} |L'\rangle\right) = a \times m + b \times p$$

Calculate probability by complex squaring total amplitude

$$Prob\left(|\phi\rangle \underset{cannot\ tell\ which\ path}{\rightarrow} |L'\rangle\right) = |a \times m + b \times p|^2$$

Final part of Rule 3 is a very important aspect of QM.

Rule that must add the amplitudes

before complex squaring

if paths indistinguishable

has **no equivalent** in normal probability calculations or classical physics.

This is a theoretical representation of what happens in any interference experiment.

Adding amplitudes before squaring

(when cannot tell which path followed)

—> two amplitudes interfere.

If both a × m and b × p are positive,

then combined amplitude bigger.

If one negative, then amplitude decreases.

In an interference experiment,

amplitudes change depending on length of path through experiment

affects sign of amplitude

—> total probability gets bigger/smaller depending on path length.

This is exactly the property we are looking for to explain interference experiments.(see later).

It is all about a Change of Basis or a Change of Language

Alternative indistinguishable path probability calculation.

Now use mathematical manipulation.

Do in detail.

Learn/practice

mathematical/algebraic procedures.

Idea that states

can be expanded over any basis is an important aspect of QM

—> part of way that **contextuality** of QM

is reflected in theory.

When we expand state,

we always choose that basis


useful in describing measurement **about to take place**

—> called going to HOME space(before measurement).

Every quantum state can be expanded in many ways,

depending on what experiment is involved.

Illustrate this idea via 2nd look at indistinguishable path experiment .



1st thing electrons in I$\phi\rangle$ come to

is (UP,DOWN) S-G magnet.

Expand state in I$U\rangle$ and I$D\rangle$ (the basis).

$$|\phi\rangle = a\,|U\rangle + b\,|D\rangle \quad \text{Expansion 1}$$

Electrons then hit (L′,R′) magnet,

sensible to expand I$U\rangle$ and I$D\rangle$ states in I$L'\rangle$ and I$R'\rangle$.

$$|U\rangle = m\,|L'\rangle + n\,|R'\rangle$$

$$|D\rangle = p\,|L'\rangle + q\,|R'\rangle$$

Expansion 2

Initial state is |ϕ⟩ and final state interested in is |L′⟩ - do not care about in-between!

Need expansion of |ϕ⟩ in |L′⟩ and |R′⟩.

Get expansion by pulling together information already have.

Plug Expansion 2 into Expansion 1.

Do details,

$$|\phi\rangle = a\,|U\rangle + b\,|D\rangle = a\,[m\,|L'\rangle + n\,|R'\rangle] + b\,[p\,|L'\rangle + q\,|R'\rangle]$$

$$= (am + bp)\,|L'\rangle + (an + bq)\,|R'\rangle$$

Result very interesting, look carefully.

$$|\phi\rangle = (am + bp)\,|L'\rangle + (an + bq)\,|R'\rangle$$

Expansion

—> amplitude for |ϕ⟩ collapse to |L′⟩ by end of experiment

(also —> amplitude for |ϕ⟩ collapse to |R′⟩).

Look closely,

amplitude (am + bp) = same result calculated using Rule 3 earlier.

If paths distinguishable,

Expansion 1 no longer valid.

Instead $|\phi\rangle$ turns into one of $|U\rangle$ or $|D\rangle$,

whereas Expansion 1 assuming both valid at same time

i.e., state has not collapsed so cannot tell which path is happening.

Same quantum state can be expanded in many ways,

depending on experiment.

Also possible to switch from one basis to another with bit of algebra.

Expanding the Dirac Language to make things easier……..

Have mainly used KETS $|..\rangle$ so now reintroduce BRAS $\langle..|$

Just an alternative way of representing/expanding state

using complex conjugates of amplitudes

rather than amplitudes themselves

$$\langle\phi| = a_1^* \langle 1| + a_2^* \langle 2| + a_3^* \langle 3| + ... + a_n^* \langle n|$$

KETS and BRAS contain same information.

RULE 4: If system starts in ⏐ϕ⟩ and ends in ⏐ψ⟩,

then amplitude for transition is

calculated by taking bra of final state

and acting on ket of initial state

(mathematically = “Braket” = inner product <—> use × for the moment)

$$amplitude\left(|\phi\rangle \rightarrow |\psi\rangle\right) = \langle\psi| \times |\phi\rangle$$

Start in $|\phi\rangle = a\,|U\rangle + b\,|D\rangle$

Rule 4 —> amplitude for collapse of ⏐ϕ⟩ into ⏐U⟩ is

$$amplitude\left(|\phi\rangle \rightarrow |U\rangle\right) = \langle U| \times |\phi\rangle = \langle U| \times \left(a\,|U\rangle + b\,|D\rangle\right)$$

Expand out

$$amplitude\left(|\phi\rangle \rightarrow |U\rangle\right) = \langle U| \times \left(a\,|U\rangle + b\,|D\rangle\right) = a\,\langle U| \times |U\rangle + b\,\langle U| \times |D\rangle$$

But we already know the answer to this question……..

Definition of state expansion

—> amplitude for ⏐ϕ⟩ changing into ⏐U⟩ is a . $|\phi\rangle = a\,|U\rangle + b\,|D\rangle$ Expansion 1

If rule consistent with already known answer, then must have

$$\langle U| \times |U\rangle = 1 \quad , \quad \langle U| \times |D\rangle = 0$$

This is totally consistent.

According to Rule 4,

$\langle U|\times|U\rangle$ = amplitude for $|U\rangle$ to change into $|U\rangle$,

which for normalized states = 1.

However, $|U\rangle$ cannot change into $|D\rangle$ (not directly),

so $\langle U|\times|D\rangle = 0$.

Two statements <=> orthonormality of (UP,DOWN) basis states (earlier)
(remember Hardness/color).

Hence

$$amplitude\left(|\phi\rangle \rightarrow |U\rangle\right) = a\,\langle U| \times |U\rangle + b\,\langle U| \times |D\rangle = a \times 1 + b \times 0 = a$$

Simplification: Bra acting on Ket, leave multiplication sign out

—> write $\langle\psi|\phi\rangle$ not $\langle\psi|\times|\phi\rangle$ <—> **Dirac Braket**.

Orthogonal Bases

$$\langle \phi \,|\, \phi \rangle = \{a_1^* \langle 1| + a_s^* \langle 2| + ... + a_n^* \langle n|\}\{a_1 |1\rangle + a_2 |2\rangle + ... + a_n |n\rangle$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{n} a_i^* a_j \langle i \,|\, j \rangle$$

all possible pairs!

States ⟩1⟨, ⟩2⟨, ⟩3⟨, etc = basis (abbreviation {⟩n⟨}).

If choose orthonormal basis = basis states, two things follow.

1. Terms ⟨n ⟩ m⟨, n not equal m vanish since ⟨n ⟩ m⟨ = 0

2. Terms ⟨1⟩1⟨ or ⟨n⟩m⟨, n = m, are ⟨1⟩1⟨=⟨n⟩n⟨=⟨m⟩m⟨=1, etc.

Remember $\langle n \,|\, m \rangle = \delta_{nm}$

Rules directly reflect experimental facts.

If states ⟩1⟨, ⟩2⟨, ⟩3⟨, etc, represent different measurement results

and experiment has separated out distinct paths

or quantitative values of a physical variable,

then one state cannot overlap with another

(they have nothing in common - they are orthogonal).

Hence amplitude for transition from ⟩m⟨ to ⟩n⟨ (m ≠ n) is zero for basis states.

—> physical basis for orthonormality of basis states corresponding to measurements.

Physicists describe any two states where ⟨n ı m⟩ = 0 as **orthogonal** states.

In good basis set,

all states are orthogonal to one another

and collection is called an **orthogonal basis**.

Assuming basis {ın⟩} is orthogonal set, calculation of ⟨ϕıϕ⟩ reduces nicely to

$$\langle \phi \,|\, \phi \rangle = \{a_1^* \langle 1| + a_s^* \langle 2| + ... + a_n^* \langle n|\}\{a_1 |1\rangle + a_2 |2\rangle + ... + a_n |n\rangle$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{n} a_i^* a_j \langle i \,|\, j \rangle \quad = 1$$

Now illustrate algebraically better way to do calculation. Use this new way later. Let

Details for the mathematically inclined ……

$$|\phi\rangle = a_1 |1\rangle + a_2 |2\rangle + \cdots + a_K |K\rangle = \sum_{i=1}^{K} a_i |i\rangle$$

K = size (dimension) of basis

$$\langle \phi| = a_1^* \langle 1| + a_2^* \langle 2| + \cdots + a_K^* \langle K| = \sum_{j=1}^{K} a_j^* \langle j|$$

$$\langle \phi \,|\, \phi \rangle = \left(\sum_{j=1}^{K} a_j^* \langle j| \right) \left(\sum_{i=1}^{K} a_i |i\rangle \right) = \sum_{j=1}^{K} \sum_{i=1}^{K} a_i a_j^* \langle j \,|\, i \rangle$$

Basis is orthonormal —> $\langle j \mid i \rangle = \delta_{ij}$ (Kronecker delta) and therefore

$$\langle \phi \mid \phi \rangle = \sum_{j=1}^{K} \sum_{i=1}^{K} a_i a_j^* \delta_{ij} = \sum_{i=1}^{K} a_i a_i^* = \sum_{i=1}^{K} |a_i|^2$$

or

$$\langle \phi \mid \phi \rangle = a_1^* a_1 + a_2^* a_2 + a_3^* a_3 + ... + a_K^* a_K = 1$$

Dirac language really is the Language(elegant) of QM .....

As starting point write |ϕ⟩ = a |U ⟩ + b |D⟩, which now becomes

$$|\phi\rangle = \langle U \mid \phi \rangle |U\rangle + \langle D \mid \phi \rangle |D\rangle$$

Since

$$\langle U \mid \phi \rangle = a \langle U \mid U \rangle + b \langle U \mid D \rangle = a \times 1 + b \times 0 = a$$

$$\langle D \mid \phi \rangle = a \langle D \mid U \rangle + b \langle D \mid D \rangle = a \times 0 + b \times 1 = b$$

Then write |U⟩ and |D⟩ in |L′⟩ and |R′⟩

$$|U\rangle = m |L'\rangle + n |R'\rangle = \langle L' \mid U \rangle |L'\rangle + \langle R' \mid U \rangle |R'\rangle$$

$$|D\rangle = p |L'\rangle + q |R'\rangle = \langle L' \mid D \rangle |L'\rangle + \langle R' \mid D \rangle |R'\rangle$$

and substitute into |ϕ⟩ expansion.

Earlier -> expression for amplitude governing transition $|\phi\rangle \rightarrow |L'\rangle$,

or (am + bp)=(ma + pb).

In Dirac language this is

$$|\phi\rangle = \langle U \mid \phi\rangle |U\rangle + \langle D \mid \phi\rangle |D\rangle$$

$$|\phi\rangle = \langle U \mid \phi\rangle \left(\langle L' \mid U\rangle |L'\rangle + \langle R' \mid U\rangle |R'\rangle\right) + \langle D \mid \phi\rangle \left(\langle L' \mid D\rangle |L'\rangle + \langle R' \mid D\rangle |R'\rangle\right)$$

$$|\phi\rangle = \left[\langle L' \mid U\rangle \langle U \mid \phi\rangle + \langle L' \mid D\rangle \langle D \mid \phi\rangle\right] |L'\rangle + \left[\langle R' \mid U\rangle \langle U \mid \phi\rangle + \langle R' \mid D\rangle \langle D \mid \phi\rangle\right] |R'\rangle$$

$$|\phi\rangle = [ma + pb]\,|L'\rangle + [na + qb]\,|R'\rangle$$

so that

$$amplitude\left(|\phi\rangle \underset{cannot\ tell\ which\ path}{\rightarrow} |L'\rangle\right) = \langle L' \mid U\rangle \langle U \mid \phi\rangle + \langle L' \mid D\rangle \langle D \mid \phi\rangle$$

$$= ma + pb$$

Beautifully illustrates how amplitudes combine together. Look closely.

1st term = amplitude takes $|\phi\rangle \rightarrow |L'\rangle$ via intermediate state $|U\rangle$.

2nd term = amplitude takes $|\phi\rangle \rightarrow |L'\rangle$ via intermediate state $|D\rangle$.

Expression = example of another important rule.

RULE 5: Any amplitude governing transition from initial state to final state via an intermediate state can be written

$$\langle final\ state \,|\, initial\ state\rangle = \langle final\ state \,|\, intermediate\ state\rangle \langle intermediate\ state \,|\, initial\ state\rangle$$

If intermediate states (labelled by i) are indistinguishable,

have to add all terms up to get overall amplitude

$$\langle final\ state \,|\, initial\ state \rangle = \sum_i \left[\langle final\ state \,|\, i \rangle \, \langle i \,|\, initial\ state \rangle \right]$$

Look carefully: I really have just inserted an IDENTITY operator

$$\sum_i |i\rangle \, \langle i| = \hat{I}$$

Rule 5 = extension of Rule 3(more formal way).

Going the Other Way

Dirac language —> amplitude for transition between initial state and final state is

$$\langle final\ state \,|\, initial\ state \rangle$$

Of course, always possible that process

goes in opposite direction from final to initial state.

Don't get confused about words initial and final; call them i and j.

RULE 6 (very important):

Amplitude from |i⟩ to |j⟩ is complex conjugate of amplitude from |j⟩ to |i⟩.

$$\langle i \mid j \rangle = \langle j \mid i \rangle^*$$

$$\langle i \mid j \rangle = \left(a^* \langle U| + b^* \langle D| \right) \left(c\,|U\rangle + d\,|D\rangle \right) = a^* c + b^* d$$

$$\langle j \mid i \rangle^* = \left[\left(c^* \langle U| + d^* \langle D| \right) \left(a\,|U\rangle + b\,|D\rangle \right) \right]^* = \left[c^* a + d^* b \right]^* = a^* c + b^* d = \langle i \mid j \rangle$$

QM Measurement (1st of many passes - we will get it correct by end of class!)

Now for most profound puzzle in QM

the process of measurement takes quantum state,

expressed as collection of possibilities,

and makes one of the possibilities ACTUAL.

Without measurement,

could not relate theory to reality.

Despite crucial nature of measurement,

still exist disputes about how it takes place in microworld.

Embracing Change

Change is built into structure of quantum state.

We have seen how state can be

expanded over set of basis states

representing different possible **outcomes** of experiment.

Once experiment performed,

original state has **collapsed** into one of basis states.

State collapse = peculiar process(more later), but is not only way quantum states change.

Most of time

for quantum systems that exist (not in experiments),

things can and do change.

Theory must also describe

normal processes where things

interact, change and develop.

Let us now look at ordinary time development

and, then come back to more mysterious, changes due to measurement.

Types of States

Quantum states divide into two groups.

State $|U\rangle$ —> particle with definite value of property —> eigenstates.

State $|\phi\rangle$ = combination of basis states

does not have definite value for measurement

—> mixed states or superpositions (exist as pure and non-pure types).

Eigenstates —> systems with definite value(physical property).

Know object in eigenstate,

can predict with absolute certainty a measurement result.

Remember:

sent electrons in |U⟩ into (UP,DOWN) S-G magnet

—> electrons emerged top path.

Add 2nd (UP,DOWN) measurement

—> electrons emerged top path.

Another property of eigenstate:

make measurement of physical property of eigenstate,

then another measurement —> no state change

—> Repeated measurement postulate.

If we choose to measure property different from that determined by eigenstate

—> cannot predict with certainty what will happen,

i.e., send |U⟩ electron into (LEFT,RIGHT) magnet,

—> do not know on which path it will emerge.

Eigenstates can have more than one physical property at same time

i.e., magenta electron with definite position.

Can measure either property,

predict with certainty what will happen,

and not change state during measurement

—> has definite values of both physical quantities.

Mixed States

$$|R\rangle = \frac{1}{\sqrt{2}}\left(|U\rangle + |D\rangle\right)$$

—> pure mixed state —> superposition !

Choose to measure electron in this state

using (UP,DOWN) S-G magnet,

get either |U⟩ or |D⟩ with 50:50 probability.

After long run of experiments,

expect to observe |U⟩ and |D⟩ with equal frequency.

Mixed states —> trouble when we want understand what QM says about the world.

Mixture is often not allowed in classical situation

i.e., Mach-Zehnder device

—> ended up with mixed state

that combined **two different paths** through device.

Classical particle follows one path

and cannot seem to be in two places at same time.

Mixed state is unavoidable in quantum analysis however.

Otherwise, we couldn’t explain experimental fact

that no photons ended up in one of detectors (for case of equal path lengths).

Quantum mixed states = central puzzle of quantum theory.

They are not some sort of average.

They do not describe an existence i.e., a blend of separate states(classical mixed state).

When we observe a quantum mixed state

—> mixture collapses into a component state.

Mixture

—> relative probability of finding component states

Mixed states —> set of **tendencies**(or **potentialities**) for something to happen.

State loaded with possibilities and measurement

—> one possibility = actuality.

Physicists disagree on the extent to which

they believe propensity or latency in state is related to physical nature of object.

Distinction between mixed state and eigenstate is not an absolute divide.

$|U\rangle$ and $|D\rangle$ eigenstates for (UP,DOWN) measurements,

but mixed states for (LEFT,RIGHT) measurements.

Similarly, $|L\rangle$ and $|R\rangle$ eigenstates for (LEFT,RIGHT) measurements,

but mixed states for (UP,DOWN) measurements.

Expectation Values

Suppose we have a large number electrons **all** in same state $|U\rangle$,

and send some at (UP,DOWN) S-G device.

Know(absolute certainty) results always are UP.

i.e., $|U\rangle$ = eigenstate of (UP,DOWN).

Send remainder of electrons at (LEFT,RIGHT) device

—> ability to predict is more limited.

Given specific electron

—> half the time emerges from LEFT channel

and half the time emerges from RIGHT channel.

No way of telling what happens to each electron

—> how **randomness** finds expression in quantum world.

We can, however, say something about the average.

Let RIGHT = +1 and LEFT = –1,

then for any set of measurements

average value for electrons is close to zero

(except for randomness and/or experimental precision).

Now consider set of electrons in state |ϕ⟩ where

$$|\phi\rangle = a\,|L\rangle + b\,|R\rangle \quad with \quad |a|^2 + |b|^2 = 1 \quad but \quad a \neq b$$

Electrons subjected to (LEFT,RIGHT) measurement,

find fraction = $|a|^2$ out LEFT

and fraction = $|b|^2$ out RIGHT.

No knowledge of each individual electron,

average value —> interesting result.

If $|a|^2$ —> result +1 and $|b|^2$ —> result –1, then average is

$$average\ value = |a|^2 \times (1) + |b|^2 \times (-1) = |a|^2 - |b|^2$$

Quantum theory:

average(set of measurements) on collection of identically prepared systems (same state)

= expectation value as we showed earlier.

Expectation value applies to a set of measurements.

For set of electrons, each electron

—> either (+1) or (–1)

—> not necessarily equal to expectation value,

which belongs to set as whole.

Some physicists say

that the connection between expectation values

and the results of set of measurements

—> view that states represent collections of systems

—-> electrons in state $|\phi\rangle$

—-> not saying each individual electron in state $|\phi\rangle$

but whole set)ensemble) has a state

—> puts to rest the view that state applies to an individual system!

Ensemble Interpretation

Some sense in that view.

Since can never tell from single measurement which state that system is in.

Sending electron at (UP,DOWN) magnet and comes out UP

—> cannot distinguish $|U\rangle$ as initial state

—> UP always

and $|R\rangle$ as initial state

—> made actual as $|U\rangle$ by measurement.

No help to repeat experiment on same electron;

once emerged from UP channel = state $|U\rangle$ (no matter how it started).

Must have a set and measure each separately.

Finding one comes out DOWN channel

—> can tell difference between |U⟩/|D⟩.

Would not allow us to tell difference between |R⟩/|L⟩, however.

Have to look at (LEFT,RIGHT) property for that.

So, clearly there is something to idea that states refer only to collections of systems.

Such view

—> instrumentalist interpretation of quantum theory.

Realist, however, wants to know what happening to each individual electron.

Now consider Operators

Need to be able to tell

if state is mixed state or eigenstate with respect to a measurement.

Mathematical machinery of quantum theory must allow for that,

otherwise can only construct state after the measurement.

An **operator** takes mathematical expression and transforms it into something else.

In quantum theory operators take one state into another

where process is governed by strict rules for each operator.

Many different operators in quantum theory

—> different jobs.

Most important —> operators representing process of measuring a physical property.

Example = (UP,DOWN) S-G operator, $\hat{S}_z$, (UP-DOWN direction is z–axis).

Role of operator = pull out of state information

about how state will react to (UP,DOWN) measurement.

Rules that govern how it works are simple.

|U⟩ and |D⟩ eigenstates of (UP,DOWN) measurements

—> (UP,DOWN) measurement

or corresponding operator does not change these states.

Thus,

$$\hat{S}_z \left|U\right\rangle = +1 \left|U\right\rangle \quad and \quad \hat{S}_z \left|D\right\rangle = -1 \left|D\right\rangle$$

Operator takes eigenstate and multiplies by number

equal to value found(on some meter) by measurement of particle in that state.

Action of operator = pull value from state.

Remember assigned values (+1) to UP and (–1) to DOWN

—> Definition of meaning of operator in this context.

Value multiplying the state = value experiment reveals with certainty

= eigenvalue of eigenstate.

Remember complete set of eigenstates for operator for physical variable

always = basis set. (Hermitian operators).

If state concerned not eigenstate of vertical(z-axis) S-G measurements,

then $\hat{S}_z$ makes a mess., i.e.,

$$\hat{S}_z |R\rangle = \hat{S}_z \left(\frac{1}{\sqrt{2}} |U\rangle + \frac{1}{\sqrt{2}} |D\rangle \right) = \left(\frac{1}{\sqrt{2}} \hat{S}_z |U\rangle + \frac{1}{\sqrt{2}} \hat{S}_z |D\rangle \right) \quad = \left(\frac{1}{\sqrt{2}} |U\rangle - \frac{1}{\sqrt{2}} |D\rangle \right) = |L\rangle$$

Action of operator on states (not eigenstates)

—> important role.

Look at this mathematical construction

$$\begin{aligned}
\langle R| \hat{S}_z |R\rangle &= \left(\frac{1}{\sqrt{2}} \langle U| + \frac{1}{\sqrt{2}} \langle D| \right) \hat{S}_z \left(\frac{1}{\sqrt{2}} |U\rangle + \frac{1}{\sqrt{2}} |D\rangle \right) \\
&= \left(\frac{1}{\sqrt{2}} \langle U| + \frac{1}{\sqrt{2}} \langle D| \right) \left(\frac{1}{\sqrt{2}} \hat{S}_z |U\rangle + \frac{1}{\sqrt{2}} \hat{S}_z |D\rangle \right) \\
&= \left(\frac{1}{\sqrt{2}} \langle U| + \frac{1}{\sqrt{2}} \langle D| \right) \left(\frac{1}{\sqrt{2}} |U\rangle - \frac{1}{\sqrt{2}} |D\rangle \right) \\
&= \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} \langle U \mid U\rangle - \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} \langle D \mid D\rangle \; = \frac{1}{2} - \frac{1}{2} = 0
\end{aligned}$$

Used fact ⟨U ı D⟩ = ⟨D ı U⟩ = 0 (property of basis states).

average value of UP-DOWN operator in the IR> state,i.e., it is equal parts UP and DOWN!

Not an especially useful calculation?

Don't jump to conclusions.

Do again using Iϕ⟩ instead of IR⟩.

Then

$$\langle\phi|\,\hat{S}_z\,|\phi\rangle = \left(a^*\langle U| + b^*\langle D|\right)\hat{S}_z\left(a|U\rangle + b|D\rangle\right)$$
$$= \left(a^*\langle U| + b^*\langle D|\right)\left(a\hat{S}_z|U\rangle + b\hat{S}_z|D\rangle\right)$$
$$= \left(a^*\langle U| + b^*\langle D|\right)\left(a|U\rangle - b|D\rangle\right)$$
$$= a^*a\langle U\mid U\rangle - b^*b\langle D\mid D\rangle = |a|^2 - |b|^2$$

average value of UP-DOWN operator in the |φ⟩ state, i.e., it is probability |a|² for +1 and probability |b|² for -1.

Mathematical constructions ⟨φ| $\hat{S}_z$ |φ⟩ or ⟨R| $\hat{S}_z$ |R⟩

—> expectation (average) values of series of measurements

made on state such as |φ⟩ or |R⟩ (happens to be 0 for |R⟩).

$\hat{S}_z$ not only example of operator

representing measurement of physical quantity.

Operators for other S-G magnet directions,

and position, momentum, energy, and all things that one can measure.

Connection of physical quantity <—> operator in quantum theory

Very different from classical physics.

Classical physics: state = collection of quantities describing object at instant.

Quantities given numerical value.

Classical laws —> rules that connect various quantities together

—> can predict future values.

Quantum mechanics: state = collection of amplitudes for object

to have values of physical quantities.

Physical quantity = operator —> expectation value.

It is average value obtained from set of measurements on identical systems;

none of systems necessarily have this expectation value.

Operators —> nothing by themselves;

need to act on states for any information extraction.

RULE 7: Every physical variable has an associated operator $\hat{O}$.

Operators have eigenstates $|\psi\rangle$ defined by

$$\hat{O}\,|\psi\rangle = a\,|\psi\rangle$$

where a = value of physical variable get if measured state $|\psi\rangle$.

Complete set eigenstates $\{|\psi\rangle\}$ = basis.

Operator associated with physical variable

—> expectation value of series of measurements

made on collection of systems in same state $|\phi\rangle$.

Given by

$$\left\langle \hat{O} \right\rangle = \langle\phi|\,\hat{O}\,|\phi\rangle$$

How can we represent operators?

Think of operator as some kind of box,

where we put vector in

and get another(different or same) vector out. **or like color/hardness boxes**

Similar to definition of function for space of numbers.

Some properties of operators in Quantum Mechanics:

$$\hat{Q}\left(|A\rangle + |B\rangle\right) = \hat{Q}\,|A\rangle + \hat{Q}\,|B\rangle \;\; (\text{linearity})$$
$$\hat{Q}\left(c\,|A\rangle\right) = c\hat{Q}\,|A\rangle \quad , \quad c = complex\ number$$
$$\langle C|\left(\hat{Q}\,|B\rangle\right) = \langle C\,|\,B'\rangle = \;\text{number} \equiv \langle C|\,\hat{Q}\,|B\rangle \;\; (\text{matrix element})$$
$$(\hat{Q}_1 + \hat{Q}_2)\,|A\rangle = \hat{Q}_1\,|A\rangle + \hat{Q}_2\,|A\rangle \;\; (\text{linearity})$$
$$(\hat{Q}_1\hat{Q}_2)\,|A\rangle = \hat{Q}_1(\hat{Q}_2\,|A\rangle) \;\; (\text{order matters})$$

Properties —> operators = LINEAR operators.

QM understood using only linear operators

—> truly amazing - simplest kind of operator that mathematicians can think of.

Nature is very efficient!!

All observables or quantities we can measure are represented by operators in QM.

Repeat the discussion from earlier.

Additional pass(many discussions in between), should greatly enhance understanding and enable us to learn a few new things.

Suppose make measurements on state |ψ⟩

of observable given by the operator $\hat{B}$

with eigenvalues/eigenvectors given by $\hat{B}\,|b_j\rangle = b_j\,|b_j\rangle \quad j = 1, 2, 3, 4, ...$

Then we can write

$$|\psi\rangle = \sum_k a_k\,|b_k\rangle$$

OK since eigenvectors of observable operator = complete set = basis

—> $\langle b_j\,|\,\psi\rangle = \langle b_j|\sum_k a_k\,|b_k\rangle = \sum_k a_k\,\langle b_j\,|\,b_k\rangle = \sum_k a_k\delta_{jk} = a_j$

—> amplitude to measure b_j

Since measurement results(= eigenvalues) are values b_k, and suppose that

b_k occurs n_k times where k=1,2,3,4,…….and where

$$\sum_k n_k = N = \text{ total number of measurements}$$

Then from the standard definition of the average value, have

$$\left\langle \hat{B} \right\rangle = \text{ average or expectation value of } \hat{B} \quad = \frac{1}{N}\sum_k n_k b_k = \sum_k \frac{n_k}{N} b_k = \sum_k b_k prob(b_k)$$

From postulates (rules) we have $prob(b_k) = |\langle b_k\,|\psi\rangle|^2$

—> $\left\langle \hat{B} \right\rangle = \sum_k b_k prob(b_k) = \sum_k b_k\,|\langle b_k\,|\,\psi\rangle|^2$

Now $\langle b_k | \psi \rangle^* = \langle \psi | b_k \rangle$.

Therefore, $|\langle b_k | \psi \rangle|^2 = \langle \psi | b_k \rangle \langle b_k | \psi \rangle$ and

$$\left\langle \hat{B} \right\rangle = \sum_k b_k \left| \langle b_k \mid \psi \rangle \right|^2 = \sum_k b_k \langle \psi \mid b_k \rangle \langle b_k \mid \psi \rangle = \langle \psi | \left[\sum_k b_k \left| b_k \right\rangle \langle b_k \right| \right] | \psi \rangle$$

Now earlier definition of expectation value which was

$$\left\langle \hat{B} \right\rangle = \langle \psi | \hat{B} | \psi \rangle$$

IMPORTANT

—> can represent operator $\hat{B}$ by expression

$$\hat{B} = \sum_k b_k \left| b_k \right\rangle \langle b_k |$$

(a result we have already used several times)

IMPORTANT

A very important way represent operator,

i.e., any operator is represented in terms of its eigenvalues/eigenvectors

—> called the **spectral decomposition** of operator.

Projection Operators

Operators of the form $\hat{P}_\beta = |\beta\rangle \langle \beta|$ = projection operators.

Can write any operator $\hat{B}$ as $\hat{B} = \sum_k b_k P_{b_k}$ where $P_{b_k} = |b_k\rangle \langle b_k|$

projection operator onto state $|b_k\rangle$.

Very important property of projection operators: $\hat{I} = \sum_k |b_k\rangle \langle b_k|$ **IMPORTANT**

$$\hat{I}\,|\beta\rangle = \sum_k |b_k\rangle \langle b_k \mid \beta\rangle = \sum_k \langle b_k \mid \beta\rangle\, |b_k\rangle = |\beta\rangle$$

Clearly it is the IDENTITY operator or an operator that leaves all kets unchanged.

Some Operators

$$\hat{O}_{color}\,|m\rangle = (+1)\,|m\rangle \quad , \quad \hat{O}_{color}\,|g\rangle = (-1)\,|g\rangle$$

—> $\hat{O}_{color} = |m\rangle \langle m| - |g\rangle \langle g|$ Does it work? We have

$$\hat{O}_{color}\,|m\rangle = (|m\rangle \langle m| - |g\rangle \langle g|)\,|m\rangle = |m\rangle$$

$$\hat{O}_{color}\,|g\rangle = (|m\rangle \langle m| - |g\rangle \langle g|)\,|g\rangle = -\,|g\rangle$$

as expected!

Note |h⟩ state has equal amounts of magenta/green.

Therefore expectation value of color operator in hard state should = zero.

This is confirmed below.

$$\langle h|\,\hat{O}_{color}\,|h\rangle = \frac{1}{\sqrt{2}}\left(\langle m| + \langle g|\right)\hat{O}_{color}\frac{1}{\sqrt{2}}\left(|m\rangle + |g\rangle\right)$$

$$= \frac{1}{2}\left[\langle m|\,\hat{O}_{color}\,|m\rangle + \langle m|\,\hat{O}_{color}\,|g\rangle + \langle g|\,\hat{O}_{color}\,|m\rangle + \langle g|\,\hat{O}_{color}\,|g\rangle\right]$$

$$= \frac{1}{2}\left[\langle m \mid m\rangle - \langle m \mid g\rangle + \langle g \mid m\rangle - \langle g \mid g\rangle\right] \quad = \frac{1}{2}\left[1 - 0 + 0 - 1\right] = 0$$

Dirac language is very powerful!

Similarly, can represent an (UP,DOWN) S-G magnet or operator $\hat{S}_z$ by

$$\hat{S}_z = |U\rangle\langle U| - |D\rangle\langle D|$$

IMPORTANT

In words, the sum of projection operators for eigenvectors times eigenvalues

Now to Figure Out How States Evolve

Operator that tells us how a state evolves in time is not directly a physical variable.

= time evolution operator $\hat{U}(t)$

—-> takes a state and moves it forward in time:

$\hat{U}(t)\,|\Psi(T)\rangle = |\Psi(T+t)\rangle$ where $|\Psi(T)\rangle$ = state of system at time T.

For $\hat{U}(t)$ to do this accurately,

it must contain all details of the behavior of system

i.e., what system is doing and how it interacts with its environment.

Example:

Consider the state |U⟩ moving through space toward S-G magnet.

Electron interacts with surroundings —> disturbs its S-G orientation.

State will evolve from |U⟩ to different state, |ϕ⟩,

in a smooth and deterministic manner over time.

Energy of electron will determine the type of evolution,

i.e., time evolution operator <—> related to energy operator for electron.

Can always describe any S-G state using {|U⟩,|D⟩} basis. $|\Psi\rangle = a\,|U\rangle + b\,|D\rangle$

So, what really happens when $\hat{U}(t)$ operator acts is

$$\hat{U}(t)\,|\Psi(T)\rangle = \hat{U}(t)\,[a(T)\,|U\rangle + b(T)\,|D\rangle] \;= [a(T+t)\,|U\rangle + b(T+t)\,|D\rangle]$$

$\hat{U}(t)$ changes amplitudes values at time T

to new ones at time T + t. (it does not change the basis vectors!!)

Now take the extra time interval t and divide into many smaller intervals, size δt.

If 100 small pieces, then t = 100 × δt.

Time evolution operator works same way for very small time intervals as for longer ones.

Applying $\hat{U}(t)$ once takes |Ψ(T)⟩ to |Ψ(T + t)⟩.

Must get same answer applying $U(\delta t)$ 100 times.

Apply $U(\delta t)$ and we get

$$\hat{U}(\delta t)\,|\Psi(T)\rangle = \hat{U}(\delta t)\,[a(T)\,|U\rangle + b(T)\,|D\rangle] \;= [a(T+\delta t)\,|U\rangle + b(T+\delta t)\,|D\rangle]$$

For small δt , a(T + δt) not very different from a(T) (same for amplitude b).

Applying $U(\delta t)$ again gives us

$$\hat{U}(\delta t)\,|\Psi(T+\delta t)\rangle = \hat{U}(\delta t)\,[a(T+\delta t)\,|U\rangle + b(T+\delta t)\,|D\rangle] \;= [a(T+2\delta t)\,|U\rangle + b(T+2\delta t)\,|D\rangle]$$

and so forth.

State's evolution from |Ψ(T)⟩ to |Ψ(T + t)⟩

takes place via continuously smooth change.

(limit as δt → 0 -> DiffEQ —> Schrodinger Equation governs evolution)

Evolution completely determined by physics of system, as expressed by $\hat{U}(t)$.

No randomness involved here.

Inclusion of Postulate 4 (collapse)

—> smooth evolution not only way state can evolve.

States can also evolve in sharp(discontinuous)

and unpredictable manner

when measurement takes place

= definition of measurement.

later we will see if this statement is really necessary or whether this view of what is happening is old and outdated!!

For the moment, however, let us continue with this old line of reasoning……

When electron in state $|\Psi\rangle$ reaches (UP,DOWN) S-G magnet,

state will “change”

into $|U\rangle$ with probability $|a|^2$

or into $|D\rangle$ with probability $|b|^2$.

depending on what we measure

After measurement, the state

has evolved into $|U\rangle$ (i.e., a→1 , b→0) or $|D\rangle$ (i.e., a→0 , b→1).

During a measurement, the amplitudes a and b have not changed continuously

(or so it seems) as they did with the $\hat{U}(t)$ operator.

Dramatic and unpredictable change in quantum state as a result of a measurement

—> **collapse or reduction of state**.

State collapse = radically different process from $\hat{U}(t)$ evolution,

can’t be broken down into smooth progression of small steps.

Mathematically impossible for equations of QM as we currently understand them,

to describe state collapse (a non-linear process cannot be described by linear operators).

Remains as add-on assumption, it cannot be predicted from within theory it seems.

A significant point is being made here **(much more on this later)**.

QM correctly describes the evolution of states in time; but doesn’t seem to describe the real world (measurement) unless(at this moment) we add the state collapse idea.

World is **not** = set of evolving possibilities.

Possibilities encoded in quantum state must be linked to actual events in world.

Collapse of state <—> measurement

—> some physicists seek an instrumentalist view of quantum states.

Argue that discontinuous change in state

shows state **only** represents information about system.

Before a dice thrown, information —> 1/6 probability of each face.

After throw, information state collapses, know which face appeared.

Dice has not changed in any physical manner.

Correspondingly when quantum state collapses —> does not necessarily signal

any physical difference in system being described,

just a change of **knowledge** about it.

To counter this, realist —> quantum state cannot just be knowledge of system,

since how can “knowledge” have direct effect on system’s behavior?

In so-called delayed choice experiments that we will discuss later

we will see that our ability/lack of ability to infer information about state of photon

has a direct result on outcome of experiment.

Therefore **knowledge(information)** must be reflection of something **real** to do with the system.

Quantum states —> weird and spooky

compared to classical reality we are used to,

but we can still believe science is revealing real truths

about world even if we find truths surprising.

SUMMARY

$\hat{U}(t)$ operator evolves state forward in time in smooth and predictable manner

$$\hat{U}(t)\,|\Psi(T)\rangle = |\Psi(T+t)\rangle$$

Exact form of $\hat{U}(t)$ operator depends on physics of system.

State collapse is sharp and unpredictable change in state

as result of measurement (state not eigenstate of physical quantity measured).

Why is State Reduction Necessary?

Reasonable to ask how this whole mess came about.

Why are we driven to construct theory that requires this odd concept of state collapse?

Argument is:

(1) Events in microworld, (reflection or transmission of photons from half-silvered mirror)
seem to be random.
—> Must represent state of microscopic objects using probabilities.
Assign number to each possibility,
and number tells us probabilities of events in future.

(2) Mach-Zehnder experiment —> numbers (amplitudes)
= complex numbers -> not probabilities.
Link between amplitudes and probabilities guessed
(probability = |amplitude|^2) —> agreement with experiment.

(3) Led us to represent quantum state of system as collection of amplitudes.

(4) In certain cases, appropriate quantum state
= mixture of states that have to be separate classically,
i.e., many experiments only understood
if the intermediate state of the photon APPEARS
to follow two **different** classical paths at same time.
Without this, interference effects would not take place
and actual experimental results could not be explained.

(5) Although quantum theory allows these mixed states to occur,
observers seem to be protected from them
—> they are never directly observed as outcomes of experiments.
i.e., photon is never observed to be on both paths at same time.

(6) Quantum state of system changes.
Prepare beam of electrons for Stern-Gerlach (S-G) experiment,
find 50% exit device along each channel,
indicating electrons were mixed states to start with.
However, if pass all electrons from one channel into another identical S-G device,
then all emerge from same channel.
1st experiment has changed quantum state from unobservable mixed state
into classical-like eigenstate that is definitely |U⟩ or |D⟩.

(7) Crux of problem.

Certain experiments only explained if classically forbidden mixtures allowed.

However, live in classical world so quantum mixed states

must collapse into more classical-like states that we can directly observe.

Much more later when we solve all these measurement confusions.

Finally we now explain the real experiment we started our entire discussion with!

Explaining the Double-Slit Experiment with Amplitudes

The quantum mechanical description of double-slit experiment

must involve a set of amplitudes

governing the passage of an electron through slits to a far detector.

Let us define the following states

$|x\rangle$ is state corresponding to electron


arriving at position x of detector (see diagram),

$|A\rangle$ corresponds to passing through slit A,

$|B\rangle$ corresponds to passing through slit B,

and $|I\rangle$ corresponds to the initial state of electron

emerging from electron gun.



Then

$$\langle x \mid I \rangle$$

is the probability amplitude we want to calculate.

We assume the electron gets from source (initial state $|I\rangle$) to detector (state $|x\rangle$)
via two possible intermediate states (either passing through slit A or slit B).

$|A\rangle$ $|B\rangle$


Then we can write

Intermediate state A Intermediate state B

$$\langle x \mid I \rangle = \langle x \mid A \rangle \langle A \mid I \rangle + \langle x \mid B \rangle \langle B \mid I \rangle$$

amplitude to calculate inserted sum over intermediate states as described earlier

here **all** possibilities -> both slits



Experimentally, we see bright and dark interference bands.

Dark band separation given by $D\lambda/a$


where λ is wavelength.

Must add together two terms in overall amplitude,

two possibilities(traveling through one slit or other)

which CANNOT be distinguished

in the context of experiment as set up.



When combine amplitudes this way interference between terms results as we will see.

If block a slit, then one of terms will disappear —> no interference.

Now probability that electron arrives at x is complex square of total amplitude.

$$|\langle x \mid I\rangle|^2 = [\langle x \mid A\rangle \langle A \mid I\rangle + \langle x \mid B\rangle \langle B \mid I\rangle]^* [\langle x \mid A\rangle \langle A \mid I\rangle + \langle x \mid B\rangle \langle B \mid I\rangle]$$

Expanding out:

bump would see if only A open — bump would see is only B open

$$|\langle x \mid I\rangle|^2 = \langle x \mid A\rangle^* \langle A \mid I\rangle^* \langle x \mid A\rangle \langle A \mid I\rangle + \langle x \mid B\rangle^* \langle B \mid I\rangle^* \langle x \mid B\rangle \langle B \mid I\rangle$$

interference terms

$$+ \langle x \mid A\rangle^* \langle A \mid I\rangle^* \langle x \mid B\rangle \langle B \mid I\rangle + \langle x \mid B\rangle^* \langle B \mid I\rangle^* \langle x \mid A\rangle \langle A \mid I\rangle$$

1st two terms give that conventional (classical) sum of probabilities

that one expect(bumps) for an electron that goes through one slit or other.

Final two terms is where interesting stuff resides: **the quantum interference terms**.

Digression: Phase and Physics

Let us review/reconsider complex numbers in different form

—> more useful form for this discussion.

The Complex Plane

Complex numbers needed if mathematical objects in QM are to represent reality.

Rethink ideas already introduced,

explore **exponential** way of writing complex numbers

and introduce **geometrical picture** to help us understand.

Sometimes convenient to think of complex number

as denoting point on special type of graph.

Mathematicians sometimes picture real numbers

lying on line (real axis) stretching out to infinity (in two directions).


As work way along line (either direction)

at some point come across the number zero.

At zero point, any points on line to right

are positive real numbers

and those to left are negative real numbers



When complex numbers appeared,

mathematicians thought in terms of two lines at right angles to one another.

A purely real number lies on horizontal line,


and a purely imaginary number (e.g., 3i or 5i or –2.1678i)

lies somewhere on vertical line running through zero point(origin).

Any two lines at right angles map out or define

a 2-dimensional (2-D) region(plane)

—> in this case, the **complex plane**.



Any complex number, a + ib,

is identified with a point on the complex plane.

To get to point walk along real line distance a,

turn left by a right angle,

and walk parallel to imaginary line distance b

—> graphical representation of the point in the complex plane.

Horizontal arrow with x next to it

-> x coordinate axis.

Normal graph -> vertical arrow with y next to it

-> y coordinate axis along vertical.

Have labelled vertical line iy for illustration (not really allowed - no such distance).

Point of the graph is to represent quantities (numbers with units)

as lengths according to some scale.

Should just put y on vertical axis —> complex number is of form


$$z = (x \text{ coordinate}) + i(y \text{ coordinate})$$

Reason not done

—> I want to emphasize

that in figures **neither** x direction nor y direction is an actual physical direction in space.

Figures just represent a geometric way of **picturing** complex numbers.



On the plane -

always more than one way of getting to particular point.

Rather than walking specific distances along two lines,

can approach same point by more direct route.


Stand on zero point(origin) looking along real line.

Turn in counterclockwise (or anticlockwise) direction

through angle θ

and walk in that direction distance R as shown

—> end up at exactly same point a + ib



Can think of line of length R

as hand of clock

and θ the angle that hand sweeps out

(starts sweeping at 3 o'clock (+ x-axis)

and goes backward (anticlockwise)!).

If R and θ get to the same place as a and b

—> must exist some formulas that connects two sets of numbers.

They are $a = R\cos\theta \ , \ b = R\sin\theta \rightarrow \frac{b}{a} = \tan\theta \ , \ R^2 = a^2 + b^2$

Note that R is **always** positive.

One does not move negative distance away from the zero point.

—> Have two ways of writing same complex number as 2-tuples,

either as z = a + ib = (a,b)

or z = (R,θ) (derive formula later).

Magnitude and Phase

1st think about R => **magnitude** of complex number.

Complex conjugate of number z = a+ib is z^* = a–ib.

If multiply complex number by its conjugate —> real number.

This number is

$$zz^* = (a+ib)(a-ib) = a^2 + b^2 = R^2$$ using Pythagorous

—> mathematical way to convert complex number into real number.

Earlier, same procedure gave us probabilities from probability amplitudes.

Now switch things around.

Many complex numbers have the same R.

Draw circle around zero point of radius R;

All points on circumference of that circle —> complex numbers with same R value.




Figure illustrates idea.

Complex numbers z = a + ib and Z = A+iB have same R

—> equal magnitudes of complex numbers.

Both lead to same absolute value.

Infinite number of complex numbers —> same absolute value.

All complex numbers with same R differ only in value of θ

—> **argument** (mathematicians) of complex number;

Physics —> call θ **the phase** of complex number.

(Will see it corresponds directly to physical phase used earlier in interference experiments).

Will turn out that magnitude of complex numbers is related to observable probability

and phase is connected to quantum interference effects.

Multiplying Complex Numbers


If can write complex number z = a + ib in form (R,θ),

then how would we write iz?

iz would be i(a + ib) = ia – b or –b + ia

-> complex numbers z and iz have same magnitude, R.

However, have different phase, i.e., see figure.



Phases of z and iz differ by 90°, i.e., see figure

$$2\alpha + 2\beta = \pi \rightarrow \alpha + \beta = \pi/2 \rightarrow orthogonal$$

Thus, whenever multiply by i,

rotate phase anticlockwise by 90°,

but leave magnitude alone.

Push these ideas further.

i is number that lies one unit along imaginary axis.

Lots of complex numbers have the same magnitude as i (i.e., R = 1)

but different phases.

Number 1 (= 1 + 0i) for example.

What happens if multiply by other numbers instead?

Take number z = a + ib and multiply by another number w = p + iq

such that $p^2 + q^2 = 1$ —> R = 1 for w.

$$zw = (a+ib)(p+iq) = ap + iaq + ibp - bq = (ap - bq) + i(aq + bp)$$

What is magnitude of new number?

Square of magnitude = square of real part + square of imaginary part.

$$R^2 = (ap - bq)^2 + (aq + bp)^2 = a^2p^2 + b^2q^2 - 2apbq - a^2q^2 - b^2p^2 - 2aqbp$$

$$= a^2p^2 + b^2q^2 + a^2q^2 + b^2p^2 = a^2(p^2 + q^2) + b^2(p^2 + q^2) \ = a^2 + b^2$$ using $p^2 + q^2 = 1$.

Thus, multiplying any complex number z by another number complex number w,

when w has magnitude 1(R=1), **doesn't change** magnitude of z.

What about phase though?

Tougher to prove - can show, however, that phases **add**.

Easy proof later using exponential form of complex numbers

and harder proof now using trigonometry (see below).

In fact, the general rule is derived as follows:

MULTIPLYING COMPLEX NUMBERS

When two complex numbers z and w are multiplied together,

magnitudes multiply and **phases add**.

$$If\ z = (R_1, \theta_1)\ and\ w = (R_2, \theta_2),\ then\ zw = (R_1 R_2, \theta_1 + \theta_2)$$

Proof:(using trigonometry)

$$z = a + ib \quad , \quad w = p + iq \qquad\qquad \tan\theta_z = \frac{b}{a} \quad , \quad \tan\theta_w = \frac{q}{p}$$

$$zw = (ap - bq) + i(aq + bp)$$


$$\tan\theta_{zw} = \frac{aq + bp}{ap - bq} = \frac{\frac{q}{p} + \frac{b}{a}}{1 - \frac{q}{p}\frac{b}{a}} = \frac{\tan\theta_z + \tan\theta_w}{1 - \tan\theta_z \tan\theta_w} = \tan(\theta_z + \theta_w) \qquad \Rightarrow \theta_{zw} = \theta_z + \theta_w$$

More Phase Stuff

Figure shows how two complex numbers

z = a + ib and –z = –a – ib

are related on the complex plane.



z and –z have same R.

Phases different by 180°

—> Multiplying z by i and again by i, as $i^2 = -1$.

Each multiplication by i shifts phase by 90°, hence 180° overall.

And Now Pulling It All Together

Complex numbers can be represented in two different, but equivalent ways

-> two numbers a and b —> z = a+ib = (a,b)

or two numbers R and θ —> (R,θ) .

Another way of writing complex numbers —> combines elements of both forms.

Need to review some other mathematics first (stated these relations earlier).

Some functions in mathematics represented by power series

—> real definitions of functions,

i.e., how we calculate them. So as we mentioned earlier….

A power series representation of a function:

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots = \sum_{k=0}^{\infty} a_k x^k$$

Special cases

$$e^{\alpha x} = 1 + \alpha + \frac{1}{2}\alpha^2 + \frac{1}{6}\alpha^3 + ... = \sum_{n=0}^{\infty} \frac{\alpha^n}{n!} x^n$$

$$\sin \alpha x = \alpha x - \frac{1}{6}(\alpha x)^3 + \frac{1}{24}(\alpha x)^5 + ... = \sum_{n=0}^{\infty} (-1)^n \frac{\alpha^{2n+1}}{(2n+1)!} x^{2n+1}$$

$$\cos \alpha x = 1 - \frac{1}{2}(\alpha x)^2 + \frac{1}{16}(\alpha x)^4 + ... = \sum_{n=0}^{\infty} (-1)^n \frac{\alpha^{2n}}{(2n)!} x^{2n}$$

Expansions are still valid if α is a complex number

—> important mathematical result for QM

Now

$$e^{i\alpha x} = \sum_{n=0}^{\infty} \frac{i^n \alpha^n}{n!} x^n = 1 + i\alpha x - \frac{\alpha^2}{2!}x^2 - i\frac{\alpha^3}{3!}x^3 + \frac{\alpha^4}{4!}x^4 + i\frac{\alpha^5}{5!}x^5 - .....$$

rearrange terms

$$= \left(1 - \frac{\alpha^2}{2!}x^2 + \frac{\alpha^4}{4!}x^4 - .....\right) + \left(i\alpha x - i\frac{\alpha^3}{3!}x^3 + i\frac{\alpha^5}{5!}x^5 - .....\right)$$

$$= \cos\alpha x + i\sin\alpha x$$ —-> Euler relation (derive in 16th century)

Then

$$\sin\alpha x = \frac{e^{i\alpha x} - e^{-i\alpha x}}{2i} \quad , \quad \cos\alpha x = \frac{e^{i\alpha x} + e^{-i\alpha x}}{2}$$

Euler relation —> define i without using square root of negative number!!

$$e^{i\pi} = \cos\pi + i\sin\pi = -1 \qquad e^{i\pi/2} = \cos\pi/2 + i\sin\pi/2 = i$$

Since $e^{a+b} = e^a e^b$ $e^a = e^{a/2}e^{a/2} \rightarrow e^{a/2} = \sqrt{e^a}$ and $(e^a)^n = e^{na}$

$$\sqrt{e^{i\pi}} = e^{i\pi/2} = \cos\pi/2 + i\sin\pi/2 = i$$

Now from earlier we had $a = R\cos\theta \ , \ b = R\sin\theta \rightarrow \frac{b}{a} = \tan\theta \ , \ R^2 = a^2 + b^2$

or

$$z = a + ib = R\cos\theta + iR\sin\theta = R(\cos\theta + i\sin\theta) = Re^{i\theta}$$ new form of complex number

Write complex conjugate z^* using exponential form.


$$z* = a - ib = R\cos\theta - iR\sin\theta = R\left[\cos(-\theta) + i\sin(-\theta)\right] = Re^{-i\theta}$$

Since using cosθ/sinθ this way

—> not directly related to geometry

—> convenient to use different measure

for θ than angle(degrees)



Mathematicians/physicists call new measure the **radian**.

Normal way measure angle

—> take circle, divide circumference into 360 pieces .

Draw lines from ends of pieces to center

—> protractor. Angle between pair of lines = 1°.

No real reason for 360 pieces. A more interesting possibility is….

Circumference = 2π × radius

—> imagine dividing circumference into 2π pieces.

Cannot do(not exactly)

since π requires an infinite number of digits to define it.

OK since not interested in making a real protractor

but just finding a mathematically convenient measure of angle.

If divided circle into 2π pieces,

then angle between pairs of lines = 1 radian.

Whole circle, = 360° = 2π radians. 90° = 1/4 circle = 2π/4 = π/2 radians.

180° = π radians, etc.

Finally, a very useful property of the exponential form of complex numbers.

Earlier, multiplication rule

—> when multiply z_1 by z_2, magnitudes multiply and phases add.

Easier proof:

$$z_1 z_2 = \mathrm{R}_1 \ \exp(i\theta_1)\mathrm{R}_2 \ \exp(i\theta_2) = (R_1 R_2)\exp\left(i(\theta_1+\theta_2)\right)$$

proof now a one-liner

Now Returning to Phase and Physics

Now we can explain electron interference pattern revealed by detector,

given the amplitudes that govern two possible routes from slits to detector.

Existence of dark bands, parts of detector where no electrons arrive

= key point we need to factor into our thinking.

When add up amplitudes for getting dark point on detector,

total amplitude must be **zero**.


How can we be sure this happens?

If thinking of water waves interfering


—> saying path difference between two routes

equivalent to multiple of half of wavelength,

so that waves arrived **out of phase**.



out of phase.



in phase.

Perhaps something similar(mathematically) is happening with amplitudes.

To explain dark band at position x on detector, must have

$$\langle x \mid I \rangle = \langle x \mid A \rangle \langle A \mid I \rangle + \langle x \mid B \rangle \langle B \mid I \rangle = 0$$

Set ⟨A|I⟩ and ⟨B|I⟩ equal ⟨slits|I⟩

I.e., if source midway between slits, these amplitudes are the ~ same. Thus

$$\langle x \mid I \rangle = [\langle x \mid A \rangle + \langle x \mid B \rangle] \langle slits \mid I \rangle = 0$$

Two ways can be true.

⟨slits | I⟩ = 0 —> electrons not getting to slits. Silly = Non-physical.

Or ⟨x | A⟩ + ⟨x | B⟩ = 0 —> related to interference.

Both true —> set up poor experiment.

If ⟨x|A⟩+⟨x|B⟩ = 0,

then ⟨x|A⟩ = –⟨x|B⟩

—> two complex numbers with same magnitude,

but opposite phases (like z and –z).

Suppose change position of one slit without moving other

—> changes length of one path compared to other

—> convert a point with destructive interference

into one with constructive interference.

To do

—> moved slit so that path altered by distance equal to half wavelength.

Question: does this alter phase or magnitude, or both?

Move slit even further

—> path difference is 3λ/2

-> destructive interference

-> two amplitudes once again, have same magnitude but opposite phases.

Unlikely simply increasing path difference

—> magnitude changes.

However, can imagine alterations to path changing phase of amplitude

remember how phase was defined(as # wavelengths in path).

If amplitude of form $Re^{i\theta}$

and θ determined by path length,

then steadily increasing path would increase angle,

taking us round and round circle in complex plane.

Mathematicians don't limit angles to ≤ 360° (or 2π radians),

i.e., angle of 370° —> once round circle and 10° more.

Comparing two amplitudes of this type could quite easily

move us from situation where their phases are identical

to out of phase and back again as path length changed.

If right, then **wavelength** associated with electron

is somehow **coded** into **amplitude phase**.

$$\lambda = \frac{h}{p}$$

So finally : An Experiment with Phase

Write two amplitudes(now just complex numbers) in the form

$$\langle x \mid A \rangle = R_1 \exp\left(i\theta_1(x,t)\right) \quad , \quad \langle x \mid B \rangle = R_2 \exp\left(i\theta_2(x,t)\right)$$

—> phase θ depends on position and time;

put into probability calculation

—> after wild algebra

(either ignore it and jump to result

or if more mathematically inclined work through algebra).

$$
\begin{aligned}
&|\langle x \mid I\rangle|^2 = |\langle x \mid A\rangle + \langle x \mid B\rangle|^2 \, |\langle slits \mid I\rangle|^2 \\
&= |R_1 \exp(i\theta_1(x,t)) + R_2 \exp(i\theta_2(x,t))|^2 \, |\langle slits \mid I\rangle|^2 \\
&= [R_1 \exp(i\theta_1(x,t)) + R_2 \exp(i\theta_2(x,t))]^* \, [R_1 \exp(i\theta_1(x,t)) + R_2 \exp(i\theta_2(x,t))] \, |\langle slits \mid I\rangle|^2 \\
&= [R_1 \exp(-i\theta_1(x,t)) + R_2 \exp(-i\theta_2(x,t))] [R_1 \exp(i\theta_1(x,t)) + R_2 \exp(i\theta_2(x,t))] \, |\langle slits \mid I\rangle|^2 \\
&= [R_1 (cos(-\theta_1(x,t)) + i\sin(-\theta_1(x,t))) + R_2 (cos(-\theta_2(x,t)) + i\sin(-\theta_2(x,t)))] \\
&\quad \times [R_1 (cos(\theta_1(x,t)) + i\sin(\theta_1(x,t))) + R_2 (cos(\theta_2(x,t)) + i\sin(\theta_2(x,t)))] \, |\langle slits \mid I\rangle|^2 \\
&= [R_1 (cos(\theta_1(x,t)) - i\sin(\theta_1(x,t))) + R_2 (cos(\theta_2(x,t)) - i\sin(\theta_2(x,t)))] \\
&\quad \times [R_1 (cos(\theta_1(x,t)) + i\sin(\theta_1(x,t))) + R_2 (cos(\theta_2(x,t)) + i\sin(\theta_2(x,t)))] \, |\langle slits \mid I\rangle|^2 \\
&= \begin{bmatrix} R_1^2 \left(cos^2(\theta_1(x,t)) + \sin^2(\theta_1(x,t))\right) + R_2^2 \left(cos^2(\theta_2(x,t)) + \sin^2(\theta_2(x,t))\right) \\ +2R_1R_2 (cos(\theta_1(x,t)) cos(\theta_2(x,t)) + \sin(\theta_1(x,t)) \sin(\theta_2(x,t))) \\ +iR_1R_2 (cos(\theta_1(x,t)) \sin(\theta_2(x,t)) + \sin(\theta_1(x,t)) \cos(\theta_2(x,t))) \\ -iR_1R_2 (cos(\theta_1(x,t)) \sin(\theta_2(x,t)) + \sin(\theta_1(x,t)) \cos(\theta_2(x,t))) \end{bmatrix} |\langle slits \mid I\rangle|^2 \\
&= \left[R_1^2 + R_2^2 + 2R_1R_2 (cos(\theta_1(x,t)) cos(\theta_2(x,t)) + \sin(\theta_1(x,t)) \sin(\theta_2(x,t)))\right] |\langle slits \mid I\rangle|^2
\end{aligned}
$$

In last line:

1st two terms

-> uniform probability for electron to arrive at detector(= classical probability result).

If slit 2(or B) blocked off

—> R_1^2 and if slit 1(or A) blocked off, probability —> R_2^2.

Last term(with cosine) -> interference pattern of light and dark bands across detector.

The Interference Term

Whole interference effect relies on

path difference between two routes from slit to screen;

size of path difference not large enough

to affect intensity of electrons arriving at screen from each slit.

So two slits close together

—> assume that $R_1 = R_2 = R$.

Using this —>

$$|\langle x \mid I\rangle|^2 = 2R^2\left[1 + (cos\left(\theta_1(x,t) - \theta_2(x,t)\right))\right]\,|\langle slits \mid I\rangle|^2$$

1st dark band at $\theta_1(x, t) - \theta_2(x, t) = \pi$

Next dark band when $\theta_1(x, t) - \theta_2(x, t) = 3\pi$; another at 5π, etc.

So interference term guarantees

series of dark bands at regular intervals across electron detector.

Theoretical formalism seems to work —> explains this particular experiment.

Spatial dependence of phase given by $\theta = px/\hbar = 2\pi x/\lambda$

If distance x is different for two amplitudes,

then get interference pattern given by

$$A' = Ae^{2\pi i x_1/\lambda} + Ae^{2\pi i x_2/\lambda}$$

Gives intensity or brightness on screen of

$$I = |A'|^2 = |A|^2 \left| \left(e^{2\pi i x_1/\lambda} + e^{2\pi i x_2/\lambda} \right) \right|^2 \quad = 4\,|A|^2 \left(1 + \cos\left(2\pi(x_2 - x_1)/\lambda\right)\right)$$

which agrees with classical result and the experiment for the interference pattern!

So our theory of QM works!

We now proceed as follows:

(1) First, we look at

(a) Time development and how to use QM
(b) Uncertainty Principle
(c) Interferometers; Delayed Choice

(2) We then apply our theory to study

(a) Entanglement
(b) Einstein-Rosen Paradox
(c) Bell's Theorem

(3) Finally we look at the measurement problem

(a) Importance of superposition and entanglement
(b) Schrodinger's cat
(c) What happens in a measurement
(d) Does anything collapse

Some Review, Some New Details and Explicit Time Evolution

Quantum Theory - How it works in general …… And some new details

Question: **How do we predict behavior of physical system using QM algorithms?**

General Approach

1. Identify vector space associated with system

    —> space where all possible physical states of system can be represented

    --> find suitable basis set.

2. Identify operators associated with measurable properties of system

    —> calculate eigenvectors and eigenvalues of observables.

3. Map out specific correspondences

    between individual physical states and individual vectors

    –> decide how to label(what we can measure) state vectors(inside ket).

4. Ascertain present state vector of system by measurements

    —> determine initial labels

    —> state preparation.

5. Time evolution of systems determined by time evolution operator $\hat{U}$

—> **deterministic** equation until next measurement.

$\hat{U}$ is specified for each system.

6. Probabilities of particular outcomes of measurement

carried out at future time calculated by a Postulate

—> probability of measuring eigenvalue b of the observable $\hat{B}$ at t

when in state $|\psi\rangle$ at t given by $|\langle b|\psi\rangle|^2$.

7. Effects of measurement taken into account using a Postulate

—> state collapses to appropriate (as determined by value measured)

eigenvector of measured observable.

8. Then [5]-[7] are just repeated over and over again……that is how you use QM.

Remember:

Postulate —> state vector "collapsed"

into eigenvector of measured observable operator.

Postulate —> how state evolves in time

—> standard or Copenhagen interpretation of quantum theory

due to Bohr, Dirac, Born and von Neumann(he was misunderstood… as we will see!)

Now we present standard way of **talking**(students of physics must master) about superpositions

and deal with some apparent contradictions

that we saw earlier in the discussion of color and hardness measurements:

Right way to think about superpositions of, say, being green **and** being magenta

—> think of them as situations where **color predictions** cannot be made

—> situations where **color talk** is unintelligible.

Talking and inquiring about color of an electron

in such circumstances (in standard view) makes **no sense** whatsoever.

—> if we follow this rule

then earlier contradictions will go away.

—> it is just not so that hard electrons

are not green and not magenta and not both and not neither,

since color talk about hard electrons

—> **no meaning at all**

and same is true for all other incompatible observables.

This is the way the world works according to the quantum physicist!

Once electron is “measured”

—> green or magenta detected,

then **“is”** green or magenta

(color talk now applies) according to standard interpretation.

Measuring color of hard electron,

then, is **not** matter of determining what color of that hard electron is

... it has **none**

it **only** has **probabilities** to have color values

if color is measured.

—> matter of “collapsing” state of measured electron

into one where color talk **applies**,

and “then” determining color of newly created, color-applicable state.

Measurements in QM (in standard view) are very **active** processes.

Not processes of merely learning something about system

—> processes which **drastically** change measured system.

Most important rule:

If going to measure some observable $\hat{C}$,

then choose as basis vectors the eigenvectors of $\hat{C}$ operator

since only with these states does $\hat{C}$–talk makes sense,

i.e., **if discussing color measurements**

than use color basis where color-talk makes sense!

Same holds true if going to measure anything else!

Some Consequences

We are allowed to use **any**

orthonormal set (number = dimension) as basis.

If other sets exist,

then they **must be** eigenvectors of other observables (not color).

Examples….. $Set \quad \#1$

$$|1\rangle = \tfrac{1}{2}|g\rangle + \tfrac{\sqrt{3}}{2}|m\rangle \text{ and } |2\rangle = \tfrac{\sqrt{3}}{2}|g\rangle - \tfrac{1}{2}|m\rangle$$

$Set \quad \#2$

$$|1\rangle = \tfrac{1}{2}|g\rangle - \tfrac{\sqrt{3}}{2}|m\rangle \text{ and } |2\rangle = \tfrac{\sqrt{3}}{2}|g\rangle + \tfrac{1}{2}|m\rangle$$

Both are basis sets

i.e., orthonormal sets

—> operators - let us call observables **direction** d and **truth** t

$\hat{d}$ —> set #1 = eigenstates with eigenvalues +1(up) and –1(down)

$\hat{t}$ —> set #2 = eigenstates of with eigenvalues +1(true) and –1(false)

$$Set \quad \#1$$
$$|1\rangle = |direction = +1\rangle = |up\rangle = \tfrac{1}{2}\,|g\rangle + \tfrac{\sqrt{3}}{2}\,|m\rangle$$
$$|2\rangle = |direction = -1\rangle = |down\rangle = \tfrac{\sqrt{3}}{2}\,|g\rangle - \tfrac{1}{2}\,|m\rangle$$

direction

$$Set \quad \#2$$
$$|1\rangle = |truth = +1\rangle = |true\rangle = \tfrac{1}{2}\,|g\rangle - \tfrac{\sqrt{3}}{2}\,|m\rangle$$
$$|2\rangle = |truth = -1\rangle = |false\rangle = \tfrac{\sqrt{3}}{2}\,|g\rangle + \tfrac{1}{2}\,|m\rangle$$

truth

Remember other basis sets:

$$|color = +1\rangle = |g\rangle = \tfrac{1}{\sqrt{2}}\,|h\rangle + \tfrac{1}{\sqrt{2}}\,|s\rangle \qquad |hardness = +1\rangle = |h\rangle = \tfrac{1}{\sqrt{2}}\,|g\rangle + \tfrac{1}{\sqrt{2}}\,|m\rangle$$
$$|color = -1\rangle = |m\rangle = \tfrac{1}{\sqrt{2}}\,|h\rangle - \tfrac{1}{\sqrt{2}}\,|s\rangle \qquad |hardness = -1\rangle = |s\rangle = \tfrac{1}{\sqrt{2}}\,|g\rangle - \tfrac{1}{\sqrt{2}}\,|m\rangle$$

Assume states represent a real physical system and think about probabilities. —> can make these statements (based on our rules):

1. probability electron is “up” will be measured to be “magenta” = 3/4
2. probability electron is “false” will be measured to be “magenta” = 1/4
3. probability electron is “soft” will be measured to be “magenta” = 1/2
4. probability electron is “green” will be measured to be “hard” = 1/2

and so on.

Do #1:

$$P(up; magenta) = |\,\langle magenta\,|\,up\rangle\,|^2 = |\frac{1}{\sqrt{2}}\,\langle m\,|\,g\rangle + \frac{\sqrt{3}}{2}\,\langle m\,|\,m\rangle\,|^2$$

$$P(up; magenta) = |\frac{1}{\sqrt{2}}(0) + \frac{\sqrt{3}}{2}(1)|^2 = |\frac{\sqrt{3}}{2}|^2 = \frac{3}{4}$$

Now some details and examples —> apply all stuff and expand knowledge and capabilities.

We begin with an example that includes only familiar classical properties

like position, velocity, momentum, energy, etc,

then return to further explain the color-hardness experiments

We know behavior of big particles(large mass)

—> rocks well described by classical mechanics of Newton

—> whatever QM theory says about microworld particles

—> ought to predict that everyday particles, subject to everyday circumstances,

behave in a Newtonian way.

Now position operator $\hat{x}$ and momentum operator $\hat{p}$ are **incompatible**

(i.e., **not simultaneously measurable**)

—> Heisenberg uncertainty principle **exists** (as we will see later)

—> $[\hat{x}, \hat{p}] = \hat{x}\hat{p} - \hat{p}\hat{x} = i\hbar \neq 0$ **commutator**

Since $\hat{x}$ = operator representing a physical observable,

a possible basis of space is set of eigenvectors of $\hat{x}$ operator.

Similarly, for eigenvectors of $\hat{p}$ operator.

and since each are Hermitian operators —> each form a possible basis.

Because corresponding operators do not commute,

a state of definite momentum (eigenvector of $\hat{p}$)

will be a **superposition** of states of definite position(eigenvectors of $\hat{x}$)

or state of definite momentum can only have **probabilities** of having definite x–values.

Similarly, state of definite position (eigenvector of $\hat{x}$)

—> **superposition** of states of definite momentum (eigenvectors of $\hat{p}$).

or state of definite position can only have **probabilities** of having definite p–values.

Vector space when $\hat{x}$ eigenvectors are basis —> coordinate representation.

Vector space when $\hat{p}$ eigenvectors are basis —> momentum representation.

Most important operator in quantum mechanics is the energy operator or Hamiltonian $\hat{H}$

—> will determine time evolution operator as we will see later.

Vector space when $\hat{H}$ eigenvectors are basis —> energy representation.

All are **equivalent** basis sets.

Choice depends on questions being asked or measurements being carried out.

Remember: Choosing an appropriate basis is called "going to HOME SPACE".

—> coordinate-talk, momentum-talk, energy-talk .....

Possible eigenvalues(allowed measured values) of $\hat{x}$ and $\hat{p}$

form **continuum** extending from $-\infty \text{ to } \infty$ (**continuous** spectrum).


Basis is infinite(non-denumerable) dimensional

—> many difficult mathematical problems during discussion in full theory.

However, for simple systems we will discuss, infinite dimension causes **no difficulties**

—> can treat almost all properties as if spaces had finite dimension.

(One exception <—> normalization - only do in an advanced class).



State $|x\rangle$ corresponds to system with definite value of position, namely, x

—> eigenvector of $\hat{x}$ operator.

State $|p\rangle$ corresponds to system with definite value of momentum, namely, p

—> eigenvector of $\hat{p}$ operator.

State $|E\rangle$ corresponds to system with definite value of energy, namely, E

—> eigenvector of $\hat{H}$ operator.

In general, set of energy eigenvalues

are both a discrete set over some range **and** a continuous set over a disjoint range.

Operators position $\hat{x}$ and $\hat{C}$ (color) are **compatible**

—> can simultaneously measure these observables.

—> can say that magenta electron located at x = 7 and that makes sense.

We write state representing this case as $|color - value\rangle\,|x - value\rangle$

i.e., $|m\rangle\,|7\rangle$ for magenta electron located at x = 7. (formal name for product is tensor-product).

When operators act on such states their operation takes the form

$$\hat{C}\,|m\rangle\,|7\rangle = -\,|m\rangle\,|7\rangle \;\text{ i.e., } \hat{C} \text{ only acts on the color space part}$$
$$\hat{x}\,|m\rangle\,|7\rangle = 7\,|m\rangle\,|7\rangle \;\text{ i.e., } \hat{x} \text{ only acts on the position space part}$$

Similar statements hold for

position and hardness, momentum and color, momentum and hardness,

energy and color, and energy and hardness.

All represent simultaneously measurable pairs.

Operator corresponding to observable energy = Hamiltonian $\hat{H}$ where $\hat{H}\,|E\rangle = E\,|E\rangle$

energy eigenvalues

energy eigenvectors

Energy basis = most fundamental in quantum theory.

We will see why shortly.

Now the quantity $\psi_E(x) = \langle x \mid E\rangle$

for state vector |E⟩ = energy **wave function** (just a "bracket" —> nothing mysterious!)

since it takes on different values for each value of x it is **function** of x —> limited use.

Satisfies famous equation that governs time and space dependence in QM,

namely, the Schrodinger equation.

Schrodinger equation in 1-dimension is an "ordinary differential equation" given by

$$-\frac{\hbar^2}{2m}\frac{d^2\psi_E(x,t)}{dx^2} + V(x)\psi_E(x,t) = E\psi_E(x,t) = i\hbar\frac{d\psi_E(x,t)}{dt}$$

Time evolution operator for simple systems

is expressed in terms of Hamiltonian operator $\hat{H}$ by relation $\hat{U} = e^{-i\hat{H}t/\hbar}$

—> **operator function** of $\hat{H}$.

In simple cases, functions of operators are easy to deal with.

If $$\hat{B}\,|b\rangle = b\,|b\rangle$$

i.e., |b⟩ = eigenstate of $\hat{B}$, then have

$$f(\hat{B})\,|b\rangle = f(b)\,|b\rangle$$ for function of operator.

—> operator argument is replaced by eigenvalue inside the function

—> energy eigenvectors have simple time dependence or time evolution, i.e.,

$$|E,t\rangle = \hat{U}\,|E,0\rangle = e^{-i\hat{H}t/\hbar}\,|E,0\rangle = e^{-iEt/\hbar}\,|E,0\rangle$$ **just replace operator with eigenvalue**

where complex exponential form —> $$e^{-iEt/\hbar} = \cos\frac{E}{\hbar}t - i\sin\frac{E}{\hbar}t$$

Using eigenvectors/eigenvalues of Hamiltonian, namely, $$\hat{H}\,|E_n\rangle = E_n\,|E_n\rangle$$

as basis —> time dependence of arbitrary state vector can be derived as follows:

This is one of the major techniques used in QM calculations.

Initial state $|\psi(0)\rangle$

Expand in energy basis (HOME space of time-evolution operator)

$$|\psi(0)\rangle = \sum_n c_n |E_n\rangle$$

Operate with time evolution operator

$$|\psi(t)\rangle = \hat{U}(t)|\psi(0)\rangle = \hat{U}(t)\sum_n c_n |E_n\rangle = \sum_n c_n \hat{U}(t)|E_n\rangle$$

Now

$$\hat{U}(t) = e^{-iHt/\hbar}$$

Therefore

$$|\psi(t)\rangle = \sum_n c_n e^{-i\hat{H}t/\hbar}|E_n\rangle$$

and we finally get (substitute eigenvalue for operator)

$$|\psi(t)\rangle = \sum_n c_n e^{-iE_n t/\hbar}|E_n\rangle$$

Short digression for the mathematically inclined(otherwise just sit and listen) to show you that you can actually already do real QM calculations.
Now repeat some earlier algebra for fun(to see it really is not difficult)

In the discrete spectrum case, we have, using a basis set {|n⟩}, that

$$|\psi\rangle = \sum_n c_n |n\rangle \quad , \quad c_m = \langle m | \psi \rangle$$

so that

same

$$|\psi\rangle = \sum_n \langle n | \psi \rangle |n\rangle = \sum_n |n\rangle \langle n | \psi \rangle = \left(\sum_n |n\rangle \langle n| \right) |\psi\rangle$$

same

which says that

$$\sum_n |n\rangle \langle n| = \hat{I}$$

identity operator

as derived earlier

and if

$$\hat{A} |a_n\rangle = a_n |a_n\rangle \quad , \quad |\psi\rangle = \sum_n c_n |a_n\rangle$$

then

$$\hat{A} |\psi\rangle = \sum_n c_n \hat{A} |a_n\rangle = \sum_n c_n a_n |a_n\rangle$$

now using clever math trick

$$\sum_n c_n a_n |a_n\rangle = \sum_n c_n \sum_m a_m |a_m\rangle \delta_{nm} = \sum_{n,m} c_n a_m |a_m\rangle \delta_{nm}$$

and substituting $\delta_{nm} = \langle a_m | a_n \rangle$ we get

$$\sum_n c_n (\hat{A}) |a_n\rangle = \sum_{n,m} c_n a_m |a_m\rangle \langle a_m | a_n\rangle = \sum_n c_n \left(\sum_m a_m |a_m\rangle \langle a_m| \right) |a_n\rangle$$

so that

$$\hat{A} = \sum_m a_m |a_m\rangle \langle a_m|$$

as derived earlier

which is **spectral representation** of operator $\hat{A}$ in terms of its eigenvalues and eigenvectors.

Now you are learning math language and we are beginning to get somewhere!

Next derivation more mathematical, but very important. **Result is important thing!**

Now we have all needed tools.

General procedure for figuring out time dependence of states

and then answering questions posed in experiments

goes as follows (go slowly — tricky algebra - j**ust follow along**):

1. Ask experiment/theoretical question — the typical form is

- if system in state $|\phi\rangle$ at t = 0, what is probability of system being in state $|b_7\rangle$ at t?

2. Assume $|b_7\rangle$ one of eigenvectors of measurable operator $\hat{B}$.

3. Solve for energy eigenvectors of system,

i.e. find Hamiltonian (energy operator) and do mathematics.

4. Write $|\phi\rangle$ in terms of energy eigenvector basis $|\phi\rangle = \sum_E c_E |E\rangle$ where $c_{E'} = \langle E' \mid \phi\rangle$

component

5. Since time dependence of energy eigenstates is easy,

can write down time dependence of state $|\phi\rangle$ as

$$|\phi, t\rangle = \sum_E c_E |E, t\rangle = \sum_E c_E e^{-iEt/\hbar} |E\rangle$$

it is that easy!

6. Write |E⟩ in terms of $\hat{B}$ eigenvector basis $|E\rangle = \sum_j d_j \, |b_j\rangle \;\text{ where } d_{\mathrm{k}} = \langle b_k \mid E\rangle$

7. Write |ϕ, t⟩ in terms of $\hat{B}$ eigenvector basis

$$|\phi, t\rangle = \sum_E c_E e^{-iEt/\hbar} \, |E\rangle = \sum_E \langle E \mid \phi\rangle \, e^{-iEt/\hbar} \sum_j \langle b_j \mid E\rangle \, |b_j\rangle$$

8. Then probability of finding b_7 at time t given by(a postulate)

$$P = |\, \langle b_7 \,|\, \phi, t\rangle \,|^2$$

We get

$$P = \left| \langle b_7| \left(\sum_E \langle E \mid \phi\rangle \, e^{-iEt/\hbar} \sum_j \langle b_j \mid E\rangle \, |b_j\rangle \right) \right|^2$$

or

$$P = \left| \left(\sum_E \langle E \mid \phi\rangle \, e^{-iEt/\hbar} \sum_j \langle b_j \mid E\rangle \, \langle b_7 \mid b_j\rangle \right) \right|^2$$

Now

$$\langle b_7 \mid b_j\rangle = \delta_{j7}$$

This means that the sum over j disappears and all j's get replaced by 7 . We get

$$P \quad = \left| \sum_E \langle b_7 \mid E \rangle \langle E \mid \phi \rangle e^{-iEt/\hbar} \right|^2$$

Final result expresses answer

in terms of initial and final states

and properties of energy eigenvectors and energy eigenvalues,

and all are known!!

Thus after several passes(to better understand things) through various topics involved,

we see that our formulation of quantum theory

is capable of making the necessary predictions.

We won't be doing this in general(that is for a regular QM course),

but it is important to know that we can!

Some more mathematical material is on the website in the the supplementary readings.

We now look at some intriguing experiments exhibiting QM phenomena in order to get an idea of how the strangely quantum world behaves and to also start to learn about non-locality which will be a central to understanding measurement.

Now to look at some important experiments —> give us hints as to how QM works!

An Interference Experiment with Photons

Direct laser beam at a half-silvered mirror.

For **intense** light beams,

such mirrors reflect half of light and allow half to pass straight through.

When intensity of laser beam high,

two beams are seen emerging from mirror,

each having 1/2 intensity of incoming beam.

Arrangement is called a **beam-splitter**.


If turn intensity of laser down(photons emerge with time gaps)

-> only one photon around at any given time,

and use pair of C(harged)C(oupled)D(evice) detectors

to detect reflected and transmitted photons,

something very interesting happens.



For each photon that leaves laser,

one photon is detected either at transmission CCD (on transmitted path)

or at reflection CCD (on reflected path).

Photons are not split in some odd manner

so that half photon goes one way at mirror and half other way. See figure.

Instead, seems to be 50:50 chance(probability)

that photon transmitted or reflected by half-silvered mirror.

No measurable difference between photons as they approach mirror,


i.e., no property seems to determine which way they will go (sound familiar).

—> Fundamental point that will come up repeatedly in context of quantum theory.

Next step is to remove detectors and replace them with two mirrors (fully silvered)

that divert two beams(change their directions by 90°)

to a second half-silvered mirror as in figure.



There are now two beams arriving at the 2nd half-silvered mirror.

One from each path through 1st silvered mirror.

At this point,

same thing happens at 2nd silvered mirror,

with 1/2 of light arriving at 2nd half-silvered

mirror passing straight through

and other half being reflected.

Two new beams emerge

and eventually travel to pair of detectors

placed at X and Y. See figure

Beam heading to detector X

is combination of light that was reflected by 1st half-silvered mirror (travelled top path),

then transmitted by 2nd half-silvered mirror,

with light that transmitted by 1st half-silvered mirror(along bottom path)

and reflected by 2nd.


Detector Y collects light that is similar mixed combination.

Arrangement of mirrors and detectors is called **Mach-Zehnder interferometer**.

Once set up, it is easy to confirm that

intensity of light reaching each detector depends critically

on **distances travelled** by light along top and bottom paths.



If equipment is finely adjusted

so that paths are exactly same length,

detector Y records no light

while detector X records all of intensity (all photons) entering experiment.

Without critical adjustment,

X and Y collect light in varying amounts:

more light at X -> less reaches Y (and vice versa).

Using classical physics methods these effects can be completely explained by saying that light is a **wave —- but this argument works only when the beam intensity is high**.

Interference as Wave Effect (reminder)

Consider ripples crossing surface of lake.

Ripples consist of places where water level is higher than normal (peaks)


and places where dropped below normal (troughs).

Wavelength of ripple is distance between successive peaks(P),

same as distance between successive troughs(T).

Frequency of wave is rate at which complete cycles (peak to trough to peak again)

pass fixed point, and period is time taken for one cycle.



Light(E/M fields) is more complicated than a water wave.

Peaks and troughs of light wave are not physical distances (like height of water wave)

but are variations in strength of fields(E and B).

Mathematically, however, they are the **same** phenomena since the equations are identical.

Thus, light waves are very sensitive measures of distance.

In interference experiment with interferometer,

divide distance travelled by light wave on route to detector into sections,

each having length equal to wavelength of wave.

Distance probably not whole number of wavelengths.

Furthermore, two different possible routes through experiment

have to be precisely same length to be precisely same number of wavelengths long.

If distances not precisely same,


light traveling along each route consequently will

have gone through different number of complete waves when gets to the detector.

As light in two beams has common source at first half-silvered mirror,

two beams will set off on different routes in phase

(i.e., in step - simplest definition of phase - both at same point on wave) with each other.



Waves labelled A and B in phase(peak to peak),

waves B and C exactly out of phase (peak to trough).

Above statement ->> set off peak for peak -

so were in phase.

By time they get to detector

two beams may no longer be in phase (different distances travelled).

One could be arriving at peak, and other at trough (B and C).

If this happens, then waves will cancel each out —> no energy entering detector

called **destructive interference**

If still in phase would add up

called **constructive interference**).

Exact cancellation only happens if waves meet precisely peak to trough

not possible for any extended length of time

due to small variations in distance (mirrors shaking slightly)

and fluctuations in laser output.

Detailed analysis of interference experiment

also takes into account what happens to light at various mirrors

-> also influence phase of waves.

When light reflects off mirror,

reflected wave out of phase with incoming wave by half wavelength.

Using λ = wavelength, wave has undergone λ/2 phase shift (shift by 1/2 wavelength)

on refection.

Slightly different with half-silvered mirror,

surface that reflects from either side mounted on thin block of glass.

Dashed line in figures indicates reflecting surface.


If reflection takes place off surface before light enters glass block,


then ordinary λ/2 phase shift takes place.

However, any light that passes through block

before reaching reflecting surface not phase shifted on reflection.

See two figures below.





Two light beams reaching detector Y will be (overall) λ/2 out of phase.

Consequently, waves B and C and will completely cancel (if travelled equal distances).

If carry out same analysis for detector X,

i.e., chart progress of waves through interferometer to detector X,

find arrive in phase provided travelled equal distances.

Most experimental setups,

paths through interferometer not exactly equal

so waves not exactly in/out of phase.

Consequently, some light reaches both X and Y.

If equipment allowed movement of fully silvered mirrors,

so relative path lengths were changed,

then variation in brightness of light in X and Y

could be studied as mirror moved - as relative paths changed.

That is the classical way of thinking using interference.

Modern version of Mach-Zehnder interferometer with Single Photons

Suppose we are able to turn down laser intensity

so that light beam made up of single photons entering apparatus at any time.

Assume have done so —> only one photon in experimental apparatus at any time.

Also have control over average rate. Note that “interference” argument is no longer valid!!

Expect photons arriving at half-silvered mirror

to have 50:50 chance of going through/reflecting off.

Another possibility is

two reduced energy photons emerge from mirror, in each direction.

Can determine experimentally what happens:

place photon detectors just after mirror in path of each possible beam.

Simple experiment produces interesting result.

Half the time, photon is reflected, and half the time it is transmitted;

and never get two photons coming out at same time.

However, no inherent difference

between those entering photons that get through and those that reflect.

No pattern to sequence,

except that after long time half reflect and half get through.

Sound familiar!

Effect is common in quantum physics.

Some aspects of nature's behavior lie beyond ability to predict (e.g., which way photon will go).

Question - does this reflect fundamentally random aspect to nature,

or is something more subtle going on that have not discovered yet?

Having established that photon reaching

1st half-silvered mirror in Mach-Zehnder interferometer

will either reflect and travel top path through device,

or transmit and follow bottom path,

Now turn attention to what happens at detector end of device.

First find, between them, detectors pick all photons that enter experiment.

Number of photons arriving at either detector in given time depends on two path lengths,

i.e., if exactly equal then no photons ever arrive at Y.

If paths not exactly equal, then find that detection rate at each detector

reflects intensity of interference pattern

that would be observed when intensity is turned up.

What do we mean by that?

Let's imagine that had arranged for path lengths

such that 70% of total light intensity entering experiment arrives at X and 30% at Y.

No double photon firings.

Experiment done under well-controlled conditions and no doubt

that photon arrival rate directly reflects an interference pattern between paths.

Doesn't sound like a problem, but there is.

If photon is small particle of light,

then how can different path lengths have any effect on one single photon?

Confirmed that photons randomly choose reflection or transmission at half-silvered mirror.

After that, surely they proceed along one path or other to detector.

Hard to imagine single photon going along both paths at same time.

Remember that is rejected by experimental results

(detectors only registered one photon at a time).

Now a wave can do this.

Can spread out throughout experiment (ripples on lake)

so that parts of wave travel along each path at same time

(i.e., wave energy divides between paths).

When two parts of wave combine at far side of experiment,

information about both paths is being compared,

which leads to interference pattern.

A single photon however, must surely have information about only one path,

so how can single photon experiments produce interference patterns?

Flaw in arguments.

The flaw in these arguments is extremely subtle

and leads to the primary issue physicists face when dealing with quantum world.

We confirmed that photons divert at half-silvered mirror by placing detectors in two paths.

However, doing this eliminated any chance of picking up an interference pattern.

If detectors have stopped photons, then they **have not** travelled any paths.

In principle, experiment does not tell anything about what happens when no detectors present.

Common sense to assume that photons do same thing with or without detectors,

but as already seen,

the interference pattern for photons(all of QM) is not a matter of common sense.

In addition, color/hardness experiments

say that it is important whether or not detectors are present!

Must investigate this further.

Place one photon detector after half-silvered mirror - in path of reflected beam.

If detect photon there, then would not get one at far side of experiment.


On other hand, if do not pick one up at detector

then it has passed through mirror rather than reflecting and should see it at far end.

Experiment is easily done and confirms

that for every photon leaving the laser the detectors pick one up either at far end

or in reflected beam as below.



Find for transmitted photons

half of arrive at Y and other half at X,

no matter what the length of paths is.

In other words, no interference takes place.

Removing detector on reflected path

(and replacing the corresponding half-silvered

mirror) opens up that route

to far side of experiment again.

At same time

it removes any direct knowledge that we might have

about behavior of photons at the half-silvered mirror.

We observe that doing this restores the interference pattern!

Remember what happened in the color/hardness experiments.

Summarizing the logic so we can expose what is happening and find the flaw.

(1) Rate of photons arriving at far side of experiment related to intensity of bright beam.

(2) Moving mirror with bright beam maps out interference pattern in detectors.

(3) Reducing intensity of beam does not affect interference pattern - instead arrival rate of photons now depends on position of mirror

(4) If set up experiment so that can tell which path taken by photon (directly or indirectly), then interference pattern is destroyed.

(5) If unable to tell paths of photons, then there is interference pattern, implies photons arriving have information about both routes through experiment.

(6) Opening up top path (removing detector) can reduce number of photons arriving at Y.

In fact, if path lengths are same, opening up top path means never get any photons at Y.

All experimental results equivalent to Young-type two-slit interference experiment

and the two-path color-hardness experiment we discussed earlier.

Changing experiment to a Delayed Choice configuration makes result even stronger!


Let us see how.

Introduce Pockels cell(PC) (can divert photons extremely fast) on one route. Details later.

—-> Trying to **find out what photons are doing while in the interferometer.**

Consider setup where PC set to divert photons(to extra detector).

Photon leaves laser and arrives at first half-silvered mirror.



If reflected,

then PC will divert it so don't see at X or Y.

However, if photon transmitted

by first half-silvered mirror,

misses PC, and turns up at either X or Y.

Either case

no interference pattern

(we have gotten which-path information).

If set PC to pass photons,

then changes what happens and get interference pattern.

For extreme case of equal path lengths, no photons ever arrive at Y.

Assume two path lengths exactly same. So have:

(1) If PC set to transmit, then get no photons at Y and all at X.

(2) If PC set to divert, then half of photons detected have equal chance of either X or Y.

This result should make us stop and think.

If photon takes lower route with PC set to divert,

then can get to X or Y.

If takes lower route with PC set to pass,

then photon can never arrive at Y.

But if takes lower route it doesn't go anywhere near PC,

so how can setting of that device affect things?

Is this further hint that somehow or other photon travels both routes at same time?

Again this should sound familiar!

Now experimentalists get very devious. Experimentalists are very clever!!

Set PC to divert photons, but while photon in flight(during experiment),

switch cell over to transmit setting.

Ability to do this very rapidly means can make change

after photon has interacted with first half-silvered mirror.

No magic in doing this. If know when photon left laser,

can estimate how long takes to get to half-silvered mirror.

Provided we switch PC after this time,

but before photon had time to reach detectors X and Y,

then can perform experiment as described.

If setting of PC has(in someway) influenced the photon,

then original setting should have determined that photon takes one path or other

and certainly not both at once.

Now think we have changed setting after decision(at 1st 1/2-silvered mirror) made by photon

NOTE: Of course, word decision is not appropriate here.

Photons do not make decisions.

Clearly, hard not to be anthropomorphic when describing experiments.

In fact, can trigger PC in random manner.

Record setting of PC and match arrival of photons at one detector or another.

Can then run experiment for many photons

and record arrival at different detector positions.

After experiment run for while,

can analyze data.

Have some photons arriving at Pockels detector

and some at far end of experiment.

Latter group sorted out into those that arrived

when PC set to divert,

and those that made it when PC set to transmit.

Remarkably, when data separated,

photons that arrived at far side with PC set to transmit show interference pattern.

Other photons that arrived with PC set to divert

(obviously committed to other path and so missed it) show no interference pattern.

In every case PC set to divert photons and switched after photon left first mirror.

With PC set to divert, photons follow one route or other.

Then switched PC, destroying ability to know which path photons travelled,

and producing interference pattern.

Hard to believe that changing setting of PC can have an influence

that seems to **travel backward in time(retrocausality)** to when photon reaches first mirror.

Last statement, was made using ordinary words, not mathematics(nothing strange is happening) —> our attempt to use words may be generating total nonsense of course!

The mathematics is clear. QM makes the correct predictions in all cases!

From both photon experiments and color/hardness experiments

we have seen that quantum physics is a contextual theory

-> adequate description of behavior of quantum object (light/electrons)

requires an understanding of whole (the context) experimental setup.

Quantum behavior depends on **context**!

Experimental results (answers to questions) **depend** on what questions we are asking!

Another Interference Experiment

Clearly, we get into difficulty with experiments

when we piece together an understanding of the “whole”

by looking at the component parts on their own.

When everything is together, things behave differently. **—> whole > sum of parts!!**

However, results of one experiment are not an accurate guide to another.

If this is case,

then conclude that photons always take only one route,

as indicated in experiments that looked for route followed.

However, know that as soon as do not have such ability

to tell path of photons

they seem to take both routes at once

or something that is equivalent to that,

or maybe nothing we can imagine with our macroworld minds/words!


Another experiment we now look at, pushed this notion further

by showing interference pattern can be destroyed

without directly influencing the photons creating it.

Experiment uses crystal known as down-shifter.

Device absorbs 1 photon to produce 2 new photons, each with half the original energy.



2 signal beams directed to detector

Laser light is sent onto 1/2-silvered mirror

and 2 beams separately

directed into down-shifters A or B.

Each down-shifter produces a

signal beam and an idler beam,

The difference between two beams is nothing

more than **subsequent** way they are used.

produce interference pattern (different path lengths)(same as Mach-Zehnder).

Idler beam from down-shifter A (now too weak to activate down shifter B) is mixed with that from down-shifter B

and both idler beams arrive at second detector.


Upshot is that every time photon leaves the laser,

it is observed that a photon of half energy arrives at each detector.

Fact that interference pattern emerges

-> in some manner,

that each photon appears to have travelled along both signal beam paths.



Say photon arrives at half-silvered mirror

and goes on top path (only that path).

Photon arrives at down-shifter A

and produces 2 further photons,

one ends up at signal detector

and other at idler.

No interference pattern

since no information carried to signal detector about other route.

Same true if photon took lower route through experiment.

Only way to get interference pattern at signal detector

is for information to arrive from both routes,

i.e., to have to be two signal beams, one from each down-shifter.

If true,

then down-shifters have to be activated

by something arriving at each one —->

makes it appear that photon from laser went both ways at half-silvered mirror.

Presence of two signal beams

doesn't imply that two photons are arriving at signal detector

i.e., measurement show that there is **only one at a time arriving**.

Most bizarre feature of experiment

is way in which interference pattern can be destroyed


if we have ability to tell path of the photons, even if don't choose to use this information.

Threat of doing this, or rather fact that

experiment(context) is set up to allow possibility, is enough to destroy interference!

Dramatically confirmed by blocking one of idler signals, say from down-shifter A

Logic here remarkable.



Whenever photon picked up

at idler detector we know must have come

from down-shifter B.

Means that photon from half-silvered mirror

must have hit down-shifter to be converted

into photon that arrives at idler detector.

Can further deduce that other photon

from down-shifter B travelled to signal

detector and therefore is photon detected there.

Arrivals correlated!


Tracing argument further back,

photon that definitely hits down-shifter B

must have come from half-silvered mirror.

There is no ambiguity about route that this

photon takes from mirror.



Nothing goes along top route; nothing produced from down-shifter A,

so interference pattern disappears.

As long as idler route from down-shifter A open,

have no way of telling which shifter a photon came from.

Ambiguity sufficient to guarantee interference pattern at signal detector.

If don't know that photon at idler detector came from B (or A),

then don't know about signal photon either.

Under those circumstances, can't say which route photon took at half-silvered mirror,

so takes both routes or "something" like that.

Seems that behavior of photon determined by **context** of experiment as whole.

Know no photons coming from down-shifter A,

why does it matter that idler route from A is blocked?

How is information conveyed back to half-silvered mirror

so as to determine what happens there? **Words FAIL us, but QM MATH works!!**

Quantum Erasers (Complementarity and Entanglement)

In earlier discussions, we saw that closing the locality loophole involved switching between different analyzer orientations while emitted photons were still in flight.

The choice between the nature of the measurement was therefore delayed with respect to the transitions that originally created the photons.

Is it possible to make this delayed choice between measuring devices of a more fundamental nature?

For example, in discussion of double-slit measurements, found that if one allows sufficient a number of photons individually to pass through slits, one at a time, interference pattern will be built up.

It **seems** that photon has knowledge of all the paths as though it passes through both slits.

As noted earlier, a skeptical physicist who places a detector over one of slits to show that photon passes through one or other does indeed prove their point - photon detected, or not detected, at one slit.

But then the interference pattern can no longer be observed - because we have which-path information.

Advocates of local hidden variable theories could argue that the photon is somehow affected by the way we choose to set up our measuring device.

It thus adopts a certain set of physical characteristics (owing to existence of hidden variables) if apparatus is set up to show particle-like behavior, and adopts different set of characteristics if apparatus is set up to show wave interference.

However, if we design an apparatus that allows us to choose between these totally different kinds of measuring device, we could delay our choice until photon was (according to local hidden variable theory) **committed** to showing one type of behavior.

Suppose that photon cannot change its mind **after** it has passed through slits, when it discovers which kind of measurement is being made (whatever last sentence given in “words” may actually mean!!)

Ultimate Delayed Choice or Quantum Eraser Experiment

QM says systems can change behavior depending on measurements made on them or in response to decision that has not yet been made.

One part of an entangled pair can affect properties of its partner instantaneously, no matter where in universe its partner happens to be.

A so-called quantum eraser experiment has now been done...it dramatizes several aspects of this quantum strangeness at once.

Experiments dramatically show non-local effects, i.e., ability of experiment in one place to somehow influence outcome of another regardless of time or distance – but without transmitting any signals.


The idea behind the quantum eraser is to make paths(like those in a 2-slit system) distinguishable, which eliminates any interference effects, but then erase the which-path information just before light reaches screen where we actually observe the interference pattern.

QM predicts that interference pattern should then reappear and it does.

Consider a quantum mystery of the following type. using words always ends too anthropomorphic

A photon approaching slits will “need to know” whether or not there is an eraser further down path(in its future), so that it can decide whether to pass through slits as a superposition of all possibilities (paths are indistinguishable) and produce an interference pattern later on screen or that should behave as if paths are distinguishable and produce no interference pattern later on screen!!


An early eraser experiment can be visualized as the two-slit experiment as shown in diagram below:



Photons passing through double slit are all vertically polarized.

By two paths as shown they can get to a recombination crystal, which remakes(without changing properties) beam that produces an interference pattern on screen, i.e., if paths are indistinguishable, then have superposition of all possible paths(2 paths in this case) and get an interference pattern.

Now insert a polarization rotator on one path only




and rotate the polarization of photons to horizontal in that path.

Since the paths now produce distinguishable photons, we get particle like behavior and the interference should disappear.

It does!

Now add the quantum eraser after recombination crystal —> a polaroid at 45°.



The polaroid filter produces equal numbers of photons with both vertical and horizontal polarization (with respect to the new direction).

Half of new vertically polarized photons come from horizontal polarized photons on top path and half of new vertically polarized photons come from vertical polarized photons on bottom path (similarly for new horizontally polarized photons).

It is impossible to tell whether photon was vertically or horizontally polarized before the eraser.

Thus, the two paths are once again indistinguishable.

If the interference pattern reappears, then the photon approaching slits somehow needs to know whether or not there is an eraser down line so can decide whether to pass through slits as superposition and produce interference effects or as mixture and produce no interference.

Again, using words here makes things sound silly!! QM mathematics is clear!!

The pattern reappears immediately thus confirming the quantum mechanical prediction.

The filter erases the which-path information caused by the rotator.

A truly astounding result.

What does this say about the classical idea that it is the two-slit system that is the real cause of the interference pattern?

Worry:

Experimenters, knowing the fundamental importance of these results, wanted to leave no possible source of controversy intact.


In the above version of experiment, there is a potential problem because the which-path information is carried by same photons that interfere, making experiment difficult to interpret.

New version of experiment...

Here which-path information is not carried by what one would naively call the interfering photon.

Instead, carried by 2nd photon and along the way it also demonstrates the striking non-local effects of QM.

Experiment looks like:



High intensity laser photons are sent into a parametric down-conversion non-linear crystal such as lithium iodate.


The crystal converts some incoming photons(green) into pairs of identical photons with lower energy and vertical polarization moving at an angle to original direction.

The photons are produced as entangled partners (in a superposition).

Measurement on one photon automatically tells us about the other with no direct measurement on 2nd photon necessary.

Twin beams exit the crystal at angle to the original path.

These are the thick red and blue lines



rotator not in place yet


They are reflected back towards the crystal by mirrors and pass straight through it to detectors (the second-pass intensity is now too weak to cause down-conversions).

These are thick red and blue lines.

However, not all laser light gets converted on 1st pass.

Some goes straight through crystal to another mirror(green) and is reflected back into the crystal(still high intensity) where the crystal creates more photon pairs which then follow same path as other beams to the detectors.

These are thin red and blue lines.




As a result, have two different beams heading towards each detector each having follow different paths.

Each pair of beams corresponds to a **separate** double slit experiment.

If there is no way to distinguish photons created on first pass through crystal from those created on 2nd pass —> both detectors should have interference patterns and they do!!

Now make one returning beam in one leg distinguishable from other by inserting polarization rotator into red path(as shown below) converting vertical to horizontal polarization in that leg.




The interference pattern in top detector vanishes instantly as it should since the two (interfering) beams now have distinguishable paths.

Now, however, also find that interference pattern disappears in bottom detector!!!!!!! Why?

We have done nothing to disturb these beams so that beams in the bottom detector still correspond to indistinguishable paths and photons!!!!

Remember, however, that photons are created in entangled pairs, so when red-path photons become labeled with which path info, the same info becomes available to blue-path photons, no matter where they are!!!

This is the concept of non-locality at work.



To erase which-path info, we now add a 45° polaroid in red path, just in front of detector.

The interference pattern immediately reappears in the top detector as it should (same as previous experiment).

It seemed however, that pattern did not reappear in bottom detector.

One might imagine this is so because erasing red-path photon information does not erase any information from blue path.

In addition, if it did reappear immediately, then we would be able to send signals faster than light with such an eraser.

However, as in EPR(we will see) experiment we find, if we bring the two data sets back together and compare them, the pattern had, in fact, been restored along both paths (need both data sets) to see the **correlations** among individual photons.

It seems like experiment is saying that the **correlations are entangled** and **not the state vectors**!!

That is the key piece of information that we have been looking for as we will see!!!!

We will have more to say about this later when we talk about measurement.

Alternatively, one can do a coincidence measurement, which only looks at those photons counted in each detector simultaneously and one can see interference pattern return directly, that is, which way and interference effects are being recorded for single photons.

Thus, inserting or removing which-path information transforms behavior of light throughout entire system simultaneously demonstrating amazing quantum eraser and dramatic non-local behavior of QM.

This experiment makes it clear that there is direct relationship between tests of complementarity(the concept that two contrasted theories, such as the wave and particle theories of light, may be able to explain a set of phenomena, although each separately only accounts for some aspects) and tests of quantum non-locality.

Interference effects are a direct manifestation of non-local behavior.

These effects can be encoded in the mathematical structure of quantum entanglement - in this case, the entanglement of the states is responsible for the interference with the state used to detect the "which way" information.

These states cannot be disentangled without forcing the system to reveal one type of behavior or other.

They cannot be disentangled to reveal both types of behavior simultaneously.

Though still a subject for debate, consensus is building that complementarity - and hence non-locality and entanglement - is the mechanism for mutual exclusivity in the dual wave-particle nature of quantum objects - what Richard Feynman described as central mystery at heart of quantum mechanics.

Uncertainty Principle

The Heisenberg Uncertainty Principle is at the center of quantum theory —> guarantees the consistency of the QM view of the microscopic world.

Yet there is a puzzle with Heisenberg's discovery:

> **Is it a statement of an epistemological limitation (what we can measure about world) or an ontological limit(fundamental property) written into world itself?**

Let us step back first and get some background ideas.

Expectation is Not Enough

Take a collection of quantum objects, all in same state $|\phi\rangle$ and measure the value of physical property(call it O) for each —> collection of data values $\{o_i\}$, i = 1,2,3,4,……

Find average value of measurements (= to expectation value of operator $\hat{O}$ associated with property O in this state).

Expectation value, however, does not tell whole story.


Also need to know how wide is range of values - how spread out around average.

Collection $\{o_i\}$ could all be very close to average value or some of

measurements might produce values considerably different

from average, not because of experimental mistake,

just due to nature of state being measured.



Whenever physicists think about a perfect set of experimental values —> **normal distribution** curve, as shown.




Figure —> results of identical measurements on collection of objects.


Normal distribution applies to all experimental results..

x–axis —> values of results obtained and y–axis —> number of times value turned up.

—> result 5 obtained 10 times and average value over all readings = 5.

However, exist quite a few times when value significantly different from 5 obtained!

Now consider figure.




—> normal distribution curve, but much narrower spread of values (peak still at 5, average still = 5).

Much higher proportion of results closer to average than before.

—-> To tell full story of set of measurement results, need more than average - also need width(spread) of distribution as well.

Width obtained by taking each value, working out how far from average and averaging results.

Mathematicians —> **standard deviation**; quantum theory —> **uncertainty**. See figure.



In classical physics, spread, or uncertainty in experimental results generally attributable to normal experimental variations one finds due to small differences in objects being tested.

Also can be attributed to variations in apparatus used or environmental factors.

Experiments in quantum physics suffer from same variations, but in addition have to face what happens when objects being tested are not in eigenstates of variables being measured.

Taking position as an example. Start with state $|\phi\rangle$

$$|\phi\rangle = \sum_x \langle x \,|\, \phi\rangle \, |x\rangle \qquad \phi(x) = \langle x \,|\, \phi\rangle$$

NOTE: I am simplifying by letting x have discrete values!

—> making position measurements on set of particles in this state we will find a range of values, with each possible x value occurring according to probability

$$\phi^*(x)\phi(x) = |\phi(x)|^2$$

ϕ(x) = expansion coefficient = component

Plotting results and number for each value —> will draw out $|\phi(x)|^2$ curve with corresponding uncertainty Δx -> measurement distribution

Uncertainty —> direct measure of range of x values over which amplitude $\phi(x)$ is significant —> range of x values highly likely to occur when one makes a measurement.

Calculations then yield the uncertainty.

$$\langle\phi|\,\hat{x}\,|\phi\rangle = \langle x\rangle \qquad \textbf{expectation value}$$

$$\Delta x = \sqrt{\langle\phi|\,\hat{x}^2\,|\phi\rangle - \left(\langle\phi|\,\hat{x}\,|\phi\rangle\right)^2} = \sqrt{\langle x^2\rangle - \langle x\rangle^2} \qquad \textbf{uncertainty}$$

Shows that Δx is as much a property of state as $\langle x\rangle$.

Now, same state could be expanded over momentum basis $|p_x\rangle$, and experimental results of momentum measurements plotted —> uncertainty Δp_x.

What about case where we have an eigenstate?

Well, if happen to start with eigenstate, then can predict with absolute certainty what result of measurement of will be.

Plot of results —> no width (aside from unavoidable experimental fluctuations), giving $\Delta p_x = 0$.

However, there is a catch.

Momentum eigenstate cannot also be position eigenstate and vice versa.

When we expand a momentum eigenstate($\Delta p_x = 0$) over position basis —> infinite number of possible positions, leading to $\Delta x = \infty$ —> if $\Delta p_x = 0$, then $\Delta x = \infty$ and vice versa!

Heisenberg's Principle

What about states (majority) that are neither position nor momentum eigenstates?

Given such state $|\phi\rangle$ have

$$|\phi\rangle = \sum_x \phi(x)\,|x\rangle \Rightarrow \langle x\rangle, \Delta x \quad , \quad |\phi\rangle = \sum_p \phi(p)\,|p\rangle \Rightarrow \langle p\rangle, \Delta p$$

i.e., position representation —> $\langle x\rangle$ and Δx and momentum representation —> $\langle p_x\rangle$ and Δp_x.

Since it is **same state** expanded in both representations there must exist a connection between Δx and Δp_x.

Connection was first demonstrated by Heisenberg.

THE HEISENBERG UNCERTAINTY PRINCIPLE $\quad \Delta x \Delta p_x \geq \dfrac{\hbar}{2}$

—-> for given state, the smaller the range of probable x values involved in position expansion, the larger range of p_x values involved in momentum expansion, and vice versa.

Key part of expression is ≥ sign —> product $\Delta x \Delta p_x$ **cannot be less than** $\hbar/2$.

Measurement results might be such that $\Delta x \Delta p_x$ is much greater than $\hbar/2$, which is fine; but can never find situation where $\Delta x \Delta p_x$ less than $\hbar/2$.

Quantum theory places a **limit on the precision** of two sets of **related** measurements.

Saying this a different way:

Δx is telling us range of x values where $\phi(x)$ is significantly large if we expand state over the position basis.

Similary, Δp_x is corresponding range of $\Psi(p_x)$ if expand same state over momentum basis.

Uncertainty principle is relating range of $\phi(x)$ to that of $\Psi(p_x)$.

Make position measurement, then most likely get value within Δx of expectation value $\langle x \rangle$.

Make momentum measurement **instead (not after)**, then most likely get value within Δp_x of $\langle p_x \rangle$.

In last statement, the "not after" part is important.

Making position measurement, **"collapses"** state into position eigenstate —> different momentum expansion than original state had.

Uncertainty principle only relates Δx and Δp_x for **same state**.

Here is possible question about the principle.

What about object that is not moving?

Is it not sitting perfectly still (precise momentum, zero) at a precise spot?

Well, ignoring idea that nothing can be perfectly still (orbiting sun, etc), the immediate point is that uncertainty principle **prevents** such state from happening in microworld.

So What?

People drew philosophical conclusions from uncertainty principle immediately.

Heisenberg targeted the principle of causality.

Classical physics —> view that powerful computer, given detailed information about positions, momenta, and forces at work on every particle in universe, could predict future of universe.

Such thinking —> doubts about free will, human mind, morality, etc…

Heisenberg's attack simple.

Uncertainty principle prevents knowing both position and momentum of any particle with sufficient accuracy, ultimately rendering predictions impossible.

Causality or notion that everything that happens is **caused by something happening previously** had been central to physics for hundreds of years.

Experiments in quantum world undermined this thinking with unpredictable outcomes and then came uncertainty principle undermining basis of all physics, namely, ability to measure with infinite accuracy.

—> nature seemed to be built on a set of principles completely contrary to approach science had been using for generations.

It is possible to regard quantum challenge as being **in** microworld and of **no** relevance to macroworld in which we live.

However, study of chaotic systems has shown that small effects do not necessarily lead to small consequences and that **no** measurement (quantum or otherwise) can, even in principle, yield sufficiently accurate results to do job.

Putting aside the philosophical debate, the uncertainty principle is **strange**.

Basic science education —> vital role of measurement and precision.

Measurement has to be a faithful rendering of what really out there, or no point in exercise.

Realism is natural state of physicist.

When run into quantum physics —> begin to doubt instincts - surely it cannot really be like that!

Some say — I'm Not Sure What Uncertainty Means...

Serious point here.

Much is written about meaning and significance of uncertainty principle.

All depends on what side of realist/instrumentalist divide one stands and on exactly what type of interpretation one favors.

Most important aspect of uncertainty principle, from a philosophical point of view, is

whether to take it to be limitation on what we can know about world

or as an inherent limitation on the world itself.

In philosophical speak, is it an epistemological(potential) statement, or is it ontological(actual)?

Whatever Heisenberg thought: first arguments used to demonstrate uncertainty principle had epistemological tinge —> famous gamma ray microscope.

Heisenberg envisaged microscope constructed to use gamma rays rather than light waves.

Microscope could be used to determine position of something as small as electron by detecting gamma rays that scattered off electron.

Path of gamma rays pinpoints location of electron.

Snag is wavelength of gamma rays.

Well known in optics —> limitation on detail that can be seen by microscope.

Limitation depends on wavelength of electromagnetic waves used.

To view level of detail sufficient to locate something as small as electron, quite high-energy gamma rays (very short wavelength) are needed.

Such gamma ray scattering off an electron would transfer a significant amount of energy

causing electron to recoil in an unspecified direction, affecting its momentum.

No way of determining energy transferred and thus, subsequent disturbance to electron's momentum.

If wanted to do that, would have to measure the gamma ray before it hit and that process would then disturb gamma ray.

Key point:

Gamma ray microscope argument assumes electron has position **and** momentum;

Problem: interaction with measuring device —> cannot accurately determine these quantities.

Every time measuring device makes measurement, it interacts with thing it is measuring.

In classical physics, scale of interactions either too small to worry us or can be calculated and accounted for in sensible way.

This cannot be said in quantum physics.

Scale of interaction too large and causes major changes to other properties of object being measured.

Also interaction is inherently unpredictable.

Any attempt to stick another measuring device in to judge scale of interaction will not work as second instrument interacts in manner that cannot be detected.

Infinite regression follows, which like most infinite regressions gets us nowhere.

An important argument which cuts once again to nature of quantum state.

If electron happens to be in an eigenstate of position (or momentum) can say with certainty what result of measuring that position (or momentum) will be.

If electron is not in eigenstate then what can we say?

In QM, one can predict various values to be revealed by measurement, and relative probabilities, but what does that tell us?

What is position (or momentum) of electron in this situation?

If take view that states refer to collections of objects, rather than individual objects, then way forward quite clear —> use statistics!

Inside collection, electrons identical, within limitations laid down by nature of quantum state.

However, could be hidden variables - do not know how to measure - which take different values in collection.

When measurement made, variables determine outcome, but because do not know values, all looks random.

Electrons have perfectly well-determined positions and momenta, but cannot know in advance what they will be since cannot access physics that determines values.

Gamma ray microscope example runs contrary to this way of thinking.

Argument is clearly dealing with single electrons, not collections.

However, does assume that positions and momenta are really there, just partially hidden.

Such hidden variable type view —> uncertainty principle expressing epistemological limits of current science.

Heisenberg himself took view that uncertainty principle was an ontological thing.

To him uncertainty principle expressed limits to which classical ideas of momentum, or position, can be applied in given situation.

Harks back to earlier argument about how classical definition of momentum cannot be applied in quantum physics.

If take this view, then have to say that object that is not in eigenstate of position simply **does not have a position**.

Concept has no meaning in such a state.

Heisenberg would have said that position of particle was a **latent or potential property**.

See supplementary readings — Taking Heisenberg's Potentia Seriously - Epperson

When measurement occurs, latency becomes actuality and classical ways of thinking can now be applied.

—> realistic view, but acknowledges that assumptions about what should be classified as real, may have to be enlarged.

To Heisenberg latent properties should be thought of as real.

Gamma ray microscope was used by Heisenberg as way of introducing uncertainty principle.

However, he held back from stating that was an ontologically valid description of what would actually happen.

Yet More Uncertainty

Have tried to show how uncertainty principle arises as natural consequence of ability to expand state in various bases.

Although illustrated with position and momentum, are many other physical variables with basis sets that can be used to expand a state.

Unless position and momentum have special significance, must be versions of uncertainty principle that apply to other physical variables also.

Basic rule:

if choose to expand state over pair of bases,

and physical variables involved

happen to be **"conjugate" or "incompatible"** variables,

then going to end up with an uncertainty principle.

State cannot be an eigenstate of two conjugate variables at same time

—> cannot get a definite value from measuring either variable,

and then uncertainty relationship must apply.

Non-Compatibility

If operator acts on its eigenstate, result is that state remains same,

but now multiplied by eigenvalue.

$$\hat{p}\,|p_1\rangle = p_1\,|p_1\rangle \quad , \quad \hat{x}\,|x_1\rangle = x_1\,|x_1\rangle$$

If state not an eigenstate, then strange things happen.

Consider two physical variables represented by operators $\hat{O}_1$ and $\hat{O}_2$,

which happen to be **compatible ==> simultaneously measurable**.

Then state $|\phi\rangle$ can be an eigenstate of both.

What happens when apply both operators on this state, one **after** other?

$$\hat{O}_1 |\phi\rangle = o_1 |\phi\rangle$$
$$\hat{O}_2 o_1 |\phi\rangle = o_1 \hat{O}_2 |\phi\rangle = o_1 o_2 |\phi\rangle$$

In addition, could also do things other way around

$$\hat{O}_2 |\phi\rangle = o_2 |\phi\rangle$$
$$\hat{O}_1 o_2 |\phi\rangle = o_2 \hat{O}_1 |\phi\rangle = o_2 o_1 |\phi\rangle$$

which says that

$$\hat{O}_2 \hat{O}_1 |\phi\rangle = \hat{O}_1 \hat{O}_2 |\phi\rangle$$

or

$$\left[\hat{O}_2 \hat{O}_1 - \hat{O}_1 \hat{O}_2\right] |\phi\rangle = 0$$

Bracket is regarded as being so important in quantum theory that given own name:

THE COMMUTATOR

$$Commutator\left(\hat{O}_2, \hat{O}_1\right) = \left[\hat{O}_2, \hat{O}_1\right] = \left[\hat{O}_2\hat{O}_1 - \hat{O}_1\hat{O}_2\right]$$

If two operator commute, then $\left[\hat{O}_2, \hat{O}_1\right] = 0$, and **can** (not must) have simultaneous eigenstates.

If don't commute, then simultaneous eigenstates not possible and uncertainty relationship follows.

RULE:

Non-compatible physical variables represented by operators that do not commute and so are linked by uncertainty principle.

$$\Delta O_1 \Delta O_2 \geq \frac{1}{2}\left\langle i\left[\hat{O}_2, \hat{O}_1\right]\right\rangle$$ **Derive later**

Consider Time Again....

Exists uncertainty relationship that does not follow directly from these arguments, namely,

ENERGY/TIME UNCERTAINTY $\Delta E \Delta\tau \geq \frac{\hbar}{2}$ **Derive later**

where τ related to characteristic development time of system, i.e., given by $\tau = \frac{\langle A\rangle}{d\langle A\rangle/dt}$ for any physical variable of system.

Let us see how this works.

Rule 7 states a link between operators and measurement of physical variables.

Have seen several examples of such operators and learned to calculate expectation values.

Where things start to go weird is over a possible $\hat{t}$ operator.

Can measure position, momentum, energy, etc, but how do you measure time?

Let us be clear about this: we can measure duration (e.g., with stopwatch) but what physical measurement tells you actual time?

Issues involved are very subtle, and not every quantum physicist would agree with what I am about to say, but here goes.

Conventional quantum mechanics is acted out on stage formed from three directions in space and one universal time (x,y,z,t).

Think of space arena as giant spider's web; with threads so closely woven that web appears to be continuous disk.

Spider able to crawl around and occupies position at any moment on web.

Position of spider is physical variable -> property of spider's state that can be measured. In quantum mechanics, position represented by operator $\hat{x}$.

In some accounts, spider position operator denoted as $\hat{q}$ to make clear **not same** as the x-value, which is part of web, not part of spider's state.

Time rather different than this.

If can picture it at all, think of it as line stretching out to infinity in both directions.

Spider on line would not be stationary, but sliding along line as time passes.

In most instances cannot conceive of experiment that would enable us to to measure spider's position on time line.

—-> time is not property of spider's state; **hence there is no time operator**.

But what about a system that is changing with time - a falling ball, for example?

Here the momentum of ball increases as time passes, suggesting that it is "time variable".

However, a physical variable that is changing does not necessarily change at a uniform rate (momentum increases at increasing rate),

and when it hits the ground, time as measured by momentum would stop,

although t continues.

There are some systems in which a physical variable corresponds in a more satisfactory way with the passage of time —> clocks.

The second hand of analog clock has angular position θ, which changes with time.

Measuring the angle gives a measure of time for system.

In a real clock there will also be an energy, which will tend to depend on θ so that we can define an energy operator $\hat{E}$ and a time operator $\hat{\theta}$, which will not commute and so lead to uncertainty relationship.

However, θ is only a physical variable of a clock and cannot be applied as time operator to any other system.

Think about Energy/Time Uncertainty another way……

Instead, ask this question: how can we tell if something has changed?

A physical system will have various properties that can be measured.

If suspect something changed system state, can go through new set of measurements to see what's going on.

How to do it in quantum physics?

Every measurement made will “collapse state” into an eigenstate, so can’t necessarily repeat measurements on same state.

If do lots of measurements on collection of identical systems in same state,

how do we tell if values are different

because of some change affecting all systems in collection,

or is just normal spread of measurements

that comes with state not an eigenstate to begin with?

Suppose we are trying to distinguish between range of values caused by inherent

uncertainty in state itself and range caused by some

causal evolution ($\hat{U}(t)$) taking place in systems.

Way to tell is to compare expectation value $\langle O \rangle$ with uncertainty ΔO.

If remeasure collection of systems

—> difference between new expectation value $\langle O' \rangle$ and old greater than ΔO

(i.e., $(\langle O' \rangle - \langle O \rangle) \geq \Delta O$),

then can be sure that system state has actually changed.

Then becomes question of how long have to leave system to evolve, in given situation,

before change can be detected in this manner.

Can work it out if employ rate of change with time (a derivative)operator, d/dt (a derivative)

In elementary physics, distance traveled by moving object is rate at which traveling (speed) multiplied by time travels for ($d = s \times t$).

Equally well , if know rate at which any quantity changing and multiply by time period, that will produce amount by which quantity has changed in that time.

If quantity happens to be an expectation value then $d\langle O \rangle/dt\ \Delta t$ is amount by which expectation value has changed in time duration Δt.

The condition for a measurable change then becomes

$$\frac{d}{dt}\langle O\rangle \Delta t \geq \Delta O \quad \text{or} \quad \Delta t \geq \frac{\Delta O}{\frac{d}{dt}\langle O\rangle}$$


—> how we should interpret Δt in energy-time uncertainty relationship.

It represents duration of time that we have to wait to be sure that energy has changed by amount at least equal to ΔE.

The Meaning of the Indeterminacy Relations

What is significance of indeterminacy relations in the world of experimental physics?

Consider experimental results shown below:



—> frequency distributions for results of independent measurements of Q and P on ensemble of similarly prepared systems,

i.e., on each of large number of similarly prepared systems one performs a single measurement (either Q **or** P).

Histograms are statistical distribution of results.

Standard deviations (variances) shown must satisfy (according to theory) relation

$$\Delta_Q \Delta_P \geq \frac{\hbar}{2}$$


They must be distinguished from resolution of individual measurements, δQ and δP.
Let me emphasize these points:

(1) Quantities Δ_Q and Δ_P are not errors of measurement.

Errors or preferably the resolutions of the Q and P measuring instruments are δQ and δP .

They are logically unrelated to Δ_Q and Δ_P and to uncertainty relations except for practical requirement that if

$$\delta Q > \Delta_Q \quad (\text{ or } \delta P > \Delta_P)$$



then will not be possible to determine Δ_Q (or Δ_P) in an experiment and experiment cannot test any uncertainty relation.

(2) An experimental test of indeterminacy relation does not involve simultaneous measurements of Q and P, but involves measurement of one or other of these dynamical variables on each independently prepared representative of particular state being studied.

Why am I being so picky here?

Quantities Δ_Q and Δ_P are often misinterpreted as errors of individual measurements.

Probably arises because Heisenberg's original paper on this subject, published in 1927, was based on early version of quantum mechanics that predates systematic formulation and statistical interpretation of quantum mechanics as it exists now.

Derivation carried out in this lecture now was not possible in 1927!

Derivation of the Uncertainty Relations in General

Very mathematical - just show how it can be done……

Given two Hermitian operators $\hat{A}$ and $\hat{B}$ and state vector $|\psi\rangle$, **define** two new operators

$$\hat{D}_A = \hat{A} - \left\langle \hat{A} \right\rangle \quad , \quad \hat{D}_B = \hat{B} - \left\langle \hat{B} \right\rangle$$

where $\langle ... \rangle$ equals average or expectation value in a state $|\psi\rangle$.

In statistical analysis of data, one uses quantity called standard or mean-square deviation as measure of uncertainty of observed quantity.

This is defined, for set of N measurements of quantity q by

$$(\Delta q)^2 = (\text{standard deviation})^2 = \frac{1}{N}\sum_{i=1}^{N}(q_i - q_{average})^2$$

$$= \frac{1}{N}\sum_{i=1}^{N}(q_i)^2 - \frac{1}{N}\sum_{i=1}^{N}(q_i q_{average}) - \frac{1}{N}\sum_{i=1}^{N}(q_{average} q_i) + \frac{1}{N}\sum_{i=1}^{N}(q_{average})^2$$

$$= (q^2)_{average} - (q_{average})^2$$

where have used

$$q_{average} = \frac{1}{N}\sum_{i=1}^{N} q_i$$

In analogy, define mean-square deviations for $\hat{A}$ and $\hat{B}$ as

$$(\Delta\hat{A})^2 = \left\langle \hat{A}^2 \right\rangle - \left\langle \hat{A} \right\rangle^2 = \left\langle \left(\hat{A} - \left\langle \hat{A} \right\rangle \right)^2 \right\rangle = \left\langle \hat{D}_A^2 \right\rangle$$
$$(\Delta\hat{B})^2 = \left\langle \hat{B}^2 \right\rangle - \left\langle \hat{B} \right\rangle^2 = \left\langle \left(\hat{B} - \left\langle \hat{B} \right\rangle \right)^2 \right\rangle = \left\langle \hat{D}_B^2 \right\rangle$$

Then we have

$$(\Delta\hat{A})^2 (\Delta\hat{B})^2 = \left\langle \hat{D}_A^2 \right\rangle \left\langle \hat{D}_B^2 \right\rangle$$

Now we assume that

$$\left[\hat{D}_A, \hat{B}\right] = \left[\hat{A}, \hat{B}\right] = \left[\hat{D}_A, \hat{D}_B\right] = i\hat{C}$$

they do not commute

where $\hat{C}$ is also Hermitian operator and let

$$|\alpha\rangle = \hat{D}_A |\psi\rangle = \left(\hat{A} - \left\langle \hat{A} \right\rangle \right) |\psi\rangle \quad , \quad |\beta\rangle = \hat{D}_B |\psi\rangle = \left(\hat{B} - \left\langle \hat{B} \right\rangle \right) |\psi\rangle$$

Then we have

$$\left(\Delta\hat{A}\right)^2 = \left\langle \hat{D}_A^2 \right\rangle = \langle\psi|\, \hat{D}_A^2\, |\psi\rangle = \left(\langle\psi|\, \hat{D}_A\right)\left(\hat{D}_A\, |\psi\rangle\right) = \langle\alpha \mid \alpha\rangle$$
$$\left(\Delta\hat{B}\right)^2 = \left\langle \hat{D}_B^2 \right\rangle = \langle\psi|\, \hat{D}_B^2\, |\psi\rangle = \left(\langle\psi|\, \hat{D}_B\right)\left(\hat{D}_B\, |\psi\rangle\right) = \langle\beta \mid \beta\rangle$$

Schwarz inequality says that for any two vectors we must have the relation

$$\langle\alpha \mid \alpha\rangle\, \langle\beta \mid \beta\rangle \geq |\langle\alpha \mid \beta\rangle|^2$$

Therefore we have

$$\left(\Delta\hat{A}\right)^2 \left(\Delta\hat{B}\right)^2 = \left\langle \hat{D}_A^2 \right\rangle \left\langle \hat{D}_B^2 \right\rangle = \langle\alpha \mid \alpha\rangle\, \langle\beta \mid \beta\rangle \geq |\langle\alpha \mid \beta\rangle|^2 \; = \left|\langle\psi|\, \hat{D}_A \hat{D}_B\, |\psi\rangle\right|^2 = \left|\left\langle \hat{D}_A \hat{D}_B \right\rangle\right|^2$$

which gives an uncertainty relation of the form

$$\left(\Delta\hat{A}\right)^2 \left(\Delta\hat{B}\right)^2 \geq \left|\left\langle \hat{D}_A \hat{D}_B \right\rangle\right|^2$$

Now

$$\left|\langle \hat{D}_A \hat{D}_B \rangle\right|^2 = \left|\langle \frac{1}{2}[\hat{D}_A, \hat{D}_B] + \frac{1}{2}\{\hat{D}_A, \hat{D}_B\}\rangle\right|^2$$

$$[\hat{D}_A, \hat{D}_B] = [\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}$$

$$\{\hat{D}_A, \hat{D}_B\} = \{\hat{A}, \hat{B}\} = \hat{A}\hat{B} + \hat{B}\hat{A}$$

$$\left|\langle \hat{D}_A \hat{D}_B \rangle\right|^2 = \left|\langle \frac{1}{2}[\hat{A}, \hat{B}] + \frac{1}{2}\{\hat{A}, \hat{B}\}\rangle\right|^2$$

and

$$\left[\hat{A}, \hat{B}\right]^\dagger = -\left[\hat{A}, \hat{B}\right] \rightarrow anti - hermitian \rightarrow expectation\ value\ is\ imaginary$$

commutator

and

$$\left\{\hat{A}, \hat{B}\right\}^\dagger = \left\{\hat{A}, \hat{B}\right\} \rightarrow hermitian \rightarrow expectation\ value\ is\ real$$

anticommutator

Therefore,

$$\left(\Delta\hat{A}\right)^2 \left(\Delta\hat{B}\right)^2 \geq \left|\left\langle \tfrac{1}{2}\left[\hat{A}, \hat{B}\right] + \tfrac{1}{2}\left\{\hat{A}, \hat{B}\right\}\right\rangle\right|^2$$
$$\left(\Delta\hat{A}\right)^2 \left(\Delta\hat{B}\right)^2 \geq \tfrac{1}{4}\left|\left\langle i\hat{C}\right\rangle + a\right|^2$$

where a = real number.

We then have

$$\left(\Delta\hat{A}\right)^2\left(\Delta\hat{B}\right)^2 \geq \tfrac{1}{4}\left|\left\langle i\hat{C}\right\rangle + a\right|^2 = \tfrac{1}{4}|a|^2 + \tfrac{1}{4}\left|\left\langle \hat{C}\right\rangle\right|^2$$
$$\left(\Delta\hat{A}\right)^2\left(\Delta\hat{B}\right)^2 \geq \tfrac{1}{4}\left|\left\langle \hat{C}\right\rangle\right|^2$$

since $|a|^2/4 \geq 0$.

If $\left[\hat{A},\hat{B}\right] = i\hat{C} = 0$ then get

$$\left(\Delta\hat{A}\right)^2\left(\Delta\hat{B}\right)^2 \geq 0$$

and have no uncertainty relation between observables. They are **compatible**!

On other hand, if $\hat{A} = \hat{x}$ and $\hat{B} = \hat{p}_x$, have

$$[\hat{x},\hat{p}_x] = i\hat{C} = i\hbar\hat{I}$$

which means

$$\left(\Delta\hat{A}\right)^2\left(\Delta\hat{B}\right)^2 \geq \frac{\hbar^2}{4} \qquad \text{or} \qquad \Delta\hat{x}\Delta\hat{p}_x \geq \frac{\hbar}{2}$$

which is Heisenberg uncertainty principle. **It is simply the Schwarz inequality!!** which every sophomore math student understands! It is in QM because we assumed QM is governed by Linear Algebra!

Hopefully, that removes any mysticism associated with the principle!

Finally we derive the Time-Energy Uncertainty Relations

Use of time-energy uncertainty relations in most textbooks is simply incorrect.

Now show (very mathematical) the most one can say about such relations.

Need to derive time-dependence of expectation values. We have

$$\left\langle \hat{Q} \right\rangle = \langle \psi(t) | \, \hat{Q} \, | \psi(t) \rangle \qquad\qquad |\psi(t)\rangle = e^{-i\hat{H}t/\hbar} \, |\psi(0)\rangle \rightarrow \langle \psi(t)| = \langle \psi(0)| \, e^{-i\hat{H}t/\hbar}$$

so that

$$\frac{d\left\langle \hat{Q} \right\rangle}{dt} = \frac{d}{dt} \langle \psi(t) | \, \hat{Q} \, | \psi(t) \rangle = \frac{d}{dt} \langle \psi(0) | \, e^{i\hat{H}t/\hbar} \hat{Q} e^{-i\hat{H}t/\hbar} \, | \psi(0) \rangle$$

$$= \langle \psi(0) | \, \frac{d}{dt} \left(e^{i\hat{H}t/\hbar} \hat{Q} e^{-i\hat{H}t/\hbar} \right) | \psi(0) \rangle$$

$$= \langle \psi(0) | \left(\frac{i\hat{H}}{\hbar} e^{i\hat{H}t/\hbar} \hat{Q} e^{-i\hat{H}t/\hbar} \right) | \psi(0) \rangle - \langle \psi(0) | \left(e^{i\hat{H}t/\hbar} \hat{Q} \frac{i\hat{H}}{\hbar} e^{-i\hat{H}t/\hbar} \right) | \psi(0) \rangle$$

$$= \langle \psi(0) | \, e^{i\hat{H}t/\hbar} \frac{i}{\hbar} \left[\hat{H}, \hat{Q} \right] e^{-i\hat{H}t/\hbar} \, | \psi(0) \rangle = \frac{i}{\hbar} \langle \psi(t) | \left[\hat{H}, \hat{Q} \right] | \psi(t) \rangle$$

or

$$\frac{d\left\langle \hat{Q} \right\rangle}{dt} = \frac{1}{i\hbar} \left\langle \left[\hat{Q}, \hat{H} \right] \right\rangle$$

Consider dynamical state of system at time t.

Let |ψ⟩ be vector representing that state. Call ΔQ, ΔE root-mean-square deviations of $\hat{Q}$ and $\hat{H}$ respectively. Applying Schwarz inequality (as above) to vectors

$$(\hat{Q} - \langle Q \rangle)\,|\psi\rangle \quad , \quad (\hat{H} - \langle H \rangle)\,|\psi\rangle$$

and carrying out same manipulations as (as earlier), find after some calculations

$$\Delta Q \Delta E \geq \frac{1}{2}\left|\left\langle \left[\hat{Q}, \hat{H}\right] \right\rangle\right|$$

with equality being realized when |ψ⟩ satisfies the equation

$$(\hat{Q} - \alpha)\,|\psi\rangle = i\gamma(\hat{H} - \varepsilon)\,|\psi\rangle$$

where α, γ and ε are arbitrary real constants (as earlier).

Putting everything together we get

$$\frac{\Delta Q}{\left|\frac{d\langle \hat{Q} \rangle}{dt}\right|}\Delta E \geq \frac{\hbar}{2} \qquad \text{or} \qquad \tau_Q \Delta E \geq \frac{\hbar}{2}$$

where we have defined

$$\tau_Q = \frac{\Delta Q}{\left|\frac{d\langle \hat{Q} \rangle}{dt}\right|}$$

τ_Q appears as the time characteristic of evolution of expectation value of $\hat{Q}$.

It is time required for center $\langle \hat{Q} \rangle$ of statistical distribution of $\hat{Q}$ to be displaced by amount equal to width ΔQ.

In other words, the time necessary for this statistical distribution to be appreciably modified.

In this way can define characteristic evolution time for each dynamical variable of system.

Let τ be shortest of times thus defined.

τ may be considered as characteristic time of evolution of system itself,

that is, whatever measurement carried out on system at an instant of time t',

the statistical distribution of results is essentially same as would be obtained at instant t,

as long as difference $|t-t'|$ is less than τ.

According to derivation, the time τ and energy spread ΔE satisfy time-energy uncertainty relation

$$\tau \Delta E \geq \frac{\hbar}{2}$$

If system in stationary state where $\frac{d\left\langle \hat{Q} \right\rangle}{dt} = 0$ no matter what $\hat{Q}$, and consequently τ

is infinite, then ΔE = 0. That is meaning of a stationary state.

Ordinary time t is just a parameter in non-relativistic QM and not an operator!

Our derivation **does not** say that

$$\Delta t \Delta E \geq \frac{\hbar}{2}$$

which is an equation that has no meaning!

Now on to EPR and Bell

Entanglement, EPR and Bell

Two threads are running through this part of lecture.

1st stems from Einstein's desire

to discredit quantum physics as fundamental theory of reality

which led to the most impressive and far-reaching confirmations of the quantum view.

2nd derives from Schrodinger desire to destroy QM and

who introduced term entanglement to physics community and showed how

entangled states lead to very bizarre features of quantum world.

Multiple passes through each topic

—> see different derivations, discussions and points of view.

Hopefully this enhances understanding!

Is Quantum theory complete?

Despite being responsible for number of key developments in theory,

Einstein was never happy with final form of QM.

1st he objected to uncertainty principle

—> developed thought experiments and arguments to get around it.

Bohr and Heisenberg made counter arguments that saved the uncertainty principle and QM many times.

Many of Einstein's proposed mechanical devices (some shown on next slide!!!!)






FIG. 6





After losing earlier arguments, Einstein then switched from trying to show QM was inconsistent

(gets different answers in similar cases)

to instead proving QM is incomplete (exist situations not covered by theory).

In 1935 Einstein thought he found way to discredit QM.

With Boris Podolsky and Nathan Rosen (EPR paper),

he constructed an argument that he thought

demonstrated the need for a deeper(more deterministic) theory to replace QM.

Paper titled

‘’Can a quantum mechanical description of physical reality be considered complete?”

Paper first defines a complete physical theory:

If theory is complete,

then every element of physical reality

must have counterpart in physical theory.

Reasonable, but we need to know what is meant by **element of physical reality**?

It is not straightforward.

EPR —> cannot guess or try to figure out elements of physical reality.

They have to be found from the results of experiments and measurements.

World is too subtle and surprising to guess correctly.

Need experiments as guide.

Still not simple.

Experiments —> huge amounts of data —> useful info + random noise.

Need to separate key features of information from random (experimental) fluctuations.

EPR directly tied theory with experiment and said following condition must be met:

if, without in any way disturbing the system, we can predict

with certainty the value of a physical quantity

then there exists an element of physical reality

corresponding to this physical quantity

We note that the part about **"disturbing the system"** —> clever **trap** for quantum theory.

Then EPR apply their condition for reality to a specific case.

Quantum theory employs state vector $|\psi\rangle$

—> systems and operators

—> physical variables $\hat{O}$.

If |ψ⟩ happens to be eigenstate of $\hat{O}$, then

$$\hat{O}\,|\psi\rangle = a\,|\psi\rangle$$ a = corresponding eigenvalue

Then can say with certainty that measurement of physical variable linked to

$\hat{O}$ —> a (one of our postulates).

Then, EPR condition —> physical variable

represented by $\hat{O}$ = element of reality (not clear if in general or only in that state?).

However, if one chooses to measure another physical variable

linked to operator $\hat{Q}$ and |ψ⟩ is not an eigenstate,

then cannot predict with certainty what value measurement will produce.

Expand |ψ⟩ over basis of eigenstates of $\hat{Q}$

and calculate expectation value for set of results,

or probability of given result.

Only way to predict with 100% certainty what measurement will give

is to disturb system by turning |ψ⟩ into one of eigenstates of $\hat{Q}$.

Thus, EPR condition —> physical variable represented by

$\hat{Q}$ cannot be element of physical reality, or, at least, not at same time as $\hat{O}$.

Specifically, electron in state |p⟩,

eigenstate of momentum operator, $\hat{p}$

—> momentum is part of electron's reality.

However, in this state, position cannot be element of electron's reality

as |p⟩ not eigenstate of $\hat{x}$.

Basis of EPR argument is the philosophical view

that one cannot create reality

by bending the electron to fit words "disturb the system".

Should be able find set of physical variables and theory

-> predict them with certainty,

as reality is out there to be found,

not something created.

Having set up conditions for reality

and illustrated them for simple quantum mechanical case,

EPR spring trap by describing a situation

where it appears quantum theory

fails to describe every element of a system's reality.

E(instein)P(odolsky)R(osen) Paradox(?) Details

Challenge to standard (Copenhagen) view of QM

—> surprising aftermath - resulted in1960s (25 years later!) work of John Bell.

1st discuss EPR and then fundamental work of Bell.

In 1935, EPR produced an argument

—> escape from the standard way of thinking about QM.

1st, they define completeness:

> **A description of the world is complete (for EPR) if nothing that is true about the world, nothing that is an element of the reality of the world is left out of the description.**

EPR **never** actually presented a prescription

for determining what all elements of reality are (**made any challenge difficult**).

Instead, they did something much narrower

(which was **sufficient** for the purposes of their argument).

They wrote down a condition for a measurable property of system at any instant

to be element of reality of system at that instant.

Condition is that:

> **if, without in any way disturbing a system, we can predict with certainty (with probability = 1) the value of a physical quantity, then there exists an element of reality corresponding to this physical quantity.**

What does this condition mean?

Consider following question:

If measurement of particular observable $\hat{O}$ of certain physical system S

were carried out at certain future time T,

what would outcome be?

Suppose a method is available so that,

prior to time T,

we can answer question with certainty.

Also, suppose method used —> no disturbance of system S.

Then(via EPR) there must be definite information(**hidden somehow**)

about outcome of a future $\hat{O}$ measurement on S at T.

For instance, suppose we just measured color of electron.

Since measurements of color are repeatable,

can predict with certainty what the outcome of later color measurement will be,

if measurement carried out.

Making prediction need not involve any further interaction with electron.

So EPR reality condition says color must, **at that moment**,

be element of reality of this electron.

—> Identical to standard way of thinking in QM! Just —> Color talk now applicable

Suppose, instead we measured hardness of electron.

To predict with certainty

what the outcome of a future measurement of color of electron would be

(if made measurement),

need to measure color of electron (interact and potentially disturb state).

EPR reality condition does not say color of electron

is element of reality of electron at that moment.

Again agrees completely with standard way of thinking! Color talk not applicable!

EPR then want to argue the following:

If predictions of QM correct,

then there must exist elements of reality of world

which have no corresponding elements

in Quantum Mechanical description of world

They are attempting to use QM against itself.

They are saying QM is missing something! It is not complete!

Their Argument: (presented in the hardness-color world by me for our benefit)

Consider system consisting of two electrons.

Electron 1 located at position 1,

and electron 2 located at position 2.

Assume that color-space state of two electrons is (was created somehow)

(note —> is a nonseparable or entangled color state - like two gloves in story)

$$|A\rangle = \frac{1}{\sqrt{2}} |green\rangle_1 |magenta\rangle_2 - \frac{1}{\sqrt{2}} |magenta\rangle_1 |green\rangle_2$$

State |A⟩ is necessarily eigenstate of some observable(always true for all states)

(a Hermitian operator) of this pair of electrons, say $\hat{O}$,

where $\hat{O}$ |A⟩ = +1|A⟩(eigenvalue = +1).

We happen to have written |A⟩ in color basis.

Now convert to hardness basis. Remember

$$|green\rangle = \frac{1}{\sqrt{2}} |hard\rangle + \frac{1}{\sqrt{2}} |soft\rangle$$
$$|magenta\rangle = \frac{1}{\sqrt{2}} |hard\rangle - \frac{1}{\sqrt{2}} |soft\rangle$$

Substituting get

$$|A\rangle = \frac{1}{\sqrt{2}}|soft\rangle_1|hard\rangle_2 - \frac{1}{\sqrt{2}}|hard\rangle_1|soft\rangle_2$$

Note it has the exact **same** nonseparable form in both bases.

In fact, it takes **same** nonseparable form in any basis! **—> a maximal entangled state**

Collapse postulate for two-particle entangled states (we develop needed extra theory)

If in state of type |A⟩, then the following rules apply during measurements.

1. If, in state |A⟩, we only measure particle 1, $|A\rangle = \frac{1}{\sqrt{2}}|green\rangle_1|magenta\rangle_2 - \frac{1}{\sqrt{2}}|magenta\rangle_1|green\rangle_2$
then if particle 1 is measured to be green,
state collapses to $|green\rangle_1|magenta\rangle_2$

i.e., at same time we know 1 is green we **also** know 2 is magenta,

WITHOUT actually measuring 2. **That is what entangled states do!!**

Similar result if measured 1 is magenta —> state would be $|magenta\rangle_1|green\rangle_2$

i.e., at the same time that we know 1 is magenta we **also** know 2 is green,

WITHOUT actually measuring 2.

2. If, in state $|A\rangle$, the probability of measuring 1 is green is still 1/2.

This is now shown below.

Probability for measurement of some property of particle 1

in a **2-particle entangled state** is calculated as follows:

(a) Step 1:

Since we not measuring particle 2,
must calculate probability for measuring 1
for all possible states of particle 2.

In this case

$$prob(1 \text{ is green}) = prob(1 \text{ is green AND } 2 \text{ is magenta}) + prob(1 \text{ is green AND } 2 \text{ is green})$$

(b) Probability (x AND y) is calculated as shown below:

$$prob(1 \text{ is green AND } 2 \text{ is magenta}) = |\langle green|_1 \langle magenta|_2 |A\rangle|^2$$

Carry out measurement of color of electron 1 (state in color basis).

Outcome of measurement either green or magenta with equal probability

(using two-particle state probability rules).

As stated above, in state A

$$prob(1\,is\,green) = prob(1\,is\,green\,and\,2\,is\,magenta) + prob(1\,is\,green\,and\,2\,is\,green)$$

since not looking at 2.

Now

$$prob(1\,is\,green\,and\,2\,is\,magenta) = |\langle green|_1 \langle magenta|_2 |A\rangle|^2 = 1/2$$

$$prob(1\,is\,green\,and\,2\,is\,green) = |\langle green|_1 \langle green|_2 |A\rangle|^2 = 0$$

Total probability to be green = 1/2. Similarly for being magenta!

Moreover, QM says(this is experimentally confirmed)

that in event outcome of measurement green,

then outcome of any subsequent measurement of color of electron 2

will necessarily be magenta

and in event outcome of measurement magenta,

then outcome of any subsequent measurement of color of electron 2

will necessarily be green.

—> a **2-particle "collapse"** occurred!

Both statements follow directly from collapse postulate for two-particle states.

Remember the 2-path experiment for color/hardness where similar things happened.

EPR assumed

(the only assumption (called **locality**) they make

on top of basic assumption that predictions of QM are correct)

that things could in principle be set up in such a way

as to guarantee that measurement of color of electron 1

produces no physical disturbance whatsoever in electron 2.

To them —> self-evident statement!

What influence could there be???

Any number of ways to satisfy this condition.

Could separate two electrons by immense distance

(no assumption says any properties of QM changes with electron separation).

Then two measurement events,

according to special relativity, could not influence each other

since c = max info speed(called spacelike separated).

Or could insert impenetrable wall between them

(nothing assumed says any properties of QM

depends on what located between electrons).

Or could set up array of detectors to verify that

no measurable signals pass from one electron to other during experiment

(QM predicts array will never register anything).

Very important point.

Locality assumption says

I cannot punch you in the nose unless my fist gets to the place where your nose is (in space and time)

Of course, something I do with my fist far from where your nose is

can cause some other fist which is near your nose to punch your nose

(i.e., something I do with my fist might signal somebody else to punch you in the nose).

Their seemingly obvious assumption is just this:

if my fist never gets anywhere near your nose then I cannot punch you in the nose directly. If you got punched in such an arrangement, then it cannot be my fist that punched you

If something I do with my fist far from your nose is cause of your getting punched in nose,

then necessarily some causal sequence of events at contiguous points in space and at contiguous points in time (propagation of some signal for instance) stretches all the way without break from whatever it was I did with my fist to your being punched in the nose.

Important thing about sequence of events

is it necessarily requires some finite time

(exact amount depends on path in space and time)

to completely unfold.

Shortest time possible would occur

if signal(s) travel with speed of light in straight line

(maximum speed of propagation of information).

So summarizing their assumption:

measurement on color 1 has no effect on measurement of color 2

if measurements are spacelike separated(cannot be connected by light)

Returning to color basis entangled state

$$|A\rangle = \frac{1}{\sqrt{2}} |green\rangle_1 |magenta\rangle_2 - \frac{1}{\sqrt{2}} |magenta\rangle_1 |green\rangle_2$$

It is clear that one can predict with certainty,

if locality true,

without disturbing electron 2,

what outcome of subsequent measurement of color 2 will be.

Measure color 1

→ know outcome of measurement of color 2

is opposite of outcome of measurement of color 1

or know color 2 without measuring it!

Reality condition then says that color is element of reality of electron 2.

i.e., color of 2 has definite value when in state |A⟩.

So both 1 and 2 have definite color values in this state!

Switch to hardness basis —> talk about hardness measurements.

$$|A\rangle = \frac{1}{\sqrt{2}} |soft\rangle_1 |hard\rangle_2 - \frac{1}{\sqrt{2}} |hard\rangle_1 |soft\rangle_2$$

Using same arguments in hardness basis

find that both 1 and 2 have definite hardness values!

Since can actually prepare states like |A⟩ in real world,

—> EPR say standard interpretation must be false. WHY?

EPR concluded both color and hardness are elements of reality of electron 2,

even though the two observables are incompatible according to QM.

Thus, formalism must be incomplete,

since some elements of physical reality of world

have no corresponding elements in QM.

Must exist hidden facts(not in postulates)

about color and hardness of 2, when in state $|A\rangle$.

Is there way out of EPR's proposed dilemma?

Nothing in QM formalism allows both color/hardness of electron 2

to be predicted with certainty simultaneously.

Similar arguments hold for electron 1.

EPR were clearly very clever!

If true, statement that system in state $|A\rangle$

constitutes incomplete description of state of pair of electrons

—> are some hidden variables.

EPR say QM predicts everything correctly, **but is still wrong.**

EPR noticed something odd about the collapse postulate for two-particle systems.

It was nonlocal.

If two particles are initially in a nonseparable state,

then a measurement carried out on one can seemingly cause changes,

instantaneously, in the quantum mechanical description of the other,

no matter how far apart two particles are or what lies in between.

Suppose pair of electrons initially in state |A⟩

and measurement of color 1 carried out.

Outcome of measurement either green or magenta(equal probabilities).

Collapse postulate for two-particle systems

says as soon as measurement over,

state of 2 will be either |magenta⟩ (if 1 was green)

or |green⟩ (if 1 was magenta)

depending on what happened in measurement.

EPR said nonlocality is disposable artifact of particular mathematical formalism,

of particular procedure for calculating statistics of outcomes of experiments

—> must be undiscovered procedures,

which give rise to same statistical properties,

but are local (no infinite speeds necessary).

30 years later, Bell showed that their suspicion was **completely wrong**!

Bell's work(discuss later)

—> proof that any attempt to be realistic

about values of observables of the pair of electrons in state $|A\rangle$,

must necessarily be nonlocal.

Result was even more serious than that!!

Bell actually proves(as we will see) that there exists genuine nonlocality in actual workings of nature, however we attempt to describe it.

Nonlocality is feature of QM,

and via Bell's theorem

will necessarily be a feature of **every** possible manner of calculating

which produces the **same** probability predictions

(which are experimentally correct) as QM.

What is this quantum nonlocality?

First, in state $|A\rangle$,

statistics(probabilities) of outcomes of measurements on 2

depend nonlocally on outcomes of measurements on 1, and vice versa.

But do statistics of outcomes of measurements on 2,

when system in state $|A\rangle$,

depend nonlocally on

whether a measurement is actually carried out on 1 (and vice versa)?

Let us figure it out.

Suppose system in state $|A\rangle$,

and measure color 2.

$$|A\rangle = \frac{1}{\sqrt{2}} |green\rangle_1 |magenta\rangle_2 - \frac{1}{\sqrt{2}} |magenta\rangle_1 |green\rangle_2$$

Using |A⟩ in color basis (color-talk)

plus standard quantum-mechanical rules

for calculating probabilities of measurement outcomes

implies outcome of measurement on 2

equally likely to be green or magenta.

Suppose system is in state |A⟩

$$|A\rangle = \frac{1}{\sqrt{2}} |green\rangle_1 |magenta\rangle_2 - \frac{1}{\sqrt{2}} |magenta\rangle_1 |green\rangle_2$$

and measure color 1,

and then measure color 2.

Measurement of color 1 equally likely to be green or magenta.

If green,

collapse postulate says subsequent measurement of color 2

will be magenta,

and if magenta,

collapse postulate says that subsequent measurement of color 2

will be green.

So, when system in state |A⟩,

outcome of a measurement of color 2 is equally likely to be green or magenta

whether or not measurement of color 1 carried out first.

Suppose system is state |A⟩ and measure hardness 1,

and then measure color 2.

$$|A\rangle = \frac{1}{\sqrt{2}} |\text{hard}\rangle_1 |\text{soft}\rangle_2 - \frac{1}{\sqrt{2}} |\text{soft}\rangle_1 |\text{hard}\rangle_2$$

Using |A⟩ in hardness basis plus probability rules

says outcome of hardness measurement on 1 equally likely to be hard or soft.

If outcome of first measurement is soft,

collapse postulate plus probability rules say outcome of second measurement (color 2)

is equally likely to be green or magenta because 2 is **now hard**.

Same result true if outcome of first measurement is hard.

So here is where we are:

When system in state $|\mathbf{A}\rangle$,

outcome of measurement of color 2

is equally likely to be green or magenta,

whether measurement of color 1 is carried out first

or measurement of hardness 1 is carried out first

or no measurement on 1 is carried out

Probabilities of various outcomes of measurement on 2

do not depend in any way

on whether or not a measurement is made on 1 first.

Since the predictions of QM are correct,

there must be non-local influences in nature

and they must be of particularly subtle kind.

Outcomes of measurements do sometimes

depend non-locally on outcomes of other, distant measurements,

but outcomes of measurements invariably **do not** depend non-locally on

whether any other distant measurements actually get carried out.

Another way (tricky…you must think about this one) to say this:

Non-local influences are so subtle that they cannot be used

to transmit any signal containing information, non-locally,

between two distant points.

They cannot encode information you want to send

on decision to make a measurement or not to make one,

or on a decision about which measurement to make,

since no such decisions can have detectable non-local effects.

NOTE: Nonlocality existence is crucial. As we will see later, the solution to the measurement questions depends on the existence of nonlocality in QM.

Look at simple experiment (1980 with photons —> it took experimentalists 35 years!)

that refutes EPR and agrees with Bell's results (will derive shortly).

Will discuss experiment again later (more detail and mathematical rigor).

Before looking at experiment we must talk about electron spin!

Digression about Electron Spin

Electron spin is real-world measurable property of electrons.

Measuring devices are Stern-Gerlach(SG) boxes, for example.

Operator for electron spin $\hat{\mathbf{S}}$ has following properties.

1. When electron measured by SG device oriented in particular direction (say z)—> SGz box, measured values for electron spin observable in z-direction are only ±1/2.
   SGz box is represented by observable $\hat{S}_z$.

2. Same result holds for any other direction, in particular, x-direction with an SGx box.
   SGx box represented by observable $\hat{S}_x$ (only ±1/2 also).

3. Corresponding eigenvectors/eigenvalues given by

$$\hat{S}_z\,|U\rangle = +1/2\,|U\rangle \quad , \quad \hat{S}_z\,|D\rangle = -1/2\,|D\rangle \qquad \hat{S}_x\,|R\rangle = +1/2\,|R\rangle \quad , \quad \hat{S}_x\,|L\rangle = -1/2\,|L\rangle$$

From earlier: $$|U\rangle = \frac{1}{\sqrt{2}}(|R\rangle + |L\rangle) \quad , \quad |D\rangle = \frac{1}{\sqrt{2}}(|R\rangle - |L\rangle)$$

Can see from mathematical formalism,

electron spin in z and x directions are **mathematically** the same as color and hardness.

One very different feature of this real-world system

is that can orient SG boxes at any arbitrary angle θ (only 2 ways in color-hardness world)

where θ = 0° correspond to the +z direction,

θ = 90° correspond to the +x direction

and θ = 180° correspond to the –z direction and so on.

Thus, any arbitrary direction states can be written(from QM studied at junior level) as

$$|U_\theta\rangle = \cos\frac{\theta}{2}|U\rangle + \sin\frac{\theta}{2}|D\rangle \qquad |D_\theta\rangle = -\sin\frac{\theta}{2}|U\rangle + \cos\frac{\theta}{2}|D\rangle$$

EPR experiment with electron spin - this is actual experiment that was done.

Consider pair of electrons in entangled state like |A⟩

where we use the observable **spin(up/down in any direction)**.

$$|A\rangle = \frac{1}{\sqrt{2}}|U\rangle_1|D\rangle_2 - \frac{1}{\sqrt{2}}|D\rangle_1|U\rangle_2 \quad = \frac{1}{\sqrt{2}}|U_0\rangle_1|D_0\rangle_2 - \frac{1}{\sqrt{2}}|D_0\rangle_1|U_0\rangle_2$$

This state arises from the decay of an atom in Spin = 0 state of an atom where spin angular momentum is conserved : 0 —> (1/2) +(-1/2) The experiment has been done!

Electrons separate in physical space without changing the state vector.

We end up with two separated electron beams.

Each beam has equal numbers of |U⟩ and |D⟩ electrons (probability = 1/2 for each component).

Each electron in one beam

is **correlated** with a partner in the other beam(started from same atom)

since if measure electron in one beam in state |U⟩,

then partner(same magnet orientations) in other beam is in state |D⟩ due to

entanglement/collapse

and vice versa.

This nonseparable state remains nonseparable

no matter what basis (direction we measure spin) we use.

A state in an arbitrary direction basis is

$$|A\rangle = \frac{1}{\sqrt{2}} |U_\theta\rangle_1 |D_\theta\rangle_2 - \frac{1}{\sqrt{2}} |D_\theta\rangle_1 |U_\theta\rangle_2$$

Now suppose we measure beam 1 in direction θ_1 and beam 2 in direction θ_2

and count number of times (note these are not necessarily the same directions)

beam 1 up **and** beam 2 up

or $=$ a MATCH

beam 1 down **and** beam 2 down

beam 1 up **and** beam 2 down

or $=$ a MISS

beam 1 down **and** beam 2 up

QM says (straightforward but messy calculation) that the experimental results depend only on the angle difference

$$\varphi = \theta_2 - \theta_1$$

and **not separately** on θ_1 and θ_2 —-> Non-locality in action!!

That is the sign that a **correlation** exists!

Results are confirmed experimentally

no matter where the two detectors are located……

in same room, or on opposite sides of city or wherever!!!!

******* Experiment has now reached 144 km apart! *******

Suppose we code the results coming from the two detectors as follows.

If $\varphi = 0$, z–axes of 2 detectors are in same direction and get

$$beam\ 1 \rightarrow \uparrow\downarrow\uparrow\downarrow\downarrow\uparrow\downarrow\downarrow\downarrow\uparrow\uparrow\downarrow\uparrow\downarrow\downarrow\uparrow \ldots\ldots\ldots$$
$$beam\ 2 \rightarrow \downarrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\downarrow\downarrow\uparrow\downarrow\uparrow\uparrow\downarrow \ldots\ldots\ldots$$

i.e., ALL MISSES (remember if 1 is up, 2 is down in this case).

Get two opposite binary data sequences in two detectors

$$1010010001101001.....$$
$$0101101110010110.....$$

If $\varphi = 180^\circ$, z–axes of 2 detectors are in opposite directions and get ALL MATCHES, i.e.,

$$beam\ 1 \rightarrow \uparrow\downarrow\uparrow\downarrow\downarrow\uparrow\downarrow\downarrow\downarrow\uparrow\uparrow\downarrow\uparrow\downarrow\downarrow\uparrow \ldots\ldots\ldots$$
$$beam\ 2 \rightarrow \uparrow\downarrow\uparrow\downarrow\downarrow\uparrow\downarrow\downarrow\downarrow\uparrow\uparrow\downarrow\uparrow\downarrow\downarrow\uparrow \ldots\ldots\ldots$$

(up in 1 is in opposite direction to up in 2 or up in 2 is down in 1).

—> get identical(not opposite) binary sequences in this case.

Quantum mechanics predicts number of MATCHES, etc varies like

$$
\begin{aligned}
[\#\mathrm{MATCHES}] &= [\#\mathrm{ELECTRONS}]\sin^2\tfrac{\varphi}{2} \\
[\#\mathrm{MISSES}] &= [\#\mathrm{ELECTRONS}]\cos^2\tfrac{\varphi}{2} \\
[\#\mathrm{MATCHES}+\#\mathrm{MISSES}] &= [\#\mathrm{ELECTRONS}][\sin^2\tfrac{\varphi}{2}+\cos^2\tfrac{\varphi}{2}] \\
&= [\#\mathrm{ELECTRONS}]
\end{aligned}
$$

as it should!

Now set angle between detector axes to be $\varphi = 120^\circ$.

Quantum mechanics predicts(from our formalism)

$$[\#\mathrm{MATCHES}] = \frac{3}{4}\,\#\mathrm{ELECTRONS}$$

What would EPR say (you must remember EPR believe in locality)?

If $\varphi = 60^\circ$, then $[\#\mathrm{MATCHES}] = \frac{1}{4}\,\#\mathrm{ELECTRONS}$

(same as quantum mechanical prediction!)

So now EPR align detectors ($\varphi = 0^\circ$).

Then rotate detector 1 such that $\varphi = 60^\circ$

—> would again get 1/4 MATCHES.

So now EPR re-align detectors ($\varphi = 0^\circ$).

Then rotate detector 2(opposite direction) such that $\varphi = -60^\circ$ and also

—> would get 1/4 MATCHES.

EPR then predict $\frac{1}{4} + \frac{1}{4} = \frac{1}{2}$ MATCHES (not $\frac{3}{4}$)

i.e., two separate measurements cannot affect/change each other
—> prior results are unchanged
That is what locality means!!

EPR —> two measurements do not influence each other and therefore totals just ADD.

Experiment says answer is 3/4!!. Quantum mechanics is correct!

—> Local view of reality due to EPR cannot be correct.

Somehow information seems to be flowing

between detectors 1 and 2 NO MATTER WHAT WE DO.

Speed of information flow(if assume any signal exists…..) might be infinite

(best measurement so far $\approx 10^7$ c = 3×10^{15} m/ sec - but might not mean anything!!).

Important to emphasize that

have not observed any non-local interactions directly,

but have only indirectly demonstrated need for them.

Each measurement is separately local,

i.e., measure any spin direction in either detector

and always get 50-50 up and down

no matter what other detector is doing.

It is the sequences that differ(in a random way)

as we change the angles.

This means, as before,

that we cannot transmit any instantaneous messages(that would violate SR).

We only find out the sequences(messages) when we bring them together,

which takes some time.

So, it seems in QM, that my QM fist can punch your QM nose without being near your QM nose!!

Bohr and EPR - A Discussion

Bohr produced a reply to EPR within 2 months of paper publication.

His counterargument uses the **contextuality** of quantum theory

and the principle of complementarity to punch holes in EPR's definition of reality.

Complementarity

Bohr had a unique, deeply thought out, and very subtle view

on nature of science and interpretation of quantum theory.

Defies any easy and compact summary.

Some writings might —> instrumentalist viewpoint —-> oversimplification of his view.

Bohr was convinced that humans could never “picture” the inner mechanisms of atoms (Feynman will have more to say about this crucial point later).

His conclusion follows from nature of our human language.

Daily experience

—> ideas and concepts we deal with comfortably and express

in language restricted to classical world; but may not extend easily into quantum world,

hence, may need to abandon classical thinking and develop new concepts.

—> need new language as we already have seen

Snag(??)

Experiments rely on gathering together instruments and equipment

very much bigger than atomic scales

—> described by everyday language (including adding technical terms for brevity).

All measurements on atoms and particles

-> numbers read from dials on measuring apparatus (macroworld scale).

Essential to success of science that equipment be described in ordinary terms

—> makes possible for others to understand and reproduce experiment.

Caught in dilemma.

Experiments should not only be capable of being described,

essentially, in everyday language, but must also be so to make science possible.

Yet, when we gather the results of experiments together —> a description of atomic world

where we find that same everyday language and ideas fail to provide explanations.

Photons act as particles in some circumstances and as waves in others or **maybe neither**.

Can find appropriate mathematics to describe situation

but does not help us **“visualize or speak”** about photons.

Bohr hoped for a way out of trap <—> principle of complementarity.

He accepted that the two completely contradictory pictures of photon

could not be tied together into one picture, but he said that does not matter,

since we never do experiment that required the use of both pictures at **same time**.

We never ask questions requiring wave and particle answers at same time.

We never set context implying wave and particle properties at same time

—> Acceptable to employ this sort of "double think"

or using only one of pair of complementary views at any one time,

depending on what was most appropriate.

Actual true nature of photon can never be expressible in any language

which we we use to describe the classical world,

but we could shadow what was going on by keeping complementary pictures in mind.

—> concepts such as position, momentum, frequency, energy, and so on,

can be defined only in context of an experiment to measure them.

If want to talk about frequency of light (wave picture)

then need to do experiment where the wave picture appropriate.

If want to talk about momentum of photon (particle picture)

then do totally different sort of experiment.

By insisting, Bohr ensured that concepts such as momentum, energy, frequency, and so on,

remained rooted in classical world, while at same time acknowledging that

they would have only a limited application in quantum world.

Bohr wrote -

> **“quantum theory forces us to adopt a new mode of description designated as complementary in sense that any given application of classical concepts precludes simultaneous use of other classical concepts which in a different connection are equally necessary for elucidation of phenomena”.**

—> Copenhagen interpretation of QM

The emphasis on relating meaning of theoretical terms

to experiment where terms are measured gives Bohr's thinking an instrumental tinge.

However, careful reading of his writings

confirms that Bohr retained the ambition of a realist tempered by conviction that atomic world could not be described by limited human experience and language.

Bohr's Reply to the EPR Argument

Bohr's main point is that measurement of particle A

may not physically disturb state of particle B,

but measurement on A does **set up the context for any information about B**.

EPR used position and momentum instead of color and hardness (Same math).

$$|A\rangle = \frac{1}{\sqrt{2}} |x\rangle_1 |p\rangle_2 - \frac{1}{\sqrt{2}} |x\rangle_2 |p\rangle_1$$

To measure momentum of A —> must build specific device.

Let it interact with something with pre-determined momentum and see what change it makes.

If want to measure momentum of device, we find the uncertainty principle gets in way.

Accurately measuring momentum of device destroys any information about where exactly it is.

In which case, cannot act as position reference point as well.

This applies to both particles.

Measurement of position needs a fixed reference point to work from.

When set up equipment to measure momentum of particle A,

automatically prevent that equipment from forming

precise reference point for position measurement of A.

Set out to measure position of particle B.

Cannot compare position of B with anything of particle A

since cannot fix position of equipment at B's end relative to A.

Have effectively blurred position of particle B by measuring momentum of particle A.

As Bohr puts it:

"If choose to measure momentum of one of particles, lose through uncontrollable displacement inevitable in such measurement any possibility of deducing from behavior of particle position of apparatus, and thus no basis whatever for predictions regarding location of other particle. "

Similar argument applies if choose to measure position of A instead.

Accurate position measurements rely on fixed objects to act as reference points.

When particle A collides with device designed to record its position,

bound to be exchange of momentum between device and particle

i.e., it interacted with measuring device.

With device fixed in place cannot measure that momentum exchange.

Any attempt to measure momentum of measuring device,

so one can figure out how much it has been changed,

is bound to run into uncertainty problems.

Need an accurate fix of position of device to measure position of A,

so cannot be sure of momentum of device.

Ruins potential momentum measurement on B.

Momentum only known relative to object.

Cup of tea on table may be stationary to us,

but if on train passing through station

—> tea in rapid motion for observer standing on platform.

If set up momentum measurement of B, have no basis to compare with A.

Bohr explains:

By allowing uncontrollable momentum to pass from first particle into support, have cut ourselves off from any future possibility of applying law of conservation of momentum to system consisting of .... two particles, and therefore lost only basis for unambiguous application of idea of momentum in predictions regarding behavior of second particle.

Key difference between philosophical positions of EPR and Bohr

—> statement **“without disturbing the system”**.

To Bohr, a property, such as momentum or position,

has meaning **only in context** of experimental equipment used to measure it.

Particle B may not come in contact with any equipment at all;

in such case any information we have about it is useless - nothing one can do with it.

If want to use information,

say to predict where second particle will be after a while

and check that information experimentally,

then second particle has to come into contact with measuring equipment,

or some other stuff has to be linked to it.

As far as complementarity concerned,

Bohr insisted that concepts of momentum and position

work only within specific experimental context.

Two particles can live either in same context,

in which case limited comparisons of same properties can be made,

or not, in which case have two separate bundles of information

that cannot be related to one another.

Einstein and Bohr

Never able to reach common ground <—-> philosophical viewpoints entirely different.

Einstein - probabilities in QM unacceptable.

Use of statistics when calculating behavior of large numbers of particles different

All particles involved have properties such as position and momentum;

but so many —> calculations could not be done,

hence need to average and use statistics to calculate system probabilities.

Bohr - probability in QM fundamental,

—> not a consequence of inability to measure/calculate.

Einstein —> would be another layer of theory underneath QM

—> hidden variables of theory

—> no probability —> QM incomplete.

Einstein(sick of arguing/losing with Bohr) turns to Schrodinger

who also said that fundamentally statistical character of theory

is simply a consequence of incompleteness of description.

Then Schrodinger, on his own, Introduced idea of Entanglement

Schrodinger argued that state of system of two particles(that have interacted)

cannot be written as simple product of individual states for each particle.

$$|\text{Particle A interacting with B}\rangle \neq |A\rangle\,|B\rangle$$

State where particles have interacted

—> **entanglement** of individual states

where we cannot say with certainty which particle is in which state.

State such as

$$|\text{singlet}\rangle = \frac{1}{\sqrt{2}}\left(|U\rangle\,|D\rangle - |D\rangle\,|U\rangle\right)$$

is an entangled state: first particle either |U⟩ or |D⟩.

Basis of EPR argument(position/momentum), and Bohm variation(spins),

is that system **starts** in entangled state.

Schrodinger saw entanglement as the defining characteristic of QM

it is **what** —> QM is totally different from classical physics.

He emphasized view by discussing the disentanglement that happens when a measurement is made.

—-> **Entanglement disappears and particles then have properties!**

Measuring spin of particle A

—> can deduce spin of B if they are in singlet state.

—> Have collapsed combined state into one of two entangled possibilities.


In Schrodinger's view disentanglement is of key (possibly **sinister in his mind**) importance

since involved in every act of measurement.

Entanglement and Measurement

Remember earlier argument about measurement/collapse on entangled states.

Possible vertical spin states of electron moving toward S-G magnet are $|U\rangle$ and $|D\rangle$ in direction appropriate to a particular device.



If use QM to describe S-G magnet(the measuring device),

possible states are going to be $|\phi_n\rangle$ (n for neutral, resting state of apparatus),

$|\phi_U\rangle$ (U for UP showing apparatus registering UP state of electron),

and $|\phi_D\rangle$(D for DOWN registering DOWN state of electron).

Might know electron in $|U\rangle$ before reaches S-G magnet (passed through one beforehand)

—> state of combined system

before interaction of electron and S-G magnet = $|U\rangle|\phi_n\rangle$. **U-beam + pointer at 0**

After interaction, if measuring device any good,

state evolved into $|U\rangle|\phi_U\rangle$ U-beam + pointer to +1

in a process governed entirely by equations of quantum theory

(Schrodinger equation or time-evolution operator $\hat{U}(t)$).

$$|U\rangle\,|\phi_n\rangle \underset{\text{evolves into}}{\Rightarrow} |U\rangle\,|\phi_U\rangle \quad or \quad |U\rangle\,|\phi_U\rangle = \hat{U}(t)\,(|U\rangle\,|\phi_n\rangle)$$

time evolution

Note: no collapse of state has taken place,

just evolution of state of S-G apparatus from $|\phi n\rangle$ to $|\phi_U\rangle$

during interaction or dial pointer moving from 0 to +1.

i.e., a pointer moved!

i.e., measurement device state changed!

First time we are treating measuring device as described by QM,

rather than incomplete statements(as earlier)

that measuring device collapses state.

If take quantum theory seriously

as an underlying theory of reality, then

have to describe measuring device, (S-G magnets), in quantum way.

—> state **does not collapse** when a measurement takes place;

Quantum states simply entangle with states of equipment!

To see why this happens, consider the following…..

Assume we do not know initial state of electron.

Have to write state as

$$|\psi\rangle = a\,|U\rangle + b\,|D\rangle$$ where $|a|^2 + |b|^2 = 1$

Then, initial joint state of electron and S-G devices is $|\psi\rangle\,|\phi_n\rangle = (a\,|U\rangle + b\,|D\rangle)\,|\phi_n\rangle$

measurement device not entangled!

Based on linearity property of QM, evolves into

$$(a\,|U\rangle + b\,|D\rangle)\,|\phi_n\rangle \Rightarrow a\,|U\rangle\,|\phi_U\rangle + b\,|D\rangle\,|\phi_D\rangle$$

—> time evolution seems to gives the entangled state that we need to deal with.

Problem is that state has not collapsed in this view!!

Is there anything we can do?

Consider what happens when passing physicist comes to apparatus

to read off measurement results.

If we consider ourselves part of same universe as everyone else

—> passing physicist described by QM as well.

Must have states such as

$$|\text{physicist has not looked yet}\rangle\,, |\text{physicist sees UP}\rangle\,, |\text{physicist sees DOWN}\rangle$$

To record results, have to interact with apparatus so that

$$(a\,|U\rangle\,|\phi_U\rangle + b\,|D\rangle\,|\phi_D\rangle)\,|\text{physicist has not looked yet}\rangle \Rightarrow$$
$$a\,|U\rangle\,|\phi_U\rangle\,|\text{physicist sees UP}\rangle + b\,|D\rangle\,|\phi_D\rangle\,|\text{physicist sees DOWN}\rangle$$

and we become part of entanglement as well!

Probably why Schrodinger felt entanglement was **sinister**:

no way to stop it from spreading until


whole universe part of a stupendous entangled state.

—> **infinite regression - which is always a bad thing!**

Schrodinger was just blindly driving down the WRONG road as we will see!!

That Damned Cat - 1st Pass

The sorry tale of Schrodinger's cat
has assumed the status of a
quantum mechanical f**airy tale**.
Amazing how thoroughly this cat
has entered quantum lore,
especially when we consider that 1 quote



is only reference to unfortunate creature in all of Schrodinger's prodigious written output.

In the standard Schrodinger's cat experiment
a small radioactive device is linked to a trigger controlling the release of poison gas.

After a period of time, quantum state of radioactive device is in superposition of states

$$|\text{decayed}\rangle + |\text{not decayed}\rangle$$

some have decayed, some have not!

State of gas and ultimately state of cat are entangled with this state(or thought to be)
—> not possible to tell in quantum description
if cat is alive or dead(more details later).

Schrodinger introduced cat as a means of

illustrating the bizarre nature of entanglement

and superposition of quantum states.

(actually wanted to ridicule the ideas so people would replace them)

Remember when we introduced superposition

—> quantum states can be combinations of states that don't make sense classically

we did so because such things were needed to describe quantum systems.

Entanglement then follows directly once that is allowed!!.

Schrodinger's cat illustrates the associated problem of scale.

At what scale do classically forbidden combinations stop being allowed? Is there a size limit?

Experimentally photon seems to travel in two directions at once,

but surely a cat cannot be alive and dead at same time!

(or easier for cat lovers to deal with —- a cat cannot travel two paths at the same time!!)

If we look inside experiment

then can see if cat alive or dead, but that doesn't improve situation.

Our state just becomes entangled and so on…….**infinite regression……..**

This discussion which is ultimately probing the measurement problem

—> crucial issue for any interpretation of quantum theory.

Bohr **dissolved the problem (in his mind) by insisting** that at

some point the measuring device or cat or person

is described classically, **breaking** the chain of entanglement. **(Remember this point!)**

I hope that you have been keeping a list of all, the items I have been telling you to remember for later!!

According to Copenhagen interpretation,

the state vector for 2-particle quantum state cat

does not disentangle state

as the particles separate in space-time.

Instead of changing into two separate vectors, one associated with each particle,

state vector remains entangled

and, when a measurement made,

state collapses instantaneously **no matter how large** separation distance might be.

That is what our version of the theory presently says!

EPR's view of physical reality

—> if two particles are isolated from each other,

then they are **no longer described** by single state vector

when a measurement is made.

Reality EPR referring to

=> local reality

and ability of particles to separate(become space like)

into 2 locally real independent physical entities

=> **Einstein separability**.

In EPR experiment,

Copenhagen interpretation or standard view of quantum theory

denies that 2 particles are Einstein separable

and thus **denies they are locally real**

until a measurement is made on one or other,

at which point both become localized and **real** (the collapse process).

Schrodinger's Cat - Another Presentation

Earlier, the collapse of state vector was presented without reference

to the point in measurement process when collapse occurs.

One might assume the collapse occurs at moment the microscopic quantum system interacts

with macroscopic measuring device.

Is this assumption justified?

Now the macroscopic measuring device is **composed** of microscopic entities.

Interactions on microscopic level are supposed to **use QM** for their description.

Suppose a microscopic quantum system

described by state vector $|\psi\rangle$ interacts with measuring instrument

(= **device which responds to interaction with quantum system**

producing macroscopic results like pointers or dials)

with two possible measurement eigenstates $|\psi_+\rangle$ and $|\psi_-\rangle$.

These eigenstates combine with the macroscopic device

to **reveal** one of 2 possible outcomes of a measurement,

deflection of pointer to left (+ result) or right (- result).

Recognizing that the instrument consists of quantum particles,

we must describe state of instrument (repeating earlier arguments - I will do this many times)

before measurement by state vector $|\phi_0\rangle$,

corresponding to central pointer position.

Total state of quantum system+measuring instrument

before measurement described by $|\Phi_0\rangle$, is

$$|\Phi_0\rangle = |\psi\rangle\,|\varphi_0\rangle = \frac{1}{\sqrt{2}}\left(|\psi_+\rangle + |\psi_-\rangle\right)|\varphi_0\rangle = \frac{1}{\sqrt{2}}\left(|\psi_+\rangle\,|\varphi_0\rangle + |\psi_-\rangle\,|\varphi_0\rangle\right)$$

just multiply out…..

where $|\psi\rangle$ expressed in terms of measurement eigenstates

(assume = orthonormal basis) so that amplitudes are

$$\langle\psi_-\mid\psi\rangle = \langle\psi_+\mid\psi\rangle = \frac{1}{\sqrt{2}}$$

i.e., —> both final pointer results are equally probable.

Description of what happens if we treat

macroscopic measuring instrument as quantum object(repeat of earlier argument).

First, how does initial state $|\Phi_0\rangle$ evolve in time during act of measurement?

Earlier, the application of time evolution operator $\hat{U}$

allowed us to calculate state vector at later time —> $|\Phi\rangle$, as

$$|\Phi\rangle = \hat{U}\,|\Phi_0\rangle \qquad \longrightarrow \qquad |\Phi\rangle = \frac{1}{\sqrt{2}}\left(\hat{U}\,|\psi_+\rangle\,|\varphi_0\rangle + \hat{U}\,|\psi_-\rangle\,|\varphi_0\rangle\right)$$

What is effect of $\hat{U}$ on these states?

If instrument interacts with quantum system

already present in one of measurement eigenstates ($|\psi_+\rangle$ say),

then total system (quantum system+instrument)

must evolve into product quantum state $|\psi_+\rangle|\phi_+\rangle$, i.e.,

or, in other words, $$\hat{U}\,|\psi_+\rangle\,|\varphi_0\rangle = |\psi_+\rangle\,|\varphi_+\rangle$$ **pointer must move to +**

if quantum system in state(probability = 1)

corresponding to definite pointer position,

then measuring device state must evolve into state **a postulate!**

where pointer is pointing to proper place.

This actually what happens in the laboratory!!

Similarly, we must have $$\hat{U}\,|\psi_-\rangle\,|\varphi_0\rangle = |\psi_-\rangle\,|\varphi_-\rangle$$ **pointer must move to -**

Using the two special case results, we have for evolution of an arbitrary initial state

$$|\Phi\rangle = \frac{1}{\sqrt{2}}\left(|\psi_+\rangle\,|\varphi_+\rangle + |\psi_-\rangle\,|\varphi_-\rangle\right)$$ **using QM linearity**

where measuring device states $|\ \varphi_+\rangle$ and $|\ \varphi_-\rangle$

correspond to pointer ending up at + or - ——> A macroscopic event!

Result suggests that measuring instrument

evolves into superposition state in which pointer has equal probability

to point either to the left or right.

——> **It is not in a definite pointer state**.

Thus, **neither the** quantum system nor the pointer has a **definite value after the measurement**.

This state will remain a superposition (**pointer does not point**)

unless we allow for collapse

so that pointer can point (take on definite value)!

That is why the extra postulate is usually added.

If this were final state of system,

then the pointer should never settle down and point somewhere!

Even been suggested by some that it would **flutter** back and forth

between two macroscopically different pointer positions. **Wow - how dumb!**

Collapsing state vector of system + measuring-device

seems to require a further measurement.

Whole argument repeated ad infinitum(infinite regression again).

Are we therefore always locked into an endless chain of measuring processes?

At what point does chain stop

or at what point does state vector “collapse”

—> pointer actually pointing(**which we know it does!!!!**)?

Problem is created by the inability to obtain the collapse of state vector

using continuous, deterministic equation of motion

i.e., via the time evolution operator.

Schrodinger called state vector $|\Phi\rangle$ entangled because,

once generated,

it was impossible to separate it into constituent parts

except by invoking a nondeterministic collapse

i.e, some discontinuous process

i.e., some non-unitary time evolution.

Such collapse is **not accounted** for in equations of orthodox quantum theory

—-> why we had to add a postulate.

Paradox of Schrodinger’s cat is designed

to show this apparent absurdity

by shifting focus of difficulty from microscopic world to macroscopic world.

Repeat(more details/ideas): Cat placed in steel chamber together with

radioactive source, a detector, a hammer mounted on pivot and bottle of prussic acid.

Chamber is closed.

From amount of radioactive material in source and known time for decay(half-life),


expected that within one hour probability is 1/2 that one atom has disintegrated (decayed).


If atom does disintegrate,

then detector is triggered sending signal to release hammer

that smashes bottle releasing prussic acid which kills the cat.

The essential ingredients are shown below





Prior to actually measuring disintegration(decay),

state vector of radioactive atom = linear superposition of measurement eigenstates,

corresponding to physical states of undecayed atom and decayed atom.

Treating measuring instrument as quantum object and using equations of QM

—> superposition of two possible outcomes of measurement.

But what about cat?

These arguments (now will see trouble we are in…)

—> must express state vector of (system + cat) as linear superposition

of products of state vectors

describing disintegrated atom and dead cat

and describing intact atom and live cat, i.e.,

$$|\Phi\rangle = \frac{1}{\sqrt{2}} \left(|\text{no decay}\rangle \, |\text{live cat}\rangle + |\text{decay}\rangle \, |\text{dead cat}\rangle \right)$$

where, for example, state vector of dead cat

—> state corresponding to

triggered detector, released hammer, smashed bottle, released prussic acid **and** dead cat.

Prior to measurement, physical state of cat must be “blurred” (whatever that word means)

—-> neither alive nor dead but some **unknown** combination of both alive and dead states.

Can perform measurement on (system+cat) by opening chamber

and thus determining the physical state.

Do we suppose(imagine) that at that point (system+cat) collapses

and we record our observation that cat is either alive or dead as appropriate?

What happens?

QM is clear!

Only result that QM predicts is that

if set up N (N large) similar systems,

then when open N boxes after 1 hour

find 1/2 the cats alive and 1/2 the cats dead.

That is what actually happens experimentally.

Millions of measurements and probability always wins!

Although obviously intended to be somewhat tongue-in-cheek,

Schrodinger's paradox nevertheless brings attention to an important difficulty we must confront.

Copenhagen interpretation

—> elements of empirical reality defined by

nature(context) of experimental apparatus we construct

to perform measurements on quantum system.

Interpretation insists we **resist temptation** to ask

what physical state particle (or cat) actually in prior to measurement

as question is **without any meaning** within this interpretation of QM.

This positivist interpretation sits uncomfortably with some scientists,

particularly those with special fondness for cats.

Thus, some have accepted the EPR argument that quantum theory is incomplete.

They set about searching for alternative theory

that allows attaching physical significance

to properties of particles

without need to specify nature of measuring instrument,

That allows us to define independent reality

and that reintroduces strict causality.

Even though searching for such a theory

might be engaging in meaningless metaphysical speculation,

they believe that it is a search that has to be undertaken.

These are hidden variables people!

Have not succeeded with this approach and as will now see

Bell/EPR arguments say it is a futile search.

John Bell and Bohm's EPR

Einstein believed in **layer** of reality underneath quantum theory

that contains hidden variables (not yet discovered)

—> properties of particles —> return to having definite values before measurement.

When you look at old-fashioned mechanical clock face,

you see hands going around,

but may not be able to see hidden mechanisms driving them.

Careful study of moving hands might

—> can figure out properties of clock,

but would have much better understanding if get inside - see workings.

In Einstein's view quantum mechanics = theory of clock hands;

—> need another theory that looks inside at mechanism

(However, would that process ever end?).

At moment one cannot see mechanism inside quantum systems

<—> have not discovered any type of hidden variables involved.

Since do not have technique for measuring hidden variables (if they exist)

we cannot make definite predictions about outcome of experiments.

When repeat experiments, we are effectively

averaging over various (unknown) possible values of variables

and this ignorance is reason —> why we **end up** using probability in QM.

—> why probability is **not** intrinsic to world

according to Einstein and other hidden variable people!!!

In 1964, John Bell demonstrated to physics community that any hidden variable theory

with EPR-type entangled states would break

another of Einstein's most cherished assumptions

about way world works: **local causality**.

Causality = view that everything happening in world

is due to something else (cause) happening first, and then if we glue it to assumption

that nothing can travel faster than light (Einstein's theory of relativity)

—> **local causality**.

When particles interact, they might influence other's properties,

i.e., measure spin state A,

—> possible that result of measurement influences properties of B:

but —> there must be time delay.

Whatever connection exists between A and B,

information flows between them only at speeds less < c.

So, if measure A and then B somewhere far enough away,

fact that A has been measured

will not have reached B at time of measurement

and so can have no influence on measurement make of B

—> **local causality.**

Particle A can only influence other particles that happen to be in its locality.

Inside its light cone!

Bell wondered if a local hidden variable theory

could properly reproduce **all results** of quantum theory,

especially when dealing with EPR-type entangled states.

Now look again at Bohm-type EPR experiment using spin states(some repetition).

Assume existence of a set of hidden variables

that determine results of spin measurements on the particles.

If create two particles in so-called "singlet" state,

$$|A\rangle = \frac{1}{\sqrt{2}} |U\rangle_1 |D\rangle_2 - \frac{1}{\sqrt{2}} |D\rangle_1 |U\rangle_2$$

—> hidden variables must have **opposite** values for each particle.

In experiment, two particles, created in singlet state,

heading toward pair of separated S-G magnets.

Assume right-hand side S-G magnet much closer to place where particles created.

If we could read these hidden values would know particle A was UP (say)

and particle B was DOWN.

Don’t know(ignorant) values —> have to mix both possibilities

or say particles in an entangled state.

If we did a series of repeated measurements

and found A **always** UP and B **always** DOWN,

then could avoid using entangled state —> state **IS** only one part of entangled state.

Since we do not see this happening in these experiments

—> we must assume that values of hidden variables

are randomly determined when particles were created.

The spins are always opposite, but which value each one has comes out randomly.

Particle A now reaches S-G 1, magnet set to arbitrary angle.

As result of measurement either get UP along this S-G angle or DOWN.

Probability of each not 50:50 unless S-G angle happens to be vertical(depends on angle).

If want to calculate probability of UP along the S-G angle,

have to look at angle between S-G 1’s axis and original vertical.

Standard quantum theoretical calculation works and details not important here.

Point we are making.

While particles are able to interact

—> possible that values of hidden variables in one particle

alter values in other particle.

When A is measured, the interaction between A and S-G 1

might alter hidden variables,

and that information could be communicated to B and

—> influence what happens when measure B.

If set things up —> S-G 1 and S-G 2 far apart,

there is not time enough for any information to travel from S-G 1 to S-G 2

before particle B reaches S-G 2.

So only things that can influence what happens when particle B reaches S-G 2

are the values of its own hidden variables,

supposedly set when particles were **created**.

Compare now to quantum mechanical point of view.

When first measurement takes place at S-G 1,

entangled state of both particles will collapse producing 50:50 UP:DOWN result

no matter what angle of S-G 1 is.

Remember an entangled state has same nonseparable form in any basis.

If at the given **moment when** spin direction of A

manifests along line in direction of magnetic field in S-G 1 ,

then at **same moment** spin direction of B must manifest itself opposite.

If S-G 1 records UP, B arrives at S-G 2 in DOWN state along axis of S-G 1.

If S-G 2 is pointing along same axis, must record DOWN for B, **only same direction fixed!**

if not, -> UP and DOWN along its axis relative to how oriented with respect to S-G 1.

all this can be calculated in QM — it works

Crucial thing is collapse of state is affecting both particles at same time.

If hidden variable theory is going to match the effects of state collapse,

information from one measurement has to get to other

and if set up experiment correctly,

—> must do so faster than speed of light.

By assuming hidden variable theory had to obey rule of local causality,

Bell was able derive formula to compare results of each measurement

over long run of trials.

Experiment can then test Bell's formula

—> either quantum description is correct **or** local hidden variables can work.

Bell's Formula

Build experiment

—> have S-G magnets that can be set to one of three angles 0°, 45° and 90°.

Allow magnets to switch between the 3 angles randomly as experiment goes on.

Then simply sit back and watch singlet states

pass through both magnets and count up results.

If results from two magnets have no influence on one another, then should find that

$$N[\text{S-G1}(0^\circ)UP, \text{S-G2}(45^\circ)DOWN] + N[\text{S-G1}(45^\circ)UP, \text{S-G2}(90^\circ)DOWN]$$

$$\geq N[\text{S-G1}(0^\circ)UP, \text{S-G2}(90^\circ)DOWN]$$

N is number of pairs of particles counted satisfying set of conditions [...],

conditions are inside square brackets, i.e., in term like

$$N[\text{S-G1}(0^\circ)UP, \text{S-G2}(45^\circ)DOWN]$$

$$N[\text{S-G1}(0^\circ)UP, \text{S-G2}(45^\circ)DOWN]$$

N is number of times A registered spin UP

in S-G1 when set to 0°

along with particle B registering DOWN

when S-G2 set at 45°.

Although predicted formula looks complicated,

can see how works with aid of figure below, which shows how Bell's formula

$$N(A, \bar{B}) + N(B, \bar{C}) \geq N(A, \bar{C})$$

is derived

Note in formula, if A means passes test A, then $\bar{A}$ means fails test A.

Making things easier, we forget about particles for a moment. and use passing/failing tests - it is same mathematics - that is what is great about mathematics use in physics!

So, replace idea of measuring particle spins

with that of passing or failing 3 different tests - A, B, and C.

i.e., Bell's derived result is a very general result (just a property of numbers and mathematics) that always should be satisfied everywhere if his classical type of reasoning is valid!

The different tests are going to stand for different angles of S-G magnets.

Passing means spin UP and failing means spin DOWN along that angle.

First test set up will be equivalent of measurement of S-G1 and second one S-G2.

Also going to assume that result of one test has no effect on next

—> assuming local causality

Symbol like N(A, $\bar{C}$) going to stand for number of times pass test A **first**

and **then** fail ($\bar{C}$) test C,


or in experimental terms number of pairs that were spin UP in S-G 1 set to angle A


and spin DOWN in S-G 2 set to angle C.

Now look at figures.

Figure broken up into three pie charts,

each stands for one part of formula.





Each pie chart divided by vertical

and horizontal lines.

Left-hand side of each vertical line


stands for passing test B,

right-hand side for failing test B.

Above horizontal line

stands for passing test C

and below for failing test C.



Scattered around segments of pie charts

are passing and failing tests A.

All three possibilities are


equally represented (same area) in pie charts.


In top left chart,

lined segments represent times

that test A passed and test B failed:

N(A,$\bar{B}$).





In top right chart, lined segments

are number of occasions that

B was passed and C failed: N(B,$\bar{C}$).

Finally, in bottom chart lined segments


are passing A but failing C: N(A,$\bar{C}$).


Look closely and compare three charts,

—> marked segments in bottom chart have


already been covered by marked regions in other

two charts - with area to spare!






$N(A,\overline{B})$



$N(B,\overline{C})$



$N(A,\overline{C})$

So, $N(A,\bar{B}) + N(B,\bar{C}) \geq N(A,\bar{C})$

—> d'Espagnat's version of Bell formula.

Startling thing about this demonstration is that it shows how few assumptions need to be used in deriving Bell's formula and that it is simple classical property of numbers and standard logic.

Formula itself has nothing to do with quantum theory.

It's simply a relationship between numbers that fall (using standard classical logic) into certain categories.

Classical physics does obey this formula!!

Amazingly, as we will see, quantum physics does not obey this formula!!

That last staement should make you sit up in your chair!!!!

Aspect's Experiment -1st pass

Although d'Espagnat's version of Bell's formula

might be relatively easy to demonstrate mathematically,

it is not easy to test in experiment.

Different version was used for test, quote here which we will not derive(too complicated):

$$|E(a,b) - E(a,b')| + E(a',b) + E(a',b') \leq 2$$

is valid for a local hidden variable theory (with E(a,b) as defined below).

This expression is once again simply a relationship between numbers that fall (using standard classical logic) into certain categories.

Vertical lines in first term simply —> absolute value.

Collection of terms on left-hand side = S parameter, Bell result —> $S \leq 2$.


Idea of experiment

—> measure S value

and see if fits quantum theoretical description which —-—> $S > 2$.

Terms in expression directly relate to experimental arrangement

in figure below which shows Aspect's experiment to test Bell's formula.



In 1982(47 years after EPR) Alain Aspect's team at Institute for Optics in Paris

performed significant test of Bell's formula(22 years after - a difficult experiment)

since it was first experiment

that allowed detector angles


to be changed while particles were in flight (figure).

Describe experiment with more details later……

For now, source S produces pair of photons in combined singlet state.

Either side of source are equivalent of S-G magnets for photons (called polarizers).

These devices, labelled a and b, check polarization of photons (equivalent of spin).



If photon entering device A(left side) is spin UP along device's angles,

detected by D+, otherwise picked up by D–.

Same true on right-hand side in device B.

Devices can be set to different angles:

left-hand one can be set to either angle a or a', right-hand one to b or b'.

Each of four detectors feeds into coincidence monitor (CM),

which counts number of times get each possibility (in coincidence => simultaneously):

N_{++} which is UP at A and UP at B N_{--} which is DOWN at A and DOWN at B

N_{+-} which is UP at A and DOWN at B N_{-+} which is DOWN at A and UP at B

Experiment run while allowing a and b

to randomly switch their angles between a and a' and b and b', respectively.

Finally use these counts to construct

$$E(a,b) = \frac{(N_{++} + N_{--}) - (N_{+-} + N_{-+})}{N_{++} + N_{--} + N_{+-} + N_{-+}}$$

Sort of an expectation value for results.

Aspect's experiment conducted with angles a = 0°, a′ = 45°, b = 22.5°, and b′ = 67.5°;

a combination predicted by quantum mechanical calculation

to break Bell's formula by largest amount.

In all detail,

it is important not to lose sight of the purpose of the experiment

and significance of Bell's formula.

If experimental information fits Bell's formula, so that S ≤ 2,

then quantum mechanics is **wrong**

and could possibly be replaced by hidden variable theory

subject to condition of local causality.

If S > 2 local causality is **broken**

and it seems that results of one measurement can **influence** results of another,

although even a signal traveling at speed of light would

not have chance to reach second measurement in time.

Experiment then represents a **crucial step** in our understanding of quantum reality.

Working with angles in Aspect's experiment, QM predicts S = 2.828.

Correcting for detector efficiency (sometimes miss things)

—> S = 2.70 = prediction of QM.

Aspect measured S = 2.697 ± 0.015

—> agreement with QM and breaking Bell's formula($S \leq 2$).

Since Aspect's pioneering experiment,

other(more accurate) such experiments —> quantum theory correct always.

In 2001, team from University of Innsbruck used

two detectors (Alice and Bob) 400 m apart to check formula.

(Now, experiments up to 144 kilometers apart)

Their detection efficiency was such that quantum mechanical prediction was S = 2.74

and they measured S = 2.73 ± 0.02.

Conditions of 2001 experiment place an even greater restriction on things,

since choice of angle at each detector was truly random,

and angles were set after photons had left source

and results stored at each detector

and compared **only after** experiment had been completed.

All of these totally rule out any communication between two photons,

or even two detectors, that could explain results.

BELL'S THEOREM - An alternative presentation to enhance understanding

Bell demonstrated that under certain conditions

quantum theory and local hidden variable theories

predict different results for same experiment on pairs of correlated particles.

This difference,

which is intrinsic to all local hidden variable theories

and is independent of exact nature of theory,

is summarized in Bell's derived inequalities.

This proof forced questions about hidden variables to immediately change character.

They were no longer academic questions about philosophy

but practical questions of profound importance for quantum theory.

Choice between quantum theory and local hidden variable theories

was no longer matter of **taste**, but matter of **correctness**.

Now for Bertlmann's Socks

Derive Bell's theorem with help of famous Dr Bertlmann

(this whimsical story is by Bell himself invented to show how "classical" it is).

Any philosopher in the street,

who has not suffered through a course in quantum mechanics,

is quite unimpressed by Einstein-Podolsky-Rosen correlations.

She can point to many examples of similar correlations in everyday life.

Bertlmann decides to subject his left socks (socks A from now on)

to 3 different tests:

$$\text{test } a \text{ washing for 1 hour at } 0.0^\circ\, C$$
$$\text{test } b \text{ washing for 1 hour at } 22.5^\circ\, C$$
$$\text{test } c \text{ washing for 1 hour at } 45.0^\circ\, C$$

He is particularly concerned about numbers of socks A that survive intact (a + result)

or are destroyed (a – result)

by prolonged washing at these different temperatures.

He denotes number of socks that pass test a and fail test b as

$$n[a+b-]$$

Being a theoretical physicist,

he knows can discover simple relationships between such numbers

without actually performing tests using real socks and real washing machines.

Makes study inexpensive and more attractive to research sponsors.

He reasons as follows: n[a + b–]

can be written as sum of numbers of socks belonging to two subsets,

one where individual socks pass test a, fail b and pass c

and one where socks pass test a, fail b and fail c, i.e.,

$$n\left[a+b-\right]=n\left[a+b-c+\right]+n\left[a+b-c-\right]$$

This works probabilistically because

$$P(C)+P(not\ C)=1$$

Also where individual socks pass test b, fail c and pass a

and one where socks pass test b, fail c and fail a and

$$n[b+c-]=n[a+b+c-]+n[a-b+c-]$$

because $P(A)+P(not\ A)=1$

From first equation follows that

$$n\left[a+b-\right] \geq n\left[a+b-c-\right]$$

since all numbers involved are ≥ 0.

Similarly, from second equation follows that

$$n\left[b+c-\right] \geq n\left[a+b+c-\right]$$

Adding last two equations gives result

$$n\left[a+b-\right]+n\left[b+c-\right] \geq n\left[a+b-c-\right]+n\left[a+b+c-\right]=n\left[a+c-\right]$$

or

$$n\left[a+b-\right]+n\left[b+c-\right] \geq n\left[a+c-\right]$$

derivation only used standard logic and inequalities among numbers

At this stage, Dr. Bertlmann notices a **flaw** in his reasoning,

which all of you will, **of course**, have spotted right at beginning.

Subjecting one of socks A to test a

will necessarily change irreversibly its physical characteristics such that,

even if it survives test,

it may not give result for test b that might be expected of brand new sock.

And, of course, if sock fails test b, it will simply not be available(destroyed) for test c.

Thus, numbers like n[a + b–], etc, have **no** practical (cannot measure in real world) relevance.

Bertlmann now remembers his socks always come in pairs.

He assumes that, apart from differences in color,

physical characteristics of each sock in pair are identical.

Thus, test performed on right sock (sock B)

can be used to predict what result of same test would be

if test had been performed on left sock(sock A),

even though test on A not actually carried out

Remember this is a counterfactual statement - relating to or expressing what has not happened or is not the case!. No test is done on A!

Must further assume that whatever test

he chooses to perform on B sock in no way

affects the outcome of any other test he might perform on A sock,

but this seems so obviously valid that

he does not give it any thought whatsoever!!!!

Can you figure out where we are in the EPR argument? Think causality…….

Bertlmann now devises three different sets of experiments

to be carried out on three samples

each containing same total number of pairs of socks.

In experiment 1, for each pair,

sock A is subjected to test a and sock B is subjected to test b .

If sock B fails test b, this implies that sock A would also have failed test b

had it been performed on sock A.

Thus, number of pairs of socks for which sock A passes test a and sock B fails test b,

which we denote by $N_{+-}(a, b)$

must be equal to (hypothetical) number of socks A which pass test

a and fail test b(same sock), i.e., $N_{+-}(a, b) = n\,[a + b-]$

In experiment 2, for each pair,

sock A is subjected to test b and sock B is subjected to test c.

Same kind of reasoning allows Bertlmann to deduce that $N_{+-}(b, c) = n\,[b + c-]$

where N+–(b,c) denotes number of pairs of socks

for which sock A passes test b and sock B fails test c.

Finally, in experiment 3, for each pair,

sock A is subjected to test a and sock B is subjected to test c.

In a similar manner, Bertlmann thus deduces that $N_{+-}(a, c) = n\,[a + c-]$

where N+–(a,c) denotes number of pairs of socks

for which sock A passes test a and sock B fails test c.

Experimental arrangements summarized below.

Experiment	Sock A Test	Sock B Test
1	a	b
2	b	c
3	a	c

Bertlmann, now using revised calculations, concludes again that we must have inequality

$$N_{+-}(a,b) + N_{+-}(b,c) \geq N_{+-}(a,c)$$

to represent the experiments.

Bertlmann generalizes this result for any batch of pairs of socks

by dividing each number by total number of pairs of socks

(same for each experiment)

to arrive at frequencies with which each joint result was obtained.

He identifies the frequencies with probabilities

for obtaining results for experiments to be performed

on any batch of pairs of socks that, statistically, have same properties.

Thus, he finds that

$$P_{+-}(a,b) + P_{+-}(b,c) \geq P_{+-}(a,c)$$

—> Bell's inequality for this experiment.

Again, just standard classical logic and properties of numbers!

Now follow above arguments again, replacing

1. socks with photons
2. pairs of socks with pairs of entangled photons
3. washing machines with polarization analyzers
4. temperatures with polarizer orientations

we still arrive at the standard Bell's inequality,

i.e., only change words in our description of experiments and not conclusions!

Three tests now refer to polarization analyzers set

with their vertical(optic) axes oriented at

a→0.0°, b→22.5°, and c→45.0°.

Different experimental arrangements summarized as follows:

Experiment	Photon A Angle	Photon B Angle	Angle Difference Δ
1	$a = 0.0°$	$b = 22.5°$	$22.5°$
2	$b = 22.5°$	$c = 45.0°$	$22.5°$
3	$a = 0.0°$	$c = 45.0°$	$45.0°$

Probabilities **predicted** by quantum theory

for given angle **difference** Δ between polaroids in each test given(using QM) by

$$P = \frac{1}{2}\sin^2\Delta$$

i.e., remember matches and misses arguments.

Putting angles above into Bell inequality get

$$\frac{1}{2}\sin^2 22.5° + \frac{1}{2}\sin^2 22.5° \geq \frac{1}{2}\sin^2 45.0°$$

or $0.1464 \geq 0.2500$ which is obviously incorrect!

Thus, for these particular arrangements of polarization analyzers,

the probability formula from quantum theory predicts results

that violate Bell's inequality.

However, all the quantum mechanical probability formula results

agree with all experimental observations!!

Implications:

Einstein could not have seen where EPR argument would lead:

from Bohm's simpler and more practical version,

to Bell's analysis in terms of local hidden variables

to Aspect's experiment and the results that have been produced since.

Quantum theory has survived every test thrown at it and thanks to Bell's argument,

now seems impossible that it will ever be replaced by any local hidden variable theory.

Of course, the option of hidden variable theory

that does not obey local causality is still on the table.

Experimental results showing correlations

between two detectors **could be explained** by communication between two particles

traveling faster than the speed of light,

something that would hardly have made Einstein feel any better about things.

Quantum theory survives Bell's test

because entangled states **collapse** at first measurement.

Remarkably this collapse affects **both** particles in entangled state

no matter how **far apart** they may be.

Bohr simply shrugged this off

as a consequence of having to use the

same experimental context for both particles,

an argument that is logically compelling

but does not seem to do justice to the remarkable properties of entangled states.

Bell's formula is broken by entangled state,

so if want to take realistic view of what's happening,


state collapse has to be real physical change


happening everywhere at once.

Perhaps we ought to stop thinking of

two entangled particles as being separate at all.

Comparison of quantum mechanics and Bell inequality from 1st experiments is shown in figures below:





Clearly, quantum mechanics is correct. Even **better** data exists now!

Most important assumption made in reasoning

which led to inequality was **Einstein separability**

or **local reality** of photons(or socks).

Inequality is quite independent of nature of any local hidden variable theory

that could be devised.

Conclusion is inescapable.

Quantum theory is incompatible with any local hidden variable theory

and hence **incompatible** with any form of **local reality**.

Means that any theory that has **same** probabilistic predictions as quantum mechanics

must be nonlocal.

Should not be too surprised by this result.

Predictions of quantum theory are based

on properties of 2-particle state vector which,

before collapsing into one of measurement eigenstates,

is delocalized or entangled over **whole** experimental arrangement.

2 particles are, in effect,

always in contact prior to measurement

and **can** therefore exhibit a degree of **correlation**

impossible for 2 Einstein separable (locally realistic) particles.


Now for some details of the Aspect experiments.

The Details of the Aspect Experiment at Orsay - Alain Aspect -Institute of Optics - Paris 1980

Light source is beam of calcium atoms, excited by two focused laser beams having wavelengths $\lambda' = 406$ nm and $\lambda'' = 581$ nm respectively.

Two-photon excitation produces state having the quantum number J = 0 (angular momentum).

When it decays, this state emits two monochromatic photons having wavelengths $\lambda_1 = 551.3$nm and $\lambda_2 = 422.7$nm respectively, in a cascade of two electronic transitions from the initial J = 0 level to the final J = 0 state, passing through an intermediate J = 1 state, as shown in figure which shows excitation and decay of calcium atom.




Calcium based light source - Energy Levels

Mean lifetime of intermediate state is 4.7ns.

Simplify terminology, call $\lambda_1 = 551.3$ nm light green, and $\lambda_2 = 422.7$ nm light violet.

Polarizers in experiment = Wollaston prisms is shown in figure where we can see its two-valued response.



Wollaston prism is made of quartz or of calcite.

It splits incident beam of natural (unpolarized) light into two beams of equal intensity, polarized at 90° to each other.

If only single unpolarized photon incident, it emerges either in state $|x\rangle$, with probability 1/2,


or in state $|y\rangle$, with probability 1/2 (50-50).


Thus, response of system is **two-valued**.

Photons are detected by photomultiplier tubes (PM) downstream from prism.

Every electric pulse from detectors corresponds to passage of a photon, allowing photons to be counted.

Experimental layout sketched in figure, which shows first Orsay experiment.





It uses coincidence circuit which registers an event whenever two photons are detected simultaneously.

In this way four separate counts are recorded simultaneously, over some given period of time.

In simpler EPR experiment(Bohm), only 2 possible responses are $(+1,-1)$ or $(-1,+1)$.

In situation realized by Aspect four different responses are possible.

N_{++} the number of coincidences corresponding to $\alpha = 1$ and $\beta = 1$, that is, to $\alpha\beta = 1$

N_{+-} the number of coincidences corresponding to $\alpha = 1$ and $\beta = -1$, that is, to $\alpha\beta = -1$

N_{-+} the number of coincidences corresponding to $\alpha = -1$ and $\beta = 1$, that is, to $\alpha\beta = -1$

N_{--} the number of coincidences corresponding to $\alpha = -1$ and $\beta = -1$, that is, to $\alpha\beta = 1$

Resolving time of coincidence circuit is 10ns, meaning that says two photons coincident if they are separated in time by no more than 10ns. Mean lifetime of the intermediate state of the calcium atom is 4.7ns —-> on average only one photon per mean lifetime in the experiment.


Therefore, after lapse of 10ns = more than twice mean lifetime, almost all atoms have decayed (actually 88%). In other words, efficiency of coincidence counter is very high. Even better now!

Experiment consists in counting, over some given time interval, the four kinds of coincidence:

N_{++}, N_{+-}, N_{-+} and N_{--}. The total number of events is $N = N_{++} + N_{+-} + N_{-+} + N_{--}$.

Accordingly, the different kinds of coincidence have probabilities

$$P_{++} = N_{++}/N \text{ corresponding to } \alpha\beta = 1$$
$$P_{+-} = N_{+-}/N \text{ corresponding to } \alpha\beta = -1$$
$$P_{-+} = N_{-+}/N \text{ corresponding to } \alpha\beta = -1$$
$$P_{--} = N_{--}/N \text{ corresponding to } \alpha\beta = 1$$



and the measured average of αβ is

$$\langle\alpha\beta\rangle = \frac{N_{++} - N_{+-} - N_{-+} + N_{--}}{N}$$


Each set of four coincidence counts corresponds to one particular relative angular setting of (A, B) prisms, and yields a mean value $\langle\alpha\beta\rangle$. But in order to determine the correlation function $\langle\gamma\rangle$ (equivalent to S parameter) used in the BCHSH form of Bell inequality, we need four mean values $\langle\alpha\beta\rangle$.

Therefore, we choose, in succession four different settings(explain) as shown in figure above four counting runs then yield the four mean values $\langle\alpha_1\beta_1\rangle$, $\langle\alpha_1\beta_2\rangle$, $\langle\alpha_2\beta_1\rangle$, $\langle\alpha_2\beta_2\rangle$ which then determine the value of $\langle\gamma\rangle$ via

$$\langle\gamma\rangle = \langle\alpha_1\beta_1\rangle + \langle\alpha_1\beta_2\rangle + \langle\alpha_2\beta_1\rangle - \langle\alpha_2\beta_2\rangle$$

The Results of the First Experiment at Orsay

Results of first Orsay experiment shown in figure. Angle θ which specifies relative setting of polarizers plotted horizontally, and mean value $\langle\gamma\rangle$ vertically.



Correlation function predicted by mathematics of quantum mechanics is

$$\langle\gamma\rangle = 3\cos 2\theta - \cos 6\theta$$

It is drawn as solid curve on graph. According to Bell-Clauser, Horne, Shimony and Holt (BCHSH) version of inequality

$$-2 \le \langle\gamma\rangle \le 2$$

so that any hidden-variable theories exclude cross-hatched regions of plane, which correspond to $\langle\gamma\rangle > 2$ or $\langle\gamma\rangle < -2$. Notice experiment goes into cross-hatched regions violating bounds!!

Clearly, can be no doubt that BCHSH inequality is violated; many of experimental points fall outside interval $[-2, 2]$. At point where violation is maximal ($\theta = 22.5^\circ$), one finds

$$\langle \gamma \rangle = 2.70 \pm 0.015$$

which represents departure of over **40 standard deviations** from extreme value of 2. What is even more convincing is precision with which experimental points lie on curve predicted by quantum mechanics - latest experiments are even more dramatic (curve and data exactly same!).

Evidently, for EPR scenario, must conclude not only that hidden-variable theories fail, but also that quantum mechanics is positively the right theory for describing observations.

The Relativistic Test

EPR experiment just described shows that measurements in A and B are **correlated**.
What is origin of correlations?


According to quantum theory, before measurement each particle pair constitutes single system extending from A to B, whose two parts are **non-separable and correlated**. This interpretation corresponds to a violation of Bell's inequality and agreement with experiment.

According to hidden-variables theories, particle pair characterized, at instant of decay, by its hidden variable, which determines correlation between polarizations measured in A and B. This interpretation satisfies Bell's inequality but disagrees with experiment.

Accordingly, Orsay experiment supports quantum interpretation (in terms of correlation between two parts A and B of a single system).

However, to reinforce conclusion, must ensure that no influence exerted in ordinary classical sense through some interaction propagated between two detectors A and B, that is, no influence which might take effect after decay at S, and which might be responsible for correlation actually observed.

Let us therefore examine Orsay apparatus in more detail as in figure below where we attempt to test **Einsteinian non-separability**.



When detectors at A and B record a coincidence, means that both have been triggered within a time interval of at most 10 ns, the resolving time of circuit.

Could it happen that, within this interval, A sends to B a signal capable of influencing the response of B?

In most favorable case, such a signal would travel with speed of light in vacuum, which according to relativity theory is upper limit on propagation speed of information, and thereby of energy.

To cover distance AB, which is 12 m in figure, such a signal would need 40 ns.

This is too long by at least 30 ns, and rules out any causal links between A and B in the sense of classical physics.

One says that interval between A and B is **space-like**.

One of advantages of Orsay experiment is that it uses a very strong light-source, allowing sufficient distance between detectors A and B while still preserving reasonable counting rates.

By increasing distance AB step by step, Aspect could check that correlation persists, even when interval between A and B becomes space-like.

This is check that guarantees that two-photon system is non-separable irrespective of distance AB.

It has become customary to speak of principle of **Einsteinian separability** in order to denote absence of correlations between two events separated by a space-like interval.

This is principle that Orsay experiment invites us to reconsider, even though our minds, used to the world at the macroscopic level, find it difficult to conceive of two microscopic photons 12m apart as a single indivisible object (**remember this point for later!!!**).

The Final Stage of the Experiment at Orsay

Though results of first Orsay experiment are unarguable and clear-cut, conclusion they invite is so startling that one should not be surprised at appearance of a last-ditch objection, which as it happens gave experimenters a great deal of trouble.

In the preceding section we discussed possible role of interactions between A and B operating after decay at S, and duly eliminated objection.

But one can also ask whether correlations might be introduced through an interaction operating **before** decay - before experiment even starts!!.

We could imagine that the decay itself is preconditioned by setting of detectors A and B, such influences taking effect through exchange of signals between detectors and source.

No such mechanism is known a priori, but we do know that, if there is one, then Einsteinian non-separability would cease to be a problem, because the mechanism could come into action long before the decay, removing any reason for expecting a minimum 30 ns delay.


Though such a scenario is very unlikely, objection is a serious one and must be taken into account;

To get around it, experimenter must be able to choose orientation of detectors A and B at random after decay has happened at S.

In more picturesque language,

we would say that two photons must leave source without knowing orientations of polarizers A and B.

Briefly put, this means that it must be possible to change detector orientations during 20 ns transits over paths SA and SB.




Solution adopted at Orsay employs periodic switching every 10ns.

These changes are governed by two independent oscillators, one for channel A and one for channel B.

The oscillators are stabilized, but however good the stabilization it cannot eliminate small random drifts that are different in the two channels, seeing that the oscillators are independent.

This ensures that the changes of orientation are random even though the oscillations are periodic, provided the experiment lasts long enough (1 to 3 hours).

The key element of the second Orsay experiment is the optical switch shown in figure below.



In a water tank,

a system of standing waves is produced by electro-acoustic excitation

at a frequency of 25 MHz

(corresponds to 10 ns between switchings).

now even better!!

Fluid keeps changing from a state of perfect rest to one of maximum agitation and back again.

In state of rest, light beam is simply transmitted.

In state of maximum agitation, fluid arranges itself into a structure of parallel and equidistant plane layers, alternately stationary (nodal planes) or agitated (antinodal planes).

Thus, one sets up a lattice of net-like diffracting planes.


Diffracted intensity is maximum at so-called Bragg angles, just as in scattering from a crystal lattice.

Here, light beam is deviated through 10^{-2} radians (angles in figure are exaggerated for effect).

The two numerical values, 25 MHz and 10^{-2} radians, suffice to show magnitude of the technical achievement.

With the acoustic power of 1 watt, the system functions as an ideally efficient switch.

The second Orsay experiment (using optical switches) is sketched in Figure below.



Now using Optical Switches

In this set-up, photons a and b leave S without knowing whether they will go - the first to A_1 or A_2 and the second to B_1 or B_2.

Second experiment is less precise than first, because light beams must be very highly collimated in order to ensure efficient switching.

Nevertheless, its results exhibit an unambiguous violation of Bell's inequality, reaching 5 standard deviations at the peak;

Moreover the results are entirely compatible with the predictions of quantum mechanics.

Some final thoughts about all these ideas before we proceed to the quantum measurement discussion.

Bell's theorem follows from classical rules for probability.

QM does not agree with the theorem.

The way QM probabilities work is different.

For example: Quantum mechanics make us wonder:

Is FALSE the SAME as NOT TRUE?

The statement

"there is a 70% chance that the proposition - if A is measured, then the result will be a - is true"

is quite different from

"if A is measured, then there is a 70% chance that the result will be a".

It is the latter that is intended in quantum mechanics by the statement:

Prob(A=alWave function) =0.7

QM works and NOTHING ELSE works!

Statements from the Pioneers

Bohr once declared when asked whether quantum mechanical algorithms could be considered as somehow mirroring an underlying quantum reality: He said :

> **There is no quantum world. There is only an abstract quantum mechanical description. It is wrong to think that the task of physics is to find out how Nature is. Physics is concerned only with what we can say about Nature.**

Heisenberg said:

> **In the experiments about atomic events we have to deal with things and facts, with phenomena that are just as real as any phenomena in daily life. But the atoms or the elementary particles are not as real; they form a world of potentialities or possibilities rather than one of real things or facts.**

Jordan declared:

> **That observations not only disturb what has to be measured, they produce it. In a measurement of position of an electron, the electron is forced to a decision.**
>
> **We compel it to assume a definite position; previously it was, in general, neither here nor there; it had not yet made its decision about a definite position.**

Clearly, since measurement is so crucial to QM, let us now look at various aspects of measurement theory.

Quantum Measurement

We now review the standard formulation of quantum mechanics that we have developed, which is based on the 4 postulates listed below(with some embellishments):

- All physical systems are represented by ket vectors $|\psi\rangle$ normalized to 1, i.e., $\langle\psi|\psi\rangle = 1$. The ket labels represent **everything that we know about the system**.

- Measurable properties of physical systems are represented by linear operators called **observables**.

So restating part of the first postulate, the ket labels represent **the values of all observables of the system that have been measured - we PREPARED the state!!!**

If a vector associated with a particular state $|\psi\rangle$ is an eigenvector, with eigenvalue α, of operator $\hat{A}$ associated with a particular measurable property of the system, i.e., if $\hat{A}\,|\psi\rangle = \alpha\,|\psi\rangle$, then the system in that state **definitely** has the value α of that measurable property.

This implies that if one performs a measurement corresponding to the observable represented by A on a system in the state $|\psi\rangle$, then with **certainty** (probability = 1) the measurement yields the value α for that measurable property.

Observables are represented by **Hermitian** operators (real eigenvalues).

Since the eigenvectors of any Hermitian operator form a complete, orthonormal set, they can be used as a **basis** for the Hilbert space of the system

Finally, if the system is in the state $|\psi\rangle$ and one measures an observable $\hat{B}$, where $|\psi\rangle$ is **not** an eigenvector of $\hat{B}$, then the **only possible results** of the measurement are one of the eigenvalues $\{b_k\}$ of $\hat{B}$.

- Dynamics of state vectors

  The state vectors of any system change with time via **deterministic** laws (similar to classical rules).

We define the **time evolution or time development operator** that governs how a state vector changes in time by the relationship

$$|A, t + \Delta t\rangle = \hat{U}(\Delta t)\,|A, t\rangle$$

or the state vector at time t+ Δt is given by the time evolution operator (Unitary) $\hat{U}$ operating on the state vector at time t.

In general, the ket labels(which contain whatever we know (have measured) about the state) are the only thing that changes.

The time evolution operator is a unitary operator since the state vector must remain normalized to 1, i.e., the vector length cannot change, and this is guaranteed by the use of a unitary operator.

The only changes to state vectors in quantum mechanics are changes in direction (phase).

The time evolution operator is related to the energy operator (this follows from time translation invariance)

$$\hat{U}(t) = e^{-i\hat{H}t/\hbar}$$

- Connection with Experiment/Measurements

We have specified above what happens when one measures a certain property of a physical system at the moment when the state vector of the system is an eigenvector of the operator representing the property - probability = 1 that we get the corresponding eigenvalue.

What if one measures a certain property of a physical system at a moment when the state vector of the system does not happen to be an eigenvector of the operator representing that property (which is most of the time) - what are the probabilities?

We needed a new assumption.

Suppose the system is in the state $|\psi\rangle$, and one carries out a measurement of a property (observable) associated with the operator $\hat{B}$.

We assume the eigenvectors of $\hat{B}$ are the vectors (states) $|b_i\rangle$, which means that $\hat{B}\,|b_i\rangle = b_i\,|b_i\rangle$, i=1,2,3,…. where the b_i are the corresponding eigenvalues.

Quantum theory now assumes that the outcome of measurement is strictly a matter of probability.

Quantum theory stipulates that the probability that the outcome of a measurement of $\hat{B}$ on the state $|\psi\rangle$ (not an eigenvector) will yield the result b_i (remember the only possible results of measurement are the eigenvalues of $\hat{B}$ no matter what state the system is in), is equal to $|\langle b_i | \psi \rangle|^2$ (the **Born rule**).

The probability is given by the absolute square of the corresponding component.

remember that —— component = amplitude

The quantum mechanics formalism based on these postulates + embellishments correctly predicts experimental results for all known experiments (over 100 years).

Some ideas implied by these rules are:

These rules imply that one **cannot** say anything definite about the value of the observable represented by $\hat{B}$ when system is in a state $|\psi\rangle$, which is NOT an eigenvector of $\hat{B}$.

One can only make probability statements.

Before one measures the observable represented by $\hat{B}$ when the system is in a state $|\psi\rangle$, which is NOT an eigenvector of $\hat{B}$, the system **does not** have a value of that observable, according to quantum theory!

Our information about any state is only a set of probabilities.

But all of your experience says that objects have values for measured quantities before they are measured, i.e., your experience tells you that the observable represented by $\hat{B}$ has a value even if we do not measure it.

That is your view (the standard classical view) about what is real and what is not real.

Quantum theory implies you are wrong in both cases!!

Where is the "collapse" postulate?

Since the system has a definite value of the observable represented by $\hat{B}$ after the measurement, i.e., a pointer points to a value or a counter clicks or a mark is registered on a piece of paper (note that these are all **irreversible occurrences** which must be the end result of any measurement) and there is no mechanism to produce a single value in the rules as presented so far, how does it happen?

Most presentations add another rule(as we did earlier) at this point called **collapse of the state vector.**

It is usually proposed that the effect of a measurement is to irreversibly change(collapse) the state vector (which was not an eigenvector of $\hat{B}$) into an eigenvector of $\hat{B}$ (corresponding to the eigenvalue just measured) so that it would be observed to have a definite (probability = 1) value for a subsequent measurement of the operator $\hat{B}$.

This extra rule says that state vector changes(discontinuously) during measurement from representing range of possibilities (superposition of all possible states) to definite state or only one possible outcome.

Which particular eigenvector it gets changed into is determined by outcome of measurement and cannot be known until then!

It cannot be predicted! It is at this point that randomness enters quantum mechanics.

I believe that this last rule is not needed and should not be added.

First of all, no real mechanism is ever given for “how” this process actually takes place, and second no specifications are given as to exactly “when” it occurs.

Also, we don’t need collapse mechanism to make any predictions confirmed by experiments.

That suggests to me that it does not exist! That it is NOT a real process!!

I will now proceed to develop a proposal for “definite outcomes“ **without** using any “collapse” rule.

The path I will follow will not be direct, but it will represent the actual tortuous path I followed to get to the conclusions.

I will meander about, repeating myself often on purpose so I can make subtle changes to the previous ideas as I learned new wrinkles along the way.

I will initially end up with one conclusion, which we will see has flaws and then I will fix the flaws and get what I now think is the correct answer at the end.

The Measurement Process(some repetition of earlier discussions)

We consider a system consisting of a quantum system (Q-system) and a measurement system (M-system).

If the meter, which we assume is initially OFF (state $|0\rangle_M$) was turned ON when quantum system was in $|+\rangle_Q$ state, then according to the above rules the combined system evolves to

$$|+\rangle_Q |0\rangle_M \rightarrow |+\rangle_Q |+1\rangle_M$$ i.e., meter (assuming a good meter) reads +1.

Similarly, if the meter turned ON when the system is in the $|-\rangle_Q$ state, then combined system evolves to

$$|-\rangle_Q |0\rangle_M \rightarrow |-\rangle_Q |-1\rangle_M$$ i.e., meter (assuming a good meter) reads –1.

This indicates that measurement, within framework of our rules, CORRELATES or ENTANGLES the dynamical variables (Q-system) being measured and the macroscopic (M-system) indicator of the meter, which we assume can be directly (macroscopically) observed (and is **irreversible**).

Let us expand(and repeat) this discussion a bit.

We have supposed above that the meter has eigenvectors (labelled by the corresponding eigenvalues), i.e., it is a quantum system also.

$$\begin{aligned} |+\rangle_M &\Rightarrow meter \text{ on}: \ reading \ +1 \\ |-\rangle_M &\Rightarrow meter \text{ on}: \ reading \ -1 \\ |0\rangle_M &\Rightarrow meter \text{ off} \end{aligned}$$

and the quantum system has eigenvectors (labelled by eigenvalues)

$$\begin{aligned} |+\rangle_Q &\Rightarrow value = +1 \\ |-\rangle_Q &\Rightarrow value = -1 \end{aligned}$$

Now suppose that the initial state of the quantum system is a **superposition**

$$|\psi\rangle = a\,|+\rangle_Q + b\,|-\rangle_Q$$

and thus the initial state of the combined system is given by

$$|initial\rangle = \left(a\,|+\rangle_Q + b\,|-\rangle_Q\right)|0\rangle_M$$

which represents the system in a superposition state and meter off.

We are interested in the evolution of this state according to QM.

We note as stated above, if, instead of the above initial state, we started with initial states

$$|A\rangle = |+\rangle_Q\,|0\rangle_M \qquad \textbf{OR} \qquad |B\rangle = |-\rangle_Q\,|0\rangle_M$$

and then turn on the meter, these states must evolve as

$$|A\rangle = |+\rangle_Q\,|0\rangle_M \longrightarrow |A'\rangle = |+\rangle_Q\,|+\rangle_M$$

$$|B\rangle = |-\rangle_Q\,|0\rangle_M \longrightarrow |B'\rangle = |-\rangle_Q\,|-\rangle_M$$

respectively, indicating that the meter measured the appropriate value (definition of good meter) since system is in eigenstate and has a definite value with certainty.

If system is in initial state corresponding to a superposition, however, then the **linearity** of quantum mechanics says must it evolve into

$$|initial\rangle = \left(a\,|+\rangle_Q + b\,|-\rangle_Q\right)|0\rangle_M \longrightarrow |final\rangle = a\,|+\rangle_Q\,|+1\rangle_M + b\,|-\rangle_Q\,|-1\rangle_M$$

We note a problem immediately,

i.e., the meter has not ended up in a state with a definite value

- it remains in a superposition of two macroscopically different pointer readings,

which is **never** observed in the real world.

Hence, if as most physicists assume,

the state vector represents a complete description of the Q-system,

there seems to be a need for the **“collapse”** rule

to fix the result and obtain **“definite”** values!

Since we **will not** be incorporating the “collapse” rule, we must proceed in a **different** way.

We **cannot use** the state vector as the fundamental object in quantum theory

since it **seems** that it must collapse (or maybe not!) for quantum mechanics to work

and we are **not** including “collapse”.

Introduce new mathematical object called the Density Operator to remedy this situation:

Now we present an **alternate** way of representing quantum states. Reasons why come later!

It is still an open question whether this alternative is a **simple** mathematical convenience or a more ontologically **true** representation.

Either way, it has a key role to play in modern interpretations of quantum theory and also leads to a possible solution to measurement problem.

This discussion is more mathematical than rest of these notes, but you will benefit if you persevere and work through material.

It is very important.

The problem, as we will now see, lies with assuming that the state vector is the proper way to represent the Q-system **during** the measurement process.

Now, the expectation value is the **average** of set of measurement results taken from a collection of systems in **same** state.

A straightforward calculation of the expectation value in a specific state takes the following form, with $\hat{O}$ being an operator representing the measurement of the specific physical variable and $|\phi\rangle$ is a state vector of some system in the collection:

$$\left\langle \hat{O} \right\rangle = \langle \phi | \, \hat{O} \, | \phi \rangle$$

If we choose **any set** of basis vectors $\{|i\rangle\}$, i = 1, 2, ... for our vector space, we can expand $|\phi\rangle$ and $\langle\phi|$ as

$$|\phi\rangle = \sum_i a_i \, |i\rangle \quad , \quad \langle \phi | = \sum_j a_j^* \, \langle j |$$

where

$$a_i = \langle i \, | \, \phi \rangle \quad , \quad a_j^* = \langle \phi \, | \, j \rangle$$

Plugging these expansions into the expression for the expectation value:

substitution

rearrangement

$$\langle \hat{O} \rangle = \sum_j a_j^* \langle j | \hat{O} \sum_i a_i | i \rangle = \sum_j \left(\sum_i a_j^* a_i \langle j | \hat{O} | i \rangle \right)$$

$$= \sum_j \left(\sum_i \langle \phi | j \rangle \langle i | \phi \rangle \langle j | \hat{O} | i \rangle \right) = \sum_j \left(\sum_i \left[\langle i | \phi \rangle \langle \phi | j \rangle \langle j | \hat{O} | i \rangle \right] \right)$$

more rearrangement

more rearrangement

Now we define a new operator $\hat{\rho} = |\phi\rangle\langle\phi|$, which, at this point, is just the **projection operator** onto the state $|\phi\rangle$.

We can then write the expectation value as

$$\langle \hat{O} \rangle = \sum_i \left(\sum_j \langle i | \hat{\rho} | j \rangle \langle j | \hat{O} | i \rangle \right)$$

substitution of new definition

Recognizing the relation $\sum_j |j\rangle\langle j| = \hat{I}$ appearing in the expression

one finds that the expectation value can be rewritten in a very interesting form

$$\langle \hat{O} \rangle = \sum_i \left(\sum_j \langle i| \hat{\rho} |j\rangle \langle j| \hat{O} |i\rangle \right) = \sum_i \left(\langle i| \hat{\rho} \left(\sum_j |j\rangle \langle j| \right) \hat{O} |i\rangle \right)$$

$$\langle \hat{O} \rangle = \sum_i \langle i| \hat{\rho} \hat{I} \hat{O} |i\rangle = \sum_i \langle i| \hat{\rho} \hat{O} |i\rangle = \mathrm{Tr}(\hat{\rho}\hat{O})$$

<— the important result!!

sum over diagonal matrix elements

i.e., the expectation value is given by the sum over the **diagonal** matrix elements of the operator product $\hat{\rho}\hat{O}$ (the symbol Tr = Trace is just shorthand for the diagonal sum).

The new operator $\hat{\rho}$ is called the **density operator**.

Why bother to introduce this new operator?

As we will see, the **real power** of density operator approach to QM comes when we have to deal with a situation in which we cannot be sure what state the system is in (just as is the case in the measurement problem!!).

Now, imagine we have a whole collection of identical systems, some in $|\phi_1\rangle$, some in $|\phi_2\rangle$, etc.

We might not know which system in which state, and might not even know how many systems are in any one state.

Example: Think about a beam of electrons that **has not** passed through any Stern-Gerlach (S-G) magnet.

Chances are that the spin states of the electrons are completely random.

Perhaps the best one can know is the probability of finding an electron in any state.

$$P_1 = \mathrm{Prob}(|\phi_1\rangle) \quad , \quad P_2 = \mathrm{Prob}(|\phi_2\rangle) \quad , \quad P_3 = \mathrm{Prob}(|\phi_3\rangle) \quad , \quad .....$$

These probabilities have **nothing to do** with quantum theory.


They simply represent our **ignorance** of the details of what is happening in the beam.

Thus, they are **not related** to any quantum amplitudes.

Given a situation like this, one should still be able to do some useful calculations.

For example, one could work out the expectation value of any measurement as follows.

If one can calculate the expectation value of each individual state, then the overall expectation value is simply given by



$$\left\langle \hat{O} \right\rangle = P_1 \left\langle \phi_1 \right| \hat{O} \left| \phi_1 \right\rangle + P_2 \left\langle \phi_2 \right| \hat{O} \left| \phi_2 \right\rangle + P_3 \left\langle \phi_3 \right| \hat{O} \left| \phi_3 \right\rangle + ... + P_n \left\langle \phi_n \right| \hat{O} \left| \phi_n \right\rangle$$

that is just the standard definition of the average value!!

Think back to original definition of expectation value which just represents the average value of measurement and this will be clear.

What we have done here is put together the weighted average of the average value for each state, **which, as we said, is just the standard definition of the overall average value**.

Now, in this case, if one constructs a density operator that is given by the expression

$$\hat{\rho} = P_1 |\phi_1\rangle \langle\phi_1| + P_2 |\phi_2\rangle \langle\phi_2| + P_3 |\phi_3\rangle \langle\phi_3| + \cdots + P_n |\phi_n\rangle \langle\phi_n|$$

sum of probability x projection operators for all possible states(not just 1 state as earlier)

then the expectation value, in this case, is still given by

$$\langle \hat{O} \rangle = \mathrm{Tr}(\hat{\rho}\hat{O})$$

Proof(for mathematically inclined):

Now
$$\langle\phi_1|\,\hat{\rho}\,|\phi_1\rangle = \langle\phi_1|\,(P_1\,|\phi_1\rangle\,\langle\phi_1| + P_2\,|\phi_2\rangle\,\langle\phi_2| + \cdots)\,|\phi_1\rangle$$

or
$$\langle\phi_1|\,\hat{\rho}\,|\phi_1\rangle = P_1\,\langle\phi_1\,|\,\phi_1\rangle\,\langle\phi_1\,|\,\phi_1\rangle + P_2\,\langle\phi_1\,|\,\phi_2\rangle\,\langle\phi_2\,|\,\phi_1\rangle + \cdots$$

but
$$\langle\phi_i\,|\,\phi_j\rangle = \delta_{ij}$$

so
$$\langle\phi_1|\,\hat{\rho}\,|\phi_1\rangle = P_1$$

similarly
$$\langle\phi_i|\,\hat{\rho}\,|\phi_j\rangle = P_i\delta_{ij}$$

then
$$\mathrm{Tr}(\hat{\rho}\hat{O}) = \sum_i \langle\phi_i|\,(\hat{\rho}\hat{O})\,|\phi_i\rangle = \sum_i \langle\phi_i|\,(\hat{\rho}\hat{I}\hat{O})\,|\phi_i\rangle = \sum_i \langle\phi_i|\,(\hat{\rho}\sum_j(|\phi_j\rangle\,\langle\phi_j|)\hat{O})\,|\phi_i\rangle$$

or
$$\mathrm{Tr}(\hat{\rho}\hat{O}) = \sum_j\sum_i \langle\phi_i|\,\hat{\rho}\,|\phi_j\rangle\,\langle\phi_j|\,\hat{O}\,|\phi_i\rangle$$

Thus
$$\mathrm{Tr}(\hat{\rho}\hat{O}) = \sum_j\sum_i P_i\delta_{ij}\,\langle\phi_j|\,\hat{O}\,|\phi_i\rangle$$

or
$$\mathrm{Tr}(\hat{\rho}\hat{O}) = \sum_i P_i\,\langle\phi_i|\,\hat{O}\,|\phi_i\rangle = \langle\hat{O}\rangle$$

proof complete!

Some notation: when the density operator takes the form $\hat{\rho} = |\phi\rangle\langle\phi|$ only one term!

it is said to represent a **pure state** and when the density operator takes the form

$$P_1 |\phi_1\rangle\langle\phi_1| + P_2 |\phi_2\rangle\langle\phi_2| + P_3 |\phi_3\rangle\langle\phi_3| + \cdots + P_n |\phi_n\rangle$$

like the beam coming out of source in S-G experiment!!

it is said to represent a **mixed state** (more about mixed states later).

A crucial example of the need for the density operator.

Follow this very carefully.

Consider a box containing a very large number of electrons, each having spin = 1/2.

This means that each electron spin can have a measurable component ±1/2 along **any** chosen measurement direction.

Now, suppose the box has a hole so that the electrons can get out and then go into a Stern-Gerlach device oriented to measure the z-components of spin(**arbitrary choice of direction**).

In order to proceed, we need to know how the box of electrons was **prepared**, i.e., what states the electrons are in or what the KET labels are…….

Let us consider **two very different cases**:

In the **first** case, we fill the box with electrons that have been prepared in a superposition state

$$|\psi\rangle_1 = \frac{1}{\sqrt{2}}(|\uparrow_z\rangle + |\downarrow_z\rangle) = |\uparrow_x\rangle$$

This preparation can be done by sending electrons through an x-oriented magnet and choosing one of the resulting beams (UP in x in this case).

Thus, in this case, **each** electron is in the the indicated superposition state of z-directions —-> **each** electron is in a superposition of “up” and “down” in the z-direction while **simultaneously** being in a definite state of spin in the x-direction (up). Hence, the x-value is known and z-value is 50-50

We then fill one box with these electrons.

In the second case, we send electrons through a z-oriented magnet and collect electrons from both beams “z-up” and “z-down” separately.

In this case we **know** that the electrons are either “z-up” OR “z-down”, i.e., they each **have** a definite value - they are **not** in a superposition.

We then fill another box with the two collections of electrons of definite z-spin electrons.
Box is now (50-50) up/down in z-direction, **BUT each electron now has a definite value!**.

Remember, the electrons in the box in this case are EITHER in the state $|\uparrow_z\rangle$ OR in the state $|\downarrow_z\rangle$

Now we proceed with the experiments.

In case (1) we observe “z-up” 50% of the time and “z-down” 50% of the time

and we know that in order to describe this system by a state vector we must say

$$|\psi\rangle_{BOX(1)} = \frac{1}{\sqrt{2}}(|\uparrow_z\rangle + |\downarrow_z\rangle) \rightarrow \text{50-50 up-down}$$

i.e., every electron in the box is in this superposition state. which is how we created them

In case (2)) we also observe “z-up” 50% of the time and “z-down” 50% of the time.

But we now have a problem because we are using state vectors as the fundamental object in the theory!

If I did not already know that each electron in the box in this case had a definite value, I would be tempted to describe this system by the same state vector as in case (1).

However, we know that is not true!

The electrons in case (2) are not each in a superposition — they all have **definite values**!

So, if I only measure z-components I cannot tell whether I have case (1) or case (2) and I **do not know how to write the state vector** for the box in case (2),

But remember how I created the electrons in case (1).

They all have a **definite** value of the x-component, namely, “x-up’.

So if I subject the electrons coming out of the box in case (1)

to an x-measurement instead of a z-measurement,

I will end up with only **one beam**!

However, in case (2), the electrons coming out of the box are either “z-up” OR “z-down”

each of which is 50-50 in the x-direction and thus I would end up with two beams after the extra x-measurement!

I get different results for 2nd measurement because they are different states!!

The different results mean that their states must be described differently in QM.

State vectors do not give us the freedom to do this unless we want to monkey around with relative phases between components, i.e., mathematically, we would need to write

$$|\psi\rangle_{BOX(1)} = \frac{1}{\sqrt{2}}(|\uparrow_z\rangle + |\downarrow_z\rangle) \quad , \quad |\psi\rangle_{BOX(2)} = \frac{1}{\sqrt{2}}(|\uparrow_z\rangle + e^{i\alpha}|\downarrow_z\rangle)$$

where α is a **completely unknown relative phase factor**, which must be averaged over during any calculations since it is different (and random) for each separate measurement (each member of ensemble).

Thus, we would be choosing to allow the two vectors to differ in the relative phase of their components and we are forced to say this is meaningful!

Let us continue…

If we use density matrices, we have a very **different story**.

For the pure state a density operator (or matrix) is defined as

$$\hat{\rho} = |\psi\rangle \langle\psi|$$

for some state vector $|\psi\rangle$, i.e., it is the pure state projection operator.

In case (1) this gives

$$\hat{\rho} = |\psi\rangle \langle\psi| = \frac{1}{2}(|1/2\rangle + |-1/2\rangle)(\langle 1/2| + \langle -1/2|)$$

or

$$\hat{\rho} = \frac{1}{2}(|1/2\rangle \langle 1/2| + |-1/2\rangle \langle 1/2| + |1/2\rangle \langle -1/2| + |-1/2\rangle \langle -1/2|)$$

—> 4 terms

projection operator cross-term cross-term projection operator

Derivation of the $\hat{\rho}$ matrix in the (+1/2, –1/2) basis: from our earlier definitions we have

$$\rho = \begin{pmatrix} \langle 1/2|\,\hat{\rho}\,|1/2\rangle & \langle 1/2|\,\hat{\rho}\,|-1/2\rangle \\ \langle -1/2|\,\hat{\rho}\,|1/2\rangle & \langle -1/2|\,\hat{\rho}\,|-1/2\rangle \end{pmatrix}$$

Now

$$\langle 1/2|\,\hat{\rho}\,|1/2\rangle$$

$$= \langle 1/2|\,\frac{1}{2}(|1/2\rangle\langle 1/2| + |-1/2\rangle\langle 1/2| + |1/2\rangle\langle -1/2| + |-1/2\rangle\langle -1/2|)\,|1/2\rangle$$

$$= \frac{1}{2}$$

and similarly for the other matrix elements and we get

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

where the diagonal matrix elements represent probabilities(50-50) and the non-zero off-diagonal matrix elements indicate that one **will observe** quantum interference effects in this system.

It is clear(look at above expression) that any pure state density operator **cannot** be written as the sum of pure state projection operators (multiplying always generates cross-terms).

In case (2), however, each electron is in a definite state (has a definite value) so that each state is separately represented by a single projection operator

—> only have the sum of two definite terms

$$\hat{\rho} = \frac{1}{2}(|1/2\rangle\langle 1/2| + |-1/2\rangle\langle -1/2|)$$

we call this form the "ignorance" density operator

and

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

same algebra

which clearly **is** the sum of pure state projection operators.

This corresponds to what is called a **mixed state**.

Note that the off-diagonals are zero so that this density operator **cannot** lead to any quantum interference effects. The electron **have** values like classical particles!!

Remember for later, this system(case (2)) is such that electrons have values so that the density operator takes this form (sum of projection operators for each value —> NO cross-terms)!!

Note that when electrons DO NOT HAVE VALUES (case (1)) the density operator has interference terms(mixed terms) and cannot be written as a sum of projection operators

A DIFFERENCE that does NOT show up(without extra phases) when using state vectors!!

We note that if we treat case (2) as pure state with the extra relative phase factor we would obtain:

$$\hat{\rho} = \frac{1}{2}\begin{pmatrix} 1 & e^{i\alpha} \\ e^{-i\alpha} & 1 \end{pmatrix} \qquad \text{which becomes} \qquad \hat{\rho} = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

when we average over α .

For comparison, let us now digress to see what happens in a **real classical system**.

Consider rolling a standard die which has possible values = 1,2,3,4,5,6 where the probability of occurrence of each value = 1/6.

In this case, the density operator representing the die will be for a mixed state (no interference effects) and the die **has** a value before/after each roll so that we have

$$\rho = \frac{1}{6}(|1\rangle\langle 1| + |2\rangle\langle 2| + |3\rangle\langle 3| + |4\rangle\langle 4| + |5\rangle\langle 5| + |6\rangle\langle 6|)$$ **a mixed state**

and the expectation value of the $\widehat{\mathrm{ROLL}}$ operator must be

$$\langle \widehat{ROLL} \rangle = \frac{1}{6}(1+2+3+4+5+6) = 3.5$$ **standard definition**

just sum of values times probabilities!!

More formally, we know that any operator can be written as sum of eigenvalues × projections operators, i.e., for the $\hat{B}$ operator introduced earlier we found we can write

$$\hat{B} = \sum_i b_i |b_i\rangle\langle b_i|$$

so that we have for the $\widehat{\mathrm{ROLL}}$ operator

$$\widehat{ROLL} = 1\,|1\rangle\langle 1| + 2\,|2\rangle\langle 2| + 3\,|3\rangle\langle 3| + 4\,|4\rangle\langle 4| + 5\,|5\rangle\langle 5| + 6\,|6\rangle\langle 6|$$

The operator product of the density operator and the $\widehat{\mathrm{ROLL}}$ operator can be written(using the orthonormality of the basis state vectors) as

$$\hat{\rho}\,\widehat{ROLL} = \frac{1}{6}(1\,|1\rangle\langle 1| + 2\,|2\rangle\langle 2| + 3\,|3\rangle\langle 3| + 4\,|4\rangle\langle 4| + 5\,|5\rangle\langle 5| + 6\,|6\rangle\langle 6|)$$

Thus, the expectation or average value is

$$\langle\widehat{ROLL}\rangle = \mathrm{Tr}(\hat{\rho}\widehat{ROLL}) = \sum_k \langle k|\,\hat{\rho}\widehat{ROLL}\,|k\rangle = \frac{1}{6}(1+2+3+4+5+6) = 3.5$$

as we obtained earlier. I can do all detailed algebra if interested after class session.

But, in this case, **we know the values are real before the measurement**,

i.e., we are using a macroscopic die with numbers we can see!

Thus, it **seems** that this **particular** form of a density operator **represents that case**.

The same was true earlier for the electrons in the box when we knew they were either “up” or “down”!

We also had a density operator of the same form!

In both cases the system **had known values before measurements**!

REMEMBER this fact for later.

If we were to add the “collapse” rule it raises a host of questions:

What exactly do we mean, physically and mathematically,

by a “collapse during measurement” of quantum system?

Does collapse occur all at one instant?

What if the state occupies a finite volume?

Wouldn’t instantaneous collapse(the entire volume changes at once) contradict special relativity?

If, instead, collapse occurs during a time interval, then what equation describes its time-evolution during that interval?

Quantum states are presumed to follow the Schrödinger equation,

which prescribes continuous time evolution.

How can instantaneous state collapse be reconciled with smooth evolution?

How can we resolve “problem of outcomes” that appears to arise

when a superposed quantum’s state is measured by “which-state” detector,

creating a so-called entangled state of the quantum and the detector

that appears to be an indefinite superposition

of two macroscopically distinct states of a composite system?

Such questions and more comprise the **quantum measurement problem**.

There are many alternative interpretations of the quantum physics mathematical formalism,

and several alternative modifications of theory,

have been proposed to resolve problem, with no consensus on a solution.

It is remarkable that, despite the unparalleled

experimental success of quantum theory across a vast range of experiments,

most of the suggested solutions **differ** from standard quantum physics

in one or more significant aspects.

Most involve **new** interpretations of the standard mathematical formalism.

Interpretations such as "human minds collapse the quantum state"

or "all possible collapses occur but only one of them occurs in our particular universe"

or even rejection of physical reality of quantum world

and assumption that quantum probabilities

(and hence changes in those probabilities,

such as quantum state collapse) are mere measures of personal degrees of belief.

yadda, yadda, yadda……

Other suggestions assume modifications of the standard mathematical formalism,

such as an additional mechanism

that causes quantum states to spontaneously collapse from time to time,

or new "hidden" and hence uncontrollable variables

that create the illusion of quantum randomness. **more yadda, yadda …..**

With all these thoughts/ideas in hand,

we will now continue this discussion of the measurement problem

and finally suggest a resolution of problem of definite outcomes

that lies **entirely** within the framework of standard quantum physics.

As I said, we will follow a my tortuous path.

First, we will spend time with one possible solution to the measurement problem.

We will arrive at a point where it "looks" like we found a solution.

But it will not be so!

The reasons will be subtle but clear.

We will then see how to correctly interpret what is happening and find in the end that von Neumann was correct in 1932 even though for 90 years papers were published deriding his ideas!

Begin with question of definitions.

What do we mean, physically, by **quantum measurement**?

First, we need a non-anthropomorphic definition of the concept known as "measurement,"
i.e., there were measurements before humans,
so let's broaden its physical definition as follows:

A "quantum measurement" means
any quantum process that results in a macroscopic effect,
regardless of whether humans or laboratories are involved.

Thus not only is an electron striking a laboratory viewing screen

and creating a visible flash a measurement, but also a cosmic-ray muon striking

and macroscopically moving a sand grain on a planet in some other galaxy

is a measurement.

To analyze a measurement,

we look at a specific experiment: suppose an electron beam passes

through a pair of double slits and then impacts a viewing screen.

Just as in Thomas Young's similar double-slit experiment using light, performed in 1801,

a pattern is formed on the viewing screen that **seems** to shows interference

between the two portions of the electron beam which are **seemingly**(classically)

coming simultaneously through the two slits:

a broad dark-and-bright striped pattern spreads out widely on the screen

- much wider than slits
- indicating regions of destructive (dark) and constructive (bright) interference.

or maybe I should say …

regions of increased electron intensity and decreased electron intensity(including 0)

On closer inspection,

the bright lines are formed by a very large number of tiny individual electron impacts,

each one making a small flash on the screen.

According to our definition, each flash is a measurement

of the position of an electron as hits the screen.

I repeat - Each electron's flash on screen is a measurement!

For the purposes of this analysis, however,

it is more informative to consider a related example of measurement,

still based on the double-slit experiment.

Suppose an electron detector is installed at the slits

and assume that the detector can detect the electron's position

as it passes through slits while disturbing each electron only **minimally** .

As it turns out, measurement,

even by such a minimally-disturbing "which-path detector", changes everything.

Exactly when the detector turns on,

the pattern of impacts on screen changes from a striped interference pattern

to a smoothly-spread-out sum of two single-slit patterns,

each showing diffraction but no interference, i.e., the probability function governing

the pattern of new flashes changes after that instant!

The interference pattern abruptly vanishes, i.e., probability has changed.

An analogous experiment has been done using light (photons) instead of electrons,
and using an interferometer rather than double-slit interference setup as we discussed earlier.

A which-path detector was randomly switched on or off
as each photon passed through this experiment;

Photons for which the detector was "off" formed an interference pattern
while photons for which the detector was "on"
formed the expected no-interference pattern.

Reminder:

If the system is in a “pure” superposition state, then the density operator takes the form

$$\hat{\rho} = \frac{1}{2}(|1/2\rangle\langle 1/2| + |-1/2\rangle\langle 1/2| + |1/2\rangle\langle -1/2| + |-1/2\rangle\langle -1/2|)$$

i.e., sum of projection operators and “cross-terms” (interference terms)

or the matrix form

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

where the diagonal matrix elements represent probabilities

and the off-diagonal matrix elements imply

that one will observe quantum interference effects in this system.

Clearly, any pure state density operator cannot be written

as the sum of pure state projection operators (because of the cross-terms).

If in a “mixed state”, then the density operator takes the form

$$\hat{\rho} = \frac{1}{2}(|1/2\rangle\langle 1/2| + |-1/2\rangle\langle -1/2|)$$

or the matrix form

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

i.e., sum of projection operators (no interference terms).

As we found in the dice example:

$$\hat{\rho} = \frac{1}{6}(|1\rangle\langle 1| + |2\rangle\langle 2| + |3\rangle\langle 3| + |4\rangle\langle 4| + |5\rangle\langle 5| + |6\rangle\langle 6|) \quad \text{mixed state}$$

We now state our first attempt at describing what happens during a measurement.

When we observe a quantum system evolve into such a “mixed state” density operator,

then the quantum system can be interpreted “classically”,

i.e., a measurement has taken place

and it has been **irreversibly recorded somewhere**.

That is the important point!!

A full “collapse” is not necessary!!

We only need to arrive at (evolve to) a particular form of the density operator!!

Before continuing let me quote Richard Feynman about what has happened.

"Somewhere in the measuring apparatus changes have occurred and the concept of a quantum amplitude is not valid anymore, i.e., if we are throwing our quantum die, then the quantum amplitude only remains a valid idea until the die comes to rest on the floor.

And that point you can't do an experiment which distinguishes interfering alternatives from just plain odds (like with classical dice).

—> we are at the "ignorance" density operator stage!

There is no point where you can say a "reduction" has taken place, i.e., there is just a murky unseen phasing out of the quantum amplitudes.

Possibly what is happening is that in the macroscopic measuring device, quantum phase information gets so "smeared out" that it is no longer definite and thus the system behaves classically.

It registered an irreversible result!

Since we are not quantum objects we may never be able to see the details of this process, but it clearly happens."

IMPORTANT!!

Now let us continue our discussion!

Possible ways to think about getting to the irreversible recording point must be found.

Two digressions —> possible ways to get to the irreversible recording —

(1) The so-called Gambler's Ruin problem (from the mathematicians)

We start by investigating the **Mathematical Problem of the Points** (seems unrelated but it is not!)

A sequence of fair coins is flipped.

Player A gets a point for every head and player B gets a point for every tail.

Player A wins if there are a heads before b tails, otherwise B wins.

Find the probability that A wins.

Let α(a,b) be the probability that A wins and

β(a,b) the probability that B wins —> $\alpha(a,b) + \beta(a,b) = 1$ someone wins!

Solution (due to Pascal and Huygens -1500's)

$$\alpha(a,b) = \frac{1}{2^{a+b-1}} \sum_{k=0}^{b-1} \binom{a+b-1}{k} \quad \text{where} \quad \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

binomial coefficient

$$k! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot k$$

An example is the **Gambler's Ruin problem**.

Just as in the problem of points, suppose that at some stage A has a counters,

and B has m + n − a counters so that total number of counters = m+n,

and let A's chances of victory at that point be v(a).

The solution when A starts with m counters is given by

$$p_A = v(m) = \frac{1 - (\beta/\alpha)^m}{1 - (\beta/\alpha)^{m+n}}$$

This works for all cases except $\alpha = \beta = 1/2$ (the 50-50 probability case).

For that special case, the solution is

$$v(a) = \frac{a}{m+n}$$

This implies that there is always a winner (hence the the name **Gambler's Ruin** for the loser)

NOTE: Even when $\alpha = \beta = 1/2$ there is still always a winner!!!!

Also note that this relation is **linear** in **a** in this special case!

Pearle's Theorem

What does this have to do with quantum collapse?

Here is a short note from Philip Pearle (a really good physicist and friend): I quote:

I soon found a charming **analogy** for collapse dynamics,

useful for providing an intuitive and non-technical **explanation of how it works**.

I happened to be browsing in Feller's book on probability

(a favorite textbook, from an undergraduate course taught by Stanislaus Ulam)

when I encountered the **gambler's ruin game**.

Two gamblers, initially possessing, respectively, a **fraction** $x_1(0)$, $x_2(0)$ of their combined wealth

(so $x_1(0) + x_2(0) = 1$)

repeatedly toss a fair coin, and the result, heads or tails,

determines which one gives one dollar to the other.

They play until one gambler loses all his money, and the game ends.

The **analogy** arises if the amount of money possessed by one gambler at any time

is proportional to the squared amplitude of one of two states

whose sum is the state vector representing the physical system undergoing collapse.

Just as one gambler loses all his money, **so one of the states loses all its amplitude**,

and as the other gambler wins all the money,

so the state vector ends up as totally described by the other state (—> collapse!!).

collapse has occurred here without any interference by gamblers!

Let Q(x) be the conditional probability that a gambler wins the game,

given that he has the fraction x of the total wealth.

If Δ is the fraction of the total wealth they exchange at each toss (i.e., Δ = \$1/total dollars),

the difference equation

$$Q(x) = \frac{1}{2}Q(x - \Delta) + \frac{1}{2}Q(x + \Delta)$$

expresses that there are two routes to win if one has fractional wealth x,

namely lose the next toss and drop to x − Δ but win thereafter,

or win the next toss and rise to x + Δ and win thereafter.

The solution of the difference equation is $Q(x) = Ax + B$, where A and B are constants.

Boundary conditions:

Since $Q(0) = 0$ (because you can't win if you have no money) and $Q(1) = 1$

(because you have won if you have all the money), then $Q(x) = x$, i.e.,

$Q(1) = 1 = A+B$ $\quad Q(0) = B= 0 \quad \text{—>} A = 1, B = 0 \quad \text{—>} Q(x) = x$

That is, if one starts with the fraction $x = x(0)$ of the money,

one has the probability $x(0)$ of attaining all the money —> you get to $x = 1$,

which is exactly collapse behavior!!

The game can be modified to have many players,

to have Δ change as the game progresses

(e.g., to get smaller as one gambler gets closer to losing), etc.

So, one may think of quantum collapse

(getting to the point where an irreversible measurement has been made)

as a gambler's ruin competition among the states in a superposition,

to see which final state wins the game (remains at the end).


I ran a simulation of the Gambler's Ruin problem using OCTAVE on a computer.


OCTAVE Code


gambruin.m


```
x1=[];
y1=[];
x=40;x1=[x1,x];
y=100-x;y1=[y1,y];
count=1;
while (x ~= 0) && (x ~= 100)
r=rand;
x=x+(r <= 0.5)-(r > 0.5);x1=[x1,x];
y=100-x;y1=[y1,y];
count=count+1;
endwhile
figure
plot(1:count,x1,'-b');
axis([1,count,0,100]);
xlabel('Step');
ylabel('Value');
title('Gambler''s Ruin')
hold on;
plot(1:count,y1,'-r');
hold off
count
```

Some sample runs are shown in next slide - **note there is always a winner(loser)**


Gambler's Ruin
Value
Step
0
10
20
30
40
50
60
70
80
90
100
200
400
600
800
1000
1200


Gambler's Ruin
Value
Step
0
10
20
30
40
50
60
70
80
90
100
100
200
300
400
500
600
700
800


Gambler's Ruin
Value
Step
0
10
20
30
40
50
60
70
80
90
100
1000
2000
3000
4000
5000
6000
7000


Gambler's Ruin
Value
Step
0
10
20
30
40
50
60
70
80
90
100
200
400
600
800
1000
1200

One state always wins out!!

Collapse occurs even though there is no direct collapse mechanism!!

An irreversible mark always appears!!

Could this be the way collapse works without a postulate or mechanism?

More Details are in supplementary reading GamblerRuin.pdf

Another way to get to the irreversible recording ……

(2) Decoherence (from the physicists)

If we have a quantum system is the state

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

then this(non-zero off-diagonal elements) states **exhibits** quantum interference effects.

“Decoherence” says that as a state interacts with macroworld in its environment (which it has to do) the off diagonal elements go to zero, i.e., the state makes transition to

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

As we already said, this state has zero quantum interference effects and a definite measurement value has been irreversibly recorded somewhere.

If we had quantum dice the state vector would be

$$|Dice\rangle = \frac{1}{\sqrt{6}}(|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle + |6\rangle) \quad \text{and} \quad \rho = |Dice\rangle\langle Dice| \quad \textbf{giving}$$

$$\begin{aligned}\rho = \frac{1}{6}\,[&|1\rangle\langle 1| + |2\rangle\langle 1| + |3\rangle\langle 1| + |4\rangle\langle 1| + |5\rangle\langle 1| + |6\rangle\langle 1| \\ &+ |1\rangle\langle 2| + |2\rangle\langle 2| + |3\rangle\langle 2| + |4\rangle\langle 2| + |5\rangle\langle 2| + |6\rangle\langle 2| \\ &+ |1\rangle\langle 3| + |2\rangle\langle 3| + |3\rangle\langle 3| + |4\rangle\langle 3| + |5\rangle\langle 3| + |6\rangle\langle 3| \\ &+ |1\rangle\langle 4| + |2\rangle\langle 4| + |3\rangle\langle 4| + |4\rangle\langle 4| + |5\rangle\langle 4| + |6\rangle\langle 4| \\ &+ |1\rangle\langle 5| + |2\rangle\langle 5| + |3\rangle\langle 5| + |4\rangle\langle 5| + |5\rangle\langle 5| + |6\rangle\langle 5| \\ &+ |1\rangle\langle 6| + |2\rangle\langle 6| + |3\rangle\langle 6| + |4\rangle\langle 6| + |5\rangle\langle 6| + |6\rangle\langle 6|]\end{aligned}$$

Thus, the quantum dice generally has a lot of quantum interference terms.

This corresponds to a density matrix of the form

$$\rho = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \\ a_{51} & a_{52} & a_{53} & a_{54} & a_{55} & a_{56} \\ a_{61} & a_{62} & a_{63} & a_{64} & a_{65} & a_{66} \end{pmatrix}$$

lots of off-diagonal terms

lots of quantum interference

Rolling the dice (a measurement) or “decoherence” if dice left alone, produces

$$\rho = \begin{pmatrix} a_{11} & 0 & 0 & 0 & 0 & 0 \\ 0 & a_{22} & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & a_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 & a_{55} & 0 \\ 0 & 0 & 0 & 0 & 0 & a_{66} \end{pmatrix}$$

where $a_{11} = a_{22} = a_{33} = a_{44} = a_{55} = a_{66} = \frac{1}{6}$

or

$$\hat{\rho} = \frac{1}{6}(|1\rangle\langle 1| + |2\rangle\langle 2| + |3\rangle\langle 3| + |4\rangle\langle 4| + |5\rangle\langle 5| + |6\rangle\langle 6|) \quad \text{mixed state}$$

and as we said earlier this indicates a measurement has occurred!! .

Maybe that is what Gambler's Ruin is doing and they are the same thing!

Now let us continue on our earlier path.

Now look closely at "which-path" experiments

In so-called "delayed-choice experiment" mentioned earlier,

there is instantaneous (to within some accuracy) fast switching between two states;

Also, each collapse is executed entirely while the photon was inside interferometer.

We can gain considerable insight by studying

how quantum theory describes a which-path measurement.

Note: it is a measurement as defined earlier,

because a detector **registers** "slit 1" or "slit 2" macroscopically for each electron.

Denote state of one electron passing through slit 1 as $|\psi_1\rangle$

and the state of one electron passing through slit 2 as $|\psi_2\rangle$.

John von Neumann, who was the first to

carefully analyze measurement in purely quantum-theoretical terms,

insisted on treating not only the measured quantum but also the macroscopic detector as quantum systems because, after all, detectors are made of atoms and they perform a quantum function by detecting individual quanta.

I agree with that view of the macroscopic detector.

Now we repeat our earlier discussion again, filling in any remaining gaps and eliminating any remaining confusions.

Accordingly, represent "ready to detect" quantum state of detector by $|ready\rangle$,

and state of detector after detecting an electron by $|1\rangle$ if $|\psi_1\rangle$ was detected,

and by $|2\rangle$ if $|\psi_2\rangle$ was detected.

A properly operating detector will surely transition from |ready⟩ to $|1\rangle$

upon measurement of electron that has been prepared

(perhaps by simply shutting slit 2) in $|\psi_1\rangle$ state .

As a limiting idealization,

we assume, with von Neumann,

that the measurement of electron prepared in the state $|\psi_1\rangle$

leaves the electron still in state $|\psi_1\rangle$ after detection.

Such a **minimally-disturbing measurement** would cause

the electron-plus-detector composite system,

initially in the composite state $|\psi_1\rangle\,|ready\rangle$,

to transition into the final state $|\psi_1\rangle\,|1\rangle$.

We summarize the process as (I will be referring to the numbered equations)

$$|\psi_1\rangle\,|ready\rangle \rightarrow |\psi_1\rangle\,|1\rangle \qquad \textbf{(1)}$$

You should make a copy of all the numbered equations for reference as I lecture

Similarly, a minimally-disturbing measurement of an electron initially prepared in $|\psi_2\rangle$ is described mathematically by

$$|\psi_2\rangle\,|ready\rangle \rightarrow |\psi_2\rangle\,|2\rangle \qquad \textbf{(2)}$$

Now suppose that both slits are open so each electron “can pass through either slit”,
and suppose the preparation and experiment (e.g. slit widths) is symmetric
with respect to the two slits.

Then the state of each electron as it approaches the slits prior to detection
must be described by a symmetric superposition

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|\psi_1\rangle + |\psi_2\rangle) \qquad \textbf{(3)}$$

But quantum physics, including its time dependence, is **linear**.

This implies that $|\psi\rangle\,|ready\rangle$ evolves according to

$$\begin{aligned} |\psi\rangle\,|ready\rangle &= \frac{1}{\sqrt{2}}(|\psi_1\rangle + |\psi_2\rangle)\,|ready\rangle \\ &= \frac{1}{\sqrt{2}}(|\psi_1\rangle\,|ready\rangle + |\psi_2\rangle\,|ready\rangle) \\ &\rightarrow \frac{1}{\sqrt{2}}(|\psi_1\rangle\,|1\rangle + |\psi_2\rangle\,|2\rangle) \end{aligned} \qquad \textbf{(4)}$$

The final state


$$|\Psi\rangle = \frac{1}{\sqrt{2}}(|\psi_1\rangle\,|1\rangle + |\psi_2\rangle\,|2\rangle) \qquad \textbf{(5)}$$

following detection is said to be **"entangled"** the state properties are now “correlated”

because it **cannot be factored** into simple product of states of the two sub-systems.

You should make a copy of all the numbered equations for reference as I lecture

As indicated suggestively in the Figure below:



when two independent quanta pass **near** each other, interact, and subsequently separate, the interaction generally **entangles** the two quanta and the entanglement then **persists** after interaction **regardless** of how far apart the two quanta might eventually travel, provided only that the two quanta experience **no further interactions**.

Despite a possibly wide spatial separation, entangled quanta have a **unity** not possessed by non-entangled quanta.

It is this unity that is the source of quantum non-locality!

Entanglement is ubiquitous in nature.


The entangled "measurement state" (5) that is at **heart** of quantum measurement is remarkably **subtle**.

To fully understand **“entanglement”**, we **first** need to understand **“superposition”**.

The quantum principle says that

> **any** linear combination(= superposition) of possible quantum states of a system,
> as in (3) and (5) for example, is also a possible quantum state of that system.

Figure on next slide pictures an experiment that demonstrates such a superposition of states.



This represents a layout of optical paths called a “Mach-Zehnder interferometer”.

Light beam enters at lower left passing through "beam splitter" BS1;

reflected beam(amplitude) makes a right angle with incoming direction


while transmitted beam(amplitude) passes straight through.

So the beam “splits” and each “half” traverses one of two paths (meaning amplitudes exist for such to happen);

Mirrors M bring paths back to crossing point as shown.

Devices called "phase shifters,

" denoted by ϕ_1 and ϕ_2, are placed into each path



The phase shifter can add

a short variable length to a path.

A second beam splitter BS2 can be placed at the final crossing point.

Without BS2, each “half”-beam(amplitude) moves straight ahead along one path

to the detector on that path.

Things get more interesting with BS2 in place.

Because 50% of each of the two beams(amplitudes) then goes to each detector,

BS2 mixes the two beams(amplitudes) together so the experiment can show interference.

The interferometer is constructed so that, when phase shifters are set to zero,

two "optical paths" (number of wavelengths, after accounting for phase changes upon reflection and refraction) from the entry point to D1 are equal


while the two optical paths to D2 differ by half a wavelength.

Thus, is found that amplitudes interfere constructively at D1 and destructively at D2,

i.e., all photons go to D1.

If one then uses ϕ_1 or ϕ_2 to add half wavelength to either path, amplitudes then interfere constructively at D2 and destructively at D1,

i.e., all photons go to D2.



As one continuously varies length of one or other path

by varying one or the other phase shifter,

one finds that amount of light(number of photons) arriving at D1 varies continuously from

100% down to 0%,

while amount arriving at D2 varies from 0% to 100%.

Two paths are clearly interfering -> can construct a normal interference pattern from the data!

Experiment is an interferometer-based analog of Young's double-slit interference experiment demonstrating the "wave" nature of light - because we have set the **experiment context** to do so!.


But, as we know, light is really just photons, and photons are indivisible.

How then does **nature** explain this experiment

when we dim the light source to a level where only **one photon at a time** traverses the

interferometer? i.e., when the detectors only record one photon at any instant of time!

After all, the photon still traverses BS1, but it cannot split in two because a **quantum is unified and cannot be split**.



With BS2 removed, one finds either D1 or D2 registers a single entire photon, randomly, with 50-50 probabilities, regardless of how the phase shifters are set.

The randomness is **absolute** - it is more random than any human macroscopic game, such as coin flips, which only mimics randomness.

Nature **has invented** quantum randomness to deal with obstacles such as beam splitters while **preserving the unity of the quantum**.

Detectors **never** register half a photon.

They **either** get either a whole photon or no photon.

What happens in single-photon experiment with BS2 present?

As discussed earlier,

beginning from equal path lengths,

which give constructive interference at D1


and destructive interference at D2,

as phase shifters vary,

probabilities of detecting a photon at D1 and D2

vary as in Figure,

which gives the percentage of photons impacting D1.



Importantly,

results do not depend on which phase shifter the experimenter chooses to vary.

Since each photon responds to changes in either path length,


each photon **must "follow both paths" - whatever that statement in words means**!

This verifies the superposition principle and shows that quanta can **seem** to be in **"two places at same time".**

This seems **paradoxical** if we **assume** photons are tiny particles,

but if we assume photons are waves it is **not** paradoxical,

i.e, each photon simply spreads along both paths, interfering with itself at D1 and D2.



One must conclude that each photon travels both paths

even when BS2 is not present to directly verify this,

because once a photon enters interferometer

it **must behave in same manner** regardless of whether BS2 is placed


or not placed at far end.

A **delayed-choice** experiment provides further evidence for this conclusion:

Since photons **"do not know"** whether BS2 will be inserted,

they must travel both paths on all trials

including those for which BS2 is not inserted - **this is connected with entanglement**.



With BS2 removed,

the situation is like the double-slit experiment

with a which-slit detector present.

Each photon is entangled

with macroscopic detectors D1 or D2 as in (5).

With BS2 present, the two paths mix and we have a situation

like double-slit experiment with no which-slit detector -

each photon follows two paths to each detector

where it interferes with itself, and we detect an interference state (3).

All of this clearly suggests that measurements affect superposed quantum states via entanglement of the superposed quantum with a detector.

Resolving Paradoxes and Understanding Measurement

The apparent paradox of Schrödinger's cat

A cat is penned up in a steel chamber, along with the following setup.

In a Geiger counter there is tiny bit of radioactive substance, so small, that in course of an hour one of its atoms decays, but also, with equal probability, none decay;

if it happens, the counter tube discharges and through a relay releases a hammer which shatters a small flask of hydrocyanic acid (poisonous vapor).

If one left the entire system to itself for an hour,

one would say that the cat still lives if no atom has decayed!

If one uses a state vector approach for understanding the entire system,

one would express this by having a living and a dead cat mixed

or smeared out in equal parts -

a superposition of "dead" and "alive" with other properties of system!

a statement that is as stupid as it sounds!!

It is typical of these cases,

that indeterminacy originally restricted to atomic domain

becomes transformed into a macroscopic indeterminacy,

which can only then be resolved by direct observation,

which prevents us from naively

accepting as valid the “blurred model" for representing reality.

Mathematically, the nucleus and the cat

have become entangled in measurement state (5),

with $|\psi_1\rangle$ and $|\psi_2\rangle$ representing the undecayed and decayed states of the nucleus

and $|1\rangle$ and $|2\rangle$ representing the alive and dead cat.

According to Schrodinger’s “understanding” of this situation,

the indeterminacy of the microscopic nuclear state "becomes transformed

into the macroscopic indeterminacy" of cat, and he could not comfortably accept this

“blurred cat state” i.e., a cat that is in a superposition of being both alive and dead.

He hoped this would say something is wrong with QM.

As we will show, standard quantum physics says that Schrodinger's 1937 understanding was **incorrect**.

The composite system (cat-plus-nucleus) is not predicted to be

in superposition of two states of a cat and two states of a nucleus.

Instead, the composite system is predicted to be in a superposition of two **correlations** between the cat and the nucleus; one where live cat is 100% correlated with undecayed nucleus, and second where dead cat is 100% correlated with decayed nucleus.

Entanglement will have transformed a pure state superposition of nuclear states to a pure state superposition of correlations between subsystem states.

We will see this is what one expects from quantum mechanics, and it is **not paradoxical**.

This so-called “problem of definite outcomes”

applies of course to more than Schrodinger’s dramatized example.

Regardless of whether measuring instrument is a which-slit detector,

a Geiger counter, or a cat, the entangled state (5) applies.

This state appears at first glance to represent a quantum superposition

in which the detector is in two macroscopically different states simultaneously.

If so, then there is an inconsistency within quantum physics,

because obviously it **cannot** be this easy to create a **macroscopic superposition**.

The question is: Is it true that (5) really represents a macroscopic superposition?

Turns out, there is more to this entangled state than meets the eye.

If one assumes the detector to be in a superposed state in its own space $a|1\rangle + b|2\rangle$,

one finds that (5) necessitates either a = 0 or b = 0 (one or the other), implying that the detector is not in an individually superposed state **within its own Hilbert space**.

The **same** applies to the detected quantum:

It is not in a superposed state $a|\psi_1\rangle + b|\psi_2\rangle$ with both $a \neq 0$ and $b \neq 0$.

The entanglement process leaves neither sub-system superposed in **own** space!

So far as I know, this simple fact has long been mostly ignored by physicists studying the measurement problem - except for a few that I work with.

The Density Operator to the Rescue

The density operator formalism for quantum physics provides a stronger version of this conclusion(because it is the more appropriate approach).

Density operator for quantum system whose state is $|\psi\rangle$(pure state) is a projection operator

$$\hat{\rho} = |\psi\rangle \langle\psi| \qquad \textbf{(6)}$$

As we saw earlier, if system is in a state whose density operator is $\hat{\rho}$, then standard quantum expectation value $\langle \hat{O} \rangle$ of an arbitrary observable $\hat{O}$ is found from

$$\langle \hat{O} \rangle = \mathrm{Tr}(\hat{\rho}\hat{O}) \qquad \textbf{(7)}$$

where "Tr" represents trace operation (sum of diagonal matrix elements).

This approach is especially useful if the quantum system is a composite of two subsystems A and B.

We define the density operator $\hat{\rho}_A$ for subsystem A alone by

$$\hat{\rho}_A = \mathrm{Tr}_B(\hat{\rho}) \qquad \textbf{(8)}$$

where "Tr_B" means that trace taken **only** over states of subsystem B (remove all traces of B).

It is then easy to show (see later)

that standard quantum expectation values for subsystem A alone

(values obtained by an observer of A **without** any knowledge of B) are

$$\langle \hat{O}_A \rangle = \mathrm{Tr}(\hat{\rho}_A \hat{O}_A) \qquad \textbf{(9)}$$

where $\hat{O}_A$ means any observable operating on system A alone

(i.e., operating within A's Hilbert space).

Applying this formulation to measurement state (5),

the reduced density operators for the quantum system (call it A) and its detector (call it B), respectively, are

$$\hat{\rho}_A = \frac{1}{2}(|\psi_1\rangle\langle\psi_1| + |\psi_2\rangle\langle\psi_2|) \qquad \textbf{(10)}$$

$$\hat{\rho}_B = \frac{1}{2}(|1\rangle\langle 1| + |2\rangle\langle 2|) \qquad \textbf{(11)}$$

Plus signs in (10) and (11) make one think of superpositions such as (3), but these are **not** superpositions.

The density operator for the superposition (3) has cross-terms:

$$\hat{\rho} = (|\psi\rangle\langle\psi|) = \frac{1}{2}(|\psi_1\rangle\langle\psi_1| + |\psi_1\rangle\langle\psi_2| + |\psi_2\rangle\langle\psi_1| + |\psi_2\rangle\langle\psi_2|) \qquad \textbf{(12)}$$

Two cross-terms, involving both $|\psi_1\rangle$ and $|\psi_2\rangle$, are missing in (10).

So (10) does not describe a system in a superposition of two quantum states.

However, (10) is precisely the density operator one should use if one **knows** the quantum system **is either** in state $|\psi_1\rangle$ **or** in state $|\psi_2\rangle$ but one didn't know which and so, due your own to lack of information,

—> you simply assign a probability of 1/2 to each of the two possibilities.

Same goes for (11).

(10) and (11) are **"classical"** probabilistic states - **analogous** to "states of knowledge" one would assign to a coin flip when you know outcome to be either heads or tails with equal probability **but don't know yet** which has occurred.

Remember the dice example also!

The situation described by a density operator such as (10) is known as a "mixture" of states $|\psi_1\rangle$ and $|\psi_2\rangle$, as distinct from a "superposition" of states observed in the Mach-Zehnder experiment and represented by (3).

Equation (9) tells us that all correct statistics for subsystem A alone can be found from standard formula (7) applied to subsystem A **alone**.

But we have just seen that (10) is the density operator one should use if one knows A to be in either $|\psi_1\rangle$ and $|\psi_2\rangle$ without knowing which.

The same goes for subsystem B and (11).

In the case of Schrodinger's cat, it follows that the observer of the cat alone sees outcomes appropriate to a cat that is **either** alive **or** dead, **not** both.

For subsystems, the interference terms are missing, and an "ensemble" of repeated trials **must** exhibit a nonsuperposed mixture rather than a superposition.

This is the clear prediction of quantum physics for the entangled state (5).

But one must be careful, because (10) and (11) are not complete descriptions of the quantum states of the nucleus or the cat.

In fact, (10) and (11) are not quantum states at all, but merely **"reduced states"** arising from the actual state (5) of the composite system when one part of the composite system is removed from the equations..

In the case of Schrodinger's cat, (10) and (11) give the correct predictions for observations of either the nucleus alone or the cat alone, but do not represent the state of either subsystem because this given by (5).

In fact, when two quanta are entangled, neither one has quantum state of its own!

But B's state of affairs is certainly **not entirely** described by (11).

Rather, it is described by the composite state (5).

Equation (11) **merely** tells us following:

If the cat and the nucleus are in state (5) then, when one looks at the cat, one is going to see a cat that is either alive or dead.

There is no claim that (11) represents the complete quantum state of the cat.

That is, there is no claim that the cat is really in either state $|1\rangle$ or state $|2\rangle$, because the state it's really in is admittedly (5).

In fact, we do have complete knowledge of the state of both A and B, namely measurement state (5).

Reduced operators admittedly do not represent the state of the composite system.

They tell us only what will observe at the nucleus and at the cat

and they tell us nothing about the correlations between these observations,

so these density operators do not tell us the real state of the system.

And so plot thickens.

The entangled state (5) properly describes both individual subsystems.

However, the plus sign in (5) signifies the superposition of two terms.

We know, however, that neither subsystem A nor subsystem B is superposed.

What is meaning of plus sign?

This superposition arose from superposition represented by (3).

We cannot logically ignore this fact -

a strategy known as "shut up and calculate" approach to quantum measurement.

Instead, we must ask:

Exactly what is superposed when two subsystems are in this entangled state?

Superpositions must preserve the all-important **unity of quantum**.

When Max Planck proposed in 1900 that electromagnetic radiation occurs in energy units of magnitude $E = h\nu$, he tacitly implied the central quantum principle:

The unity of an individual quantum.

Energy (electromagnetic energy in case of radiation) comes in spatially extended bundles, each having a definite and identical quantity of energy.

One cannot have half a quantum, or 2.7 quanta.

You must have either 0 or 1 or 2 etc. quanta.

This is a fairly natural notion - apparently nature prefers to sub-divide the universe into a countable or even a finite set of entities as opposed to an uncountable continuum.

The spatial extension of these bundles **then** implies nonlocality.

If we have one quantum and destroy it (by transforming it to something else),

we **must** destroy all of it **everywhere**,

simultaneously, because we **cannot**, at any time, have **just** part of quantum.

Louis de Broglie put it perfectly in 1924, regarding another kind of quantum, namely the electron:

> **The energy of an electron is spread over all space with a strong concentration in a very small region. ...That which makes an electron an atom of energy is not its small volume that it occupies in space – I repeat it occupies all space – but the fact that it is indivisible, that it constitutes a unit.**

When one transforms state of a quantum, one **must** transform the entire extended quantum all at once.

Hence there are observed quantum jumps in experiments.

Furthermore, composite entangled systems such as atoms **also** behave in a unified fashion.

This unity is the source of nonlocality seen in experiments involving entangled pairs of photons.

Nonlocality is exactly what one would expect, given the unity and spatial extension of quantum and unitary (i.e. unity-preserving) nature of entanglement process.

Standard nonrelativistic quantum theory seems to prescribe two kinds of time evolution:

collapse upon measurement,

and Schrodinger equation between measurements.

The key feature of Schrodinger equation is that it prescribes a so-called "unitary" time evolution,

meaning time evolution that preserves pure states,

i.e. transforms unit Hilbert space vectors into other unit vectors.

Some ideas that are required physically by the unity of quantum are expressed as follows:

If quantum is described by pure quantum state at t = 0, it should **remain** pure at later times.

This notion prompts us to **ask** whether the measurement process also preserves pure states.

At least in case of idealized process described in (4), the answer is **"yes"** because **both** "before" and "after" states are pure.

Measurement state (5), since it is pure, **represents** a highly unified state of affairs, **even** though one of its subsystems is a macroscopic detector.

Thus, one suspects that this state, like its progenitor (3), is truly a superposition in which the superposed terms **represent** two situations or states of the same object.

But precisely what is that object, i.e., **what** is superposed?

We have seen that the states of subsystem A are **not** superposed,

nor are the states of subsystem B.

The conventional interpretation (which, as we will see, is subtly incorrect) of a product state such as $|\psi_1\rangle|1\rangle$ is that it represents a state of a composite system AB in which subsystem A is in state $|\psi_1\rangle$ while B is in state $|1\rangle$ (we used such a state in the two-path experiments).

In this case, (5) would represent a superposition in which AB is simultaneously in the state $|\psi_1\rangle|1\rangle$ and also in the state $|\psi_2\rangle|2\rangle$.

The situation for Schrodinger's cat would be: live cat and undecayed nucleus superposed with dead cat and decayed nucleus.

This is at least as physically outrageous as a live cat superposed with a dead cat, and it contradicts physical implications (a cat that is either alive or dead) of reduced states (10) and (11) as described earlier.

Something is still wrong!

Let us repeat some of our discussion and provide more details to work it all out.

Some Repetition and More Intricate Details

Proposal:

The solution to the so-called quantum measurement problem is completely contained within standard quantum mechanics and needs no elaborate new structures and interpretations.

Remember the standard discussion from earlier:

The following discussion is at limits of knowledge you have learned about QM, but I thought you should see what researchers including myself are doing at the present time.

The details of the derivations are not important, ONLY the conclusions.

Total system T = quantum system S (states $|s_i\rangle$) + measuring device A (states $|a_i\rangle$), where $|a_0\rangle =$ state of measuring device "off"

The unitary time evolution rule then says

$|s_1\rangle\,|a_0\rangle \to |s_1\rangle\,|a_1\rangle$ unitary evolution $\qquad$ $|s_2\rangle\,|a_0\rangle \to |s_1\rangle\,|a_2\rangle$ unitary evolution

The linearity of QM then says that

$$|\psi\rangle_{SA} = (c_1 |s_1\rangle + c_2 |s_2\rangle) |a_0\rangle = c_1 |s_1\rangle |a_0\rangle + c_2 |s_2\rangle |a_0\rangle \rightarrow c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

In this derivation

$$|\psi\rangle_S = c_1 |s_1\rangle + c_2 |s_2\rangle$$

is state of quantum system(superposition), while

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

is state of combined system(entanglement/superposition).

The quantum measurement problem is the realization that the state

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

is **not** observed as outcome of measurement!

What is seen is not a so-called superposition, but either

$$|s_1\rangle\,|a_1\rangle \;\text{ or }\; |s_2\rangle\,|a_2\rangle$$

That is the so-called **"problem of outcomes"**.

The state

$$|\psi\rangle_{SA} = c_1\,|s_1\rangle\,|a_1\rangle + c_2\,|s_2\rangle\,|a_2\rangle$$

is usually referred to as a superposition, which is **misleading**.

Entanglement **outweighs** superposition as the **defining** feature of this state.

Without entanglement-correlations - we would **not** have a measurement problem!

Careful investigation of this state in a 2007 experiment(Roch, et al) (a Wheeler-type delayed choice experiment) demonstrates its strikingly non-local character.

We will discuss the experiment in detail shortly.

A photon jumps from state $|\psi\rangle_S = c_1 |s_1\rangle + c_2 |s_2\rangle$ to state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ precisely when A(measuring device) switches on and while photon is still inside interferometer, and jumps from $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ to $|\psi\rangle_S = c_1 |s_1\rangle + c_2 |s_2\rangle$ when A switches off during the experiment.

Quantum jumps, removal of interferences, and the observed non-locality are due to entanglement.

Of course, an entangled state = superposition, but it is a very **special** superposition.

To call $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ simply a “superposition” **misses** the crucial physics of entanglement and makes all the difference in the understanding of the state.

Entanglement is the characteristic trait of quantum mechanics; one that enforces the entire departure from classical lines of thought.

It is well known that, for 2-part systems, **all** non-product states exhibit non-locality (Bell inequality).

Thus when S and A are entangled $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$, they **share** a so-called non-local **channel**.

The measurement state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ is very subtle.

Although $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ is often called a superposition of S and/or superposition of A, **neither** case is true.

Thus, when S and A in measurement state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$, neither S nor A in a superposition. $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ = superposition, but neither a superposition of S nor of A and also not superposition of states of composite system SA.

In $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$, S is in both states $|s_1\rangle$ and $|s_2\rangle$ simultaneously, as is known from observed interference between 2 states.

However, in case of $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ **experiment** now shows that SA is not in both two-part states $|s_1\rangle |a_1\rangle$ and $|s_2\rangle |a_2\rangle$ simultaneously, **but only simultaneously in two correlations between the states.**

The entanglement of **two** systems is quite **different** from superposition of **one** system.

Experiments(Rarity) demonstrated the **precise** sense in which $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ **represents** a superposition.

Experiments answer question "given that S and A are in state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$, **what** (if any) entities interfere and **what** is nature of interference?"

Let us see exactly how……..

The Local State Solution(due to Jauch) of the Problem of Definite Outcomes

Consider a **single** quantum S(electron or photon), passing through double-slit experiment, with a "downstream" viewing screen.

Suppose an ideal "which-slit detector" A is present so that, upon detection, S and A become entangled in measurement state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ with **orthogonal** detection states $|a_i\rangle$.

Imagine S and A are possibly separated by meters or kilometers.

Complete observation of experiment **requires** two “local observers”, The 1st observer is S and the 2nd observer is A.

Such a non-local setup **has been carried** out experimentally.

What did the **local** observers observe this experiment?

A well-known prediction of quantum physics says

that the 1st observer observes implications of the **“local state of S,”**

represented by a **“reduced”** density operator,

where the degrees of freedom for the 2nd system

are averaged over(removed) by the “trace” (Tr) operation (see below)

$$\rho_S = \mathrm{Tr}_A(\rho_{SA}) = |c_1|^2 |s_1\rangle \langle s_1| + |c_2|^2 |s_2\rangle \langle s_2|$$

and 2nd observer would observe the implications of **“local state of A”**, represented by the reduced density operator

$$\rho_A = \mathrm{Tr}_S(\rho_{SA}) = |c_1|^2 |a_1\rangle \langle a_1| + |c_2|^2 |a_2\rangle \langle a_2|$$

where a density operator gives probabilities via the relation

$$P(b) = \mathrm{Tr}(\rho P_b) = \mathrm{Tr}(\rho \left|b\right\rangle \left\langle b\right|)$$

The “local state of S” is found by **completely removing** from the density operator any effects of A and vice versa.

This is the important idea!

Derivation of reduced density operator (for completeness)

$$\rho_{SA} = \left|\psi_{SA}\right\rangle \left\langle\psi_{SA}\right| = (c_1 \left|s_1\right\rangle \left|a_1\right\rangle + c_2 \left|s_2\right\rangle \left|a_2\right\rangle)(c_1^* \left\langle s_1\right| \left\langle a_1\right| + c_2^* \left\langle s_2\right| \left\langle a_2\right|)$$

$$\begin{aligned}
\rho_S &= \mathrm{Tr}_A(\rho_{SA}) = \sum_{k=1}^{2} \left\langle a_k\right| \rho_{SA} \left|a_k\right\rangle \\
&= \sum_{k=1}^{2} \left\langle a_k\right| \left(\left|\psi\right\rangle_{SA\ \ SA}\left\langle\psi\right|\right) \left|a_k\right\rangle \\
&= \sum_{k=1}^{2} \left\langle a_k\right| \left((c_1 \left|s_1\right\rangle \left|a_1\right\rangle + c_2 \left|s_2\right\rangle \left|a_2\right\rangle)(c_1^* \left\langle s_1\right| \left\langle a_1\right| + c_2^* \left\langle s_2\right| \left\langle a_2\right|)\right) \left|a_k\right\rangle \\
&= \sum_{k=1}^{2} (c_1 \left|s_1\right\rangle \delta_{k1} + c_2 \left|s_2\right\rangle \delta_{k2})(c_1^* \left\langle s_1\right| \delta_{k1} + c_2^* \left\langle s_2\right| \delta_{k2}) \\
&= \left|c_1\right|^2 \left|s_1\right\rangle \left\langle s_1\right| + \left|c_2\right|^2 \left|s_2\right\rangle \left\langle s_2\right|
\end{aligned}$$

where we have used orthonormality of measurement states via $\langle a_i \,|\, a_j \rangle = \delta_{ij}$

We note that the Tr (trace) operation = sum over all designated states

removes all knowledge of the designated(traced) system from the equation.

This is clearly useful i**f we do not know much** about the reduced system.

A similar result holds for ρ_A .

Now, continuing our discussion.

Clearly, the reduced states are **mixtures**, not superpositions.

QM predicts both LOCAL observers **find(see)** mixtures **not** superpositions.

An ensemble of experimental trials verifies

this via mixed-state patterns in agreement with the assertion made earlier

that neither S nor A is in a superposition.

For a different example,

$$\rho_A = \mathrm{Tr}_S(\rho_{SA}) = |c_1|^2 |a_1\rangle \langle a_1| + |c_2|^2 |a_2\rangle \langle a_2|$$

predicts Schrödinger's cat is in a mixture of either dead or alive,

not a superposition of both dead and alive.

The **local states**

$$\rho_S = \mathrm{Tr}_A(\rho_{SA}) = |c_1|^2 |s_1\rangle \langle s_1| + |c_2|^2 |s_2\rangle \langle s_2|$$

and

$$\rho_A = \mathrm{Tr}_S(\rho_{SA}) = |c_1|^2 |a_1\rangle \langle a_1| + |c_2|^2 |a_2\rangle \langle a_2|$$

must be taken seriously as implying outcomes predicted to be observed at the two sites.

The local states **cannot be dismissed** simply

by the argument that the only "real" state is the "global state"

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

There is no contradiction between the predicted local **mixtures**

$$\rho_S = \mathrm{Tr}_A(\rho_{SA}) = |c_1|^2 |s_1\rangle \langle s_1| + |c_2|^2 |s_2\rangle \langle s_2|$$

$$\rho_A = \mathrm{Tr}_S(\rho_{SA}) = |c_1|^2 |a_1\rangle \langle a_1| + |c_2|^2 |a_2\rangle \langle a_2|$$

and the unitarily-evolving global **pure state** .

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

Important:

We are saying: To consider the combined system SA as a single system only evolving unitarily **misses** the essential physics of nonlocality.

Haroche and Raimond weighed in on the question of

when a two-part system should be considered as a composite of two subsystems,

versus when it should be considered a single system B = SA.

“The composite system should be considered as single whenever the binding between the parts is much stronger than the interactions involved in dynamics, so that the internal structure of the composite system is left unchanged as it travels through the experiment.”

By this criterion, SA not single system.

Not only does the relation between S and A change during experiment, the relation is entangled and thus nonlocal.

The implication is that $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ **must be** considered an entanglement of two separate systems S and A, not superposition of single composite system SA.


That's the basics of the theory at this point in the discussion(at this point in my tortuous development! **It follows completely from standard QM.**


Not sure yet! Let us do it again!

Now for even more details and some possible new interpretations

Start new discussion with a new experiment as we always should!!!!!

Roch's "delayed choice" experiments used a Mach-Zehnder interferometer (1st figure) rather than the logically equivalent double-slit setup, to observe photons. It is shown in detail in the 2nd figure on next slide.





While a photon is on the **48-meter-long** interferometer paths,

a quantum-based random number generator “decided” whether the second beam splitter (positioned at the end of paths - note the spatial separation) would be incorporated or omitted,

i.e., it decided whether detectors would not or would (respectively) determine “which path.”

In trials incorporating second beam splitter, an interference pattern is observed,

indicating the photon passed through device as superposition along both paths.

On trials omitting second beam splitter, no interference is observed, indicating the photon passed through device along one or other path.

The two parallel paths were 5 millimeters apart.

Precisely (so far as the experiment could determine)
when the second beam-splitter switched from “on” to “off,”
the photon collapsed in **mid-flight**
from being on both paths to being on one or the other path.

The quantum jump was **correlated** with and **coincident** with
the incorporation or the omission of the second beam splitter
(i.e., with the **decision** to not entangle **or** to entangle the detectors with the photon).

All experimental results **are** just as predicted correctly by the reduced states

$$\rho_A = \mathrm{Tr}_S(\rho_{SA}) = |c_1|^2 |a_1\rangle \langle a_1| + |c_2|^2 |a_2\rangle \langle a_2|$$

$$\rho_S = \mathrm{Tr}_A(\rho_{SA}) = |c_1|^2 |s_1\rangle \langle s_1| + |c_2|^2 |s_2\rangle \langle s_2|$$

Switching the second beam-splitter off entangles the photon and the detector in measurement state

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

causing the photon to collapse from a superposition

$$|\psi\rangle_{S} = c_1 |s_1\rangle + c_2 |s_2\rangle$$

into the mixture

$$\rho_S = \mathrm{Tr}_A(\rho_{SA}) = |c_1|^2 |s_1\rangle \langle s_1| + |c_2|^2 |s_2\rangle \langle s_2|$$

that is observed at the detector.

At this point we are in the diagonal density matrix stage.

Thus, we are doing a classical measurement(like dice) and it should be interpreted as such!

This should resolve problem of definite outcomes!

This was my belief at this point in my tortuous path to understanding!

Quantum theory predicts and experiment verifies that, with the detector in operation,

observers of S and of A find them to be in definite mixtures,

not indefinite superpositions.

Any strategy of imagining the quantum and the detector to be widely separated

obviously changes nothing

- it does not matter whether the quantum and the detector are close together or far apart.

The key to understanding quantum measurements comes from

understanding nonlocal relationship that develops between S and A

when they evolve unitarily into entangled measurement state

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

In experiments, this global state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$

violates Bell's inequality, implying instantaneous non-local transfer of **correlations** across arbitrarily large distances.

Not information transfers, but
CORRELATIONS which we can only see later!

Without entanglement one would observe different results

- the interferences would not disappear as in the experiments.

But entanglement “decoheres-collapses” coherent states

so that S and A impact their detectors randomly.

Locally, entanglement “decoheres” each photon so that they exhibit definite outcomes.

But quantum dynamics is unitary, implying that the global state

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

remains coherent despite the incoherence of subsystems.

Since the individual photons are now incoherently mixed, what has happened to coherence?

Answer:

It is the experimentally observed global coincidence measurements

which compare the impact points of entangled pairs.

In the double-slit experiment with two screens, each photon "knows" the impact point (i.e., phase shift) of other photon and instantly adjusts own impact point in order to form an interference pattern as a function of difference between two photons' phase shifts!

This is strikingly non-local, and the experimental results violate Bell's inequality.

Thus the coherence of the entangled state resides in correlations between subsystems, rather than in the subsystems themselves.

Entanglement **transforms** the coherence of states of S into coherence of correlations **between** states of S and A, **allowing** S and A to exhibit definite outcomes while **preserving** the global coherence as demanded by unitary evolution.

We can now answer the question:

Precisely what is superposed and what interferes in the measurement state?

The answer is surprisingly simple:

Only the correlations between S and A are superposed.

Thus the measurement state $|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$ should be read as:

The state $|s_1\rangle$ is positively correlated with the state $|a_1\rangle$,
and the state $|s_2\rangle$ is positively correlated with the state $|a_2\rangle$

Only correlations are superposed, not states.

When the superposition $|\psi\rangle_S = c_1 |s_1\rangle + c_2 |s_2\rangle$ of S entangles with states of A, the superposition **shifts**, from a superposition of states of S to superposition of correlations between S and A, so S can be in an incoherent mixture **while** maintaining unitary global dynamics.

This is how nature resolves problem of definite outcomes.

So the coherence exhibited by the measurement state

$$|\psi\rangle_{SA} = c_1 |s_1\rangle |a_1\rangle + c_2 |s_2\rangle |a_2\rangle$$

must be **invisible** to local observers,

and **yet show up** in the global measurement state in order to **preserve** unitary dynamics.

One could regard this as the **underlying reason** why entanglement (i.e., measurement)

must shift coherence from the states of S and A to the correlations between S and A.

"Collapse" or the "appearance of definite values" (a reduced density operator)

can be viewed as a consequence of the measurement state's nonlocality

plus special relativity's ban on instant signaling.

We note that the global measurement state is a very **different** animal from local states.

While the local mixed states are immediately observed at both local sites,

the global state can be "observed" only at some time after measurement by traveling to both local sites, gathering data from both, and then assembling information and noting the correlations between two sets of data.

Our conclusions so far:

When the detector measures the superposed quantum,

quantum physics predicts that the states actually observed are

local (i.e. mixed or reduced) states of subsystems,

it is not the superposed global state that follows from Schrödinger's equation.

The local states directly observed in the measurement

must contain **no** hint of nonlocal correlations

between two subsystems since this would violate relativity's prohibition on instant signaling.

Thus the local states describe what **actually** happens at both subsystems.

The global state predicts these local states(they can be derived from it),

and **also** predicts indirectly-observable (by gathering global data at **later** time)

nonlocal correlations between these states.

Continuing along my path to final understanding…..

An even more dramatic experiment — Experimental nonlocality and entanglement

Now we demonstrate that, according to standard quantum theory (**and** experiment),

measurement state (5) represents none of paradoxical situations we have mentioned.

Again let us repeat material from earlier - understanding this topic is so important that repetition is warranted.

The **unity** of the quantum suggests that the measurement state (5) represents a unified,

hence superposed and pure, quantum state of composite system.

We asked the question: precisely what is superposed?

Studied simple (i.e. non-composite) superposition (3) via interference exhibited in M-Z experiment.

Varying the length of either path 1 or path 2

created varying interference effects in detectors,

demonstrating each photon really must travel both paths to its detector.

Quantum theory agrees entirely with these conclusions,

as can be shown by using photon wavelengths

to show that path differences correctly predict interferences observed at each detector.

This implies that to understand the measurement state,

one needs to find and analyze entanglement experiments that demonstrate interference.

This has been done for several decades in connection with quantum non-locality.

Many nonlocal interference experiments have been done

beginning with Clauser and Freedman,

culminating in experiments demonstrating nonlocality across great distances

and that simultaneously closed all possible loopholes in all previous experiments.

By now, it is well known that entangled state (5) predicts nonlocal effects

between two subsystems, and that phase variations of either subsystem

cause instantaneous(non-signaling), i.e., non-local, readjustments(correlations)

of the possibly-distant other subsystem.

When macroscopic systems are involved (i.e., cats) we have a problem.

it is not easy to **vary the phase of cat**,

and as we saw in the Mach-Zehnder experiment,

one **cannot** understand superposition without varying phases of the superposed parts.


Thus, all nonlocality experiments carried out with pairs of simpler quanta such as photons.


The best nonlocal entanglement experiments

most appropriate for investigating measurement

were conducted nearly simultaneously by Rarity,Tapster and Ou, Zou, Wang (RTO).

Figure shows layout for these "RTO" experiments.





The “Source" creates entangled photon pairs by "parametric down-conversion”, a process which we discussed earlier.

The RTO experiment is **two back-to-back MZ-interferometer experiments** but with first beam splitter for each photon located **inside** the source of entangled photons.

Without entanglement, each single photon (either A or B) would interfere with itself at its own detectors according to own phase shifters ϕ_A or ϕ_B.

In the experiment, two entangled photons are emitted into a superposition of solid paths connecting detectors A1 and B1, and dashed paths connecting detectors A2 and B2.

Remember that the two photons are already entangled when emitted.

Entanglement changes everything.


No longer does either photon interfere with itself at own detectors.

Instead, photons are entangled in measurement state (5) with $|\psi_1\rangle$ and $|\psi_2\rangle$

representing (say) solid-line and dashed-line states of A and $|1\rangle$ and $|2\rangle$

representing solid-line and dashed-line states of B,


although in RTO experiments **neither** subsystem is macroscopic.



Each photon **now** acts like a which-path detector for the other photon.

Recall the double-slit experiment:

When the which-slit detector is switched on, the pattern on screen switches abruptly from a striped interference pattern indicating the pure state nature of each electron across both slits, to phase-independent sum of two non-interfering single-slit patterns.



Entanglement between the electron and the which-slit detector breaks the pure state into two single-slit parts, so that measured electron comes through either slit 1 or slit 2.

This suggests that in the RTO experiment, the entanglement should break the pure-state superposition (3) into two non-interfering parts.


This is exactly what is observed.

Both photons impact their detectors as random 50-50 mixtures, just like a flipped coin.

Entanglement **breaks** the single-photon pure state (3) observed in the Mach-Zehnder experiment, causing each photon to behave "incoherently" with no dependence on its phase setting.

But (5) is pure state.

Where has the phase dependence gone?



The **answer** lies in the phase-dependent but nonlocal relationship observed **between** the solid and the dashed branches.


This phase dependence is observed experimentally in coincidence (or **correlation**) measurements comparing the detections of entangled pairs.

The “flipped coins” mentioned above turn out to be correlated with each other.


This phase dependence across two separated subsystems is **essential to preserve** unity of (now entangled) quantum.

This is **not** an easy experiment to perform:

The source creates a stream of photon pairs, and one must compare the impact of single photon A at detectors A1, A2 with impact of corresponding entangled photon B at detectors B1, B2.




RTO figured out how to do this, with result shown in Figure.



The figure graphs the **degree of correlation** between A and B.

This is a **measure of the agreement** between the outcomes at A's detectors and B's detectors.



A correlation of +1 means perfect, or 100%, agreement:

Either both sets of detectors register outcome 1 (i.e., A1 and B1 click)

or both register outcome 2.

The opposite extreme is a correlation of -1, meaning 100% disagreement: If one detector registers 1, the other registers 2.

Either correlation, +1 or -1, implies that either photon's outcome is **predictable** from the other photon's outcome.

A correlation of zero means one photon's outcome does not at all determine other's outcome: Each photon has random 50-50 chance of either outcome regardless of the other photon.

Correlations between 0 and +1 mean the outcomes are more likely to agree than to disagree, with larger correlations denoting higher probability of agreement; for example, correlation of +0.5 means 75% probability of agreement.

Similarly, correlations between 0 and -1 mean outcomes are more likely to disagree than to agree; a correlation of -0.5 means a 75% probability of disagreement.

The RTO experiment agrees entirely with the predictions of standard quantum physics.

When accounting is made of optical paths for both photons, QM obtains the following result:


$$P(correlated) = P(A1\ and\ B1) + P(A2\ and\ B2) = \frac{1}{2}[1 + \cos(\varphi_B - \varphi_A)]$$

$$P(anticorrelated) = P(A1\ and\ B2) + P(A2\ and\ B1) = \frac{1}{2}[1 - \cos(\varphi_B - \varphi_A)]$$

where P(correlated) is single-trial probability that A's and B's detectors will agree,

and P(anticorrelated) is a single-trial probability that A's and B's detectors will disagree.

The degree of correlation, defined as
P(correlated) – P(anticorrelated),
is then simply $\cos(\phi_B - \phi_A)$, as graphed in last figure.



In 1964, John Bell, as we discussed earlier, published a ground-breaking article stating the sufficient condition for a statistical theory such as quantum physics to meet a condition known as "locality."


He defined locality to mean **"that the result of measurement on one system be unaffected by operations on distant system with which it has interacted in past"**.

Bell expressed a sufficient condition in the form of inequality that any local theory must obey.

He demonstrated that certain statistical predictions of quantum physics violate Bell's inequality, i.e. quantum physics makes nonlocal predictions.

The results in last figure implies the case in point: the figure violates Bell's inequality at all phase differences $\phi_B - \phi_A$ other than 0, π, and 2π.

Let me **<u>underline</u>** meaning of this:





Violation of Bell's inequality means that the statistics of the measurements on photon A - photon A's **"statistical behavior"** - is necessarily affected by the setting of photon B's phase shifter.

In fact, even without Bell's condition, the nonlocality of the experiment is intuitively obvious.

Here's why:

Suppose we set the phase shifters to zero and that all four optical paths (two solid, two dashed) in are then equal; thus $\phi_B - \phi_A$ is zero.





Without the two beam splitters BS,

the two photons emitted into the solid pair and the dashed pairs of paths

would impact either detectors A1 and B1 or A2 and B2

because of the symmetry of experiment and conservation of momentum.

This is neither surprising nor nonlocal,

and would happen even if the photons were not entangled.

But the beam splitter is a **randomizing** device

that mixes the solid and dashed paths;

—> any photon passing through it has 50-50 chance of reflection or transmission.

With non-entangled photons and both beams splitters in place, there would then be no correlation between photon A's outcome and B's outcome because the two photons are independent of each other.

With entanglement, the correlation is perfect.

How does one photon know which path the other photon took at the other photon's beam splitter? **poor question!!**

Each photon is now "detecting" the quantum state of the other photon, from a distance that could be large.

Perfect correlation certainly "feels" nonlocal even though (as mentioned above) this perfect correlation at $\phi_B - \phi_A = 0$ does not violate Bell's inequality.

Note that such a violation is a sufficient but not a necessary condition for nonlocality.

Non-locality is **written** all over the RTO experiment.

Each photon "knows" which direction the other photon takes at its beam splitter and adjusts its selection accordingly.

The key nonlocal feature of graph, which is simply a cosine function, has ($\phi_B - \phi_A$) as its independent variable.

Thus any desired shift in correlations can be made by an observer at either of

possibly-widely-separated phase shifters.

Bell suspected that this kind of situation meant that observer A (call her Alice)

could use her phase shifter to alter outcomes that would have occurred at both her own and observer B's (call him Bob) detector and, following up on this hypothesis, he derived his inequality involving probabilities at both Alice's and Bob's detectors

which, **if violated**, implied that both photons must have readjusted their states.

Such a readjustment is just what we expect,

given the **unity** of the quantum

and thus the unity of atoms and other entangled systems such as our two photons.

The two photons form a single **“bi-quantum", an "atom of light”**, in the pure state (5).

When Alice varies her phase shifter, both photons "know" both path lengths and **readjust** their behavior accordingly to produce the proper correlations.

Analogously, a single photon "knows" both path lengths in single-photon interferometer experiment.

Finally, we come to the central question of the discussion:

What is actually superposed in the entangled superposition (5)?

A Mach-Zehnder experiment tests the simple superposition (3), while the RTO experiment tests the entangled superposition (5).

We know what is superposed in Mach-Zehnder, namely quantum states $|\psi_1\rangle$ (path 1) and $|\psi_2\rangle$ (path 2).

This is deduced from the effect that either phase shifter has on both states.

Now consider the RTO experiment.

What is the effect of shifting either phase shifter?

One thing that does not change is the state ("local state" would be a better term, as discussed earlier) of either photon A or photon B:

As we know, both photons remain in 50-50 mixtures regardless of either phase setting.

What does change with variations in either phase shifter is the correlations between A and B.

With $\phi_B - \phi_A = 0$ they have perfect correlation:

Either A1 and B1 (which we denote (11)) or A2 and B2 (denoted (22)).

As we vary either ϕ_B or ϕ_A we obtain non-zero probabilities of anti-correlated individual trials,

denoted (12) (outcomes A1 and B2) and (21) (A2 and B1).

When the non-local phase angle difference ($\phi_B - \phi_A$) reaches $\pi/2$,

we have zero correlation, and when it reaches π we have perfect anti-correlation.

Table 1 summarizes the crucial points in more detail.

Simple superposition:		**Entangled superposition of two sub-systems:**		
φ	State of photon	$\varphi_B - \varphi_A$	State of each photon	Correlation between the two photons
0	100% "1", 0% "2"	0	50-50 "1" or "2"	100% corr, 0% anti
$\pi/4$	71% "1", 29% "2"	$\pi/4$	50-50 "1" or "2"	71% corr, 29% anti
$\pi/2$	50% "1", 50% "2"	$\pi/2$	50-50 "1" or "2"	50% corr, 50% anti
$3\pi/4$	29% "1", 71% "2"	$3\pi/4$	50-50 "1" or "2"	29% corr, 71% anti
π	0% "1", 100% "2"	π	50-50 "1" or "2"	0% corr, 100% anti

Table 1. In a simple superposition, the photon's state varies with phase angle. In an entangled superposition, the relationship between states of the two photons varies, while individual states of both photons are phase-independent (or "mixed").

The column titled "simple superposition" shows how the superposition state of single photon (M-Z) varies from "100% state 1" to "100% state 2" as phase angle between two states varies.

Column titled "entangled superposition of two subsystems" shows that the state of each photon remains unchanged throughout entire range of both phase settings, while nonlocal correlation between the states of two photons varies from "100% correlated" to "zero correlation" and then to "100% anticorrelated" as either of the two local phase angles varies.

So, once again, what is superposed in the RTO experiment?

The hallmark of a superposition is the dependence on phase difference between the objects superposed.

But Table 1 exhibits **no** such phase dependence of the states of the two photons.

Simple superposition:		**Entangled superposition of two sub-systems:**		
φ	State of photon	$\varphi_B - \varphi_A$	State of each photon	Correlation between the two photons
0	100% "1", 0% "2"	0	50-50 "1" or "2"	100% corr, 0% anti
$\pi/4$	71% "1", 29% "2"	$\pi/4$	50-50 "1" or "2"	71% corr, 29% anti
$\pi/2$	50% "1", 50% "2"	$\pi/2$	50-50 "1" or "2"	50% corr, 50% anti
$3\pi/4$	29% "1", 71% "2"	$3\pi/4$	50-50 "1" or "2"	29% corr, 71% anti
π	0% "1", 100% "2"	π	50-50 "1" or "2"	0% corr, 100% anti

Table 1. In a simple superposition, the photon's state varies with phase angle. In an entangled superposition, the relationship between states of the two photons varies, while individual states of both photons are phase-independent (or "mixed").

Each photon remains in an unchanging 50-50 mixture of own "path 1" and "path 2" states

- a situation that is radically at odds with the true superposition of path 1 and path 2 exhibited by the M-Z experiment.

Thus, in the entangled RTO state, **neither** photon is superposed.

We see here the source of the "classical" or non-superposed
nature of reduced density operators (Eqs. (10) and (11)),
not to mention the non-superposed
and hence non-paradoxical nature of Schrodinger's cat.

Examination of the phase-dependence of the measurement state (5), as demonstrated by nonlocality experiments such as the RTO experiment,

reveals the true nature of Schrodinger's cat.

The last column of Table 1 shows us what actually is superposed when two subsystems are entangled in measurement state (5).

Simple superposition:		**Entangled superposition of two sub-systems:**		
φ	State of photon	$\varphi_B - \varphi_A$	State of each photon	Correlation between the two photons
0	100% "1", 0% "2"	0	50-50 "1" or "2"	100% corr, 0% anti
$\pi/4$	71% "1", 29% "2"	$\pi/4$	50-50 "1" or "2"	71% corr, 29% anti
$\pi/2$	50% "1", 50% "2"	$\pi/2$	50-50 "1" or "2"	50% corr, 50% anti
$3\pi/4$	29% "1", 71% "2"	$3\pi/4$	50-50 "1" or "2"	29% corr, 71% anti
π	0% "1", 100% "2"	π	50-50 "1" or "2"	0% corr, 100% anti

Table 1. In a simple superposition, the photon's state varies with phase angle. In an entangled superposition, the relationship between states of the two photons varies, while individual states of both photons are phase-independent (or "mixed").

Since the correlations between two photons vary sinusoidally as the non-local phase angle between the two photons varies, clearly these are correlations between the states of two photons, and not the states themselves, that are interfering.

Entanglement has shifted superposition, from states of one photon A (Eq. (3), M-Z) to correlations between photon A and photon B (Eq. (5), RTO).

More learning, More repetition, More Details and Some Tentative Conclusions

We are getting closer……

In order to resolve the problem of definite outcomes of measurements, aka Schrodinger's cat,

our discussion analyzed the entangled state (5) of a microscopic quantum

and its macroscopic measuring apparatus.

This state is a superposition of two composite entities $|\psi_1\rangle|1\rangle$ and $|\psi_2\rangle|2\rangle$,

with a phase angle between these entities that can range over 2π radians.

To resolve problem of definite outcomes we must ask:

Precisely what does the composite superposition (5) actually superpose, physically?

In order to understand a simple non-composite superposition (3),

we looked at the effect of varying the phase angle between

superposed entities $|\psi_1\rangle$ and $|\psi_2\rangle$

in an experimental setting such as M-Z interferometer.

The theoretically predicted and experimentally observed results then made it
obvious that the quantum whose state is (3) flows
simultaneously along two separate paths described by $|\psi_1\rangle$ and $|\psi_2\rangle$.

To understand superposition (5), one should proceed similarly by studying situations
in which the phase angle between superposed entities $|\psi_1\rangle|1\rangle$ and $|\psi_2\rangle|2\rangle$ varies.

Theorists and experimentalists studying the phenomenon of nonlocality
have been doing this for decades, but quantum foundations specialists have not particularly
noticed how the work is connected with the measurement problem.

In fact, the nonlocal aspects of the state (5) have been studied
since Bell's 1964 theoretical paper and Clauser and Freedman's 1972 experiment.

The 1990 experiments of Rarity and Tapster, and of Ou, Zou, Wang, and Mandel,
furnish an ideal vehicle for such an analysis and are the central feature of this discussion

One lesson of this analysis is that, in order to understand the measurement problem, one must understand the significance of nonlocality.

This is because the key measurement state (5) that caused Schrodinger and other experts so much concern has nonlocal characteristics.

It must be understood as a superposition of correlations, rather than a superposition of states, but this cannot become apparent until one considers the effect of variations in phase angle between its superposed terms.

Such variations are not part of the measurement process itself because measurements are designed to take place at zero phase angle.


Experimental or theoretical studies of such phase variations will have nonlocal ramifications, because such variations are inherently nonlocal!

This situation would have prevented Schrodinger in 1935, or indeed anyone prior to Bell's 1964 paper and the experimental confirmations of reality of nonlocality beginning in 1972, from understanding entangled superposition (5).

It's worth emphasizing that, when two subsystems are entangled in measurement state (5), neither subsystem is superposed.

Only the correlations between subsystems are superposed.

In RTO experiments, the two correlations in question are represented by solid and dashed paths connecting pairs outcomes.



A pair of photons entangled in state (5) follows both of these paths simultaneously.

The subsystems themselves, however, are not in superpositions, but instead, in indeterminate mixtures of definite states.

Thus observers of either subsystem will observe only definite outcomes, as predicted by local mixtures (10) and (11).

RTO experiments are the entangled analog of the M-Z interferometer experiment: a pair of back-to-back interferometer experiments, with entangled pair of quanta of which one quantum passes through each interferometer.

As we said earlier, the experiment and its theoretical analysis shows that, when superposed photon A becomes entangled with second photon B to form state (5), the nonlocal aspect of A's superposition is transferred to correlations between A and B.

Thus, an entangled state such as (5) is neither a superposition of states of A nor of states of B, but is instead a superposition of correlations between states of A and states of B.

To see most clearly, compare simple superposition (3) with entangled superposition (5).

In the simple superposition, the state observed by a "which- state" detector varies smoothly from 100% $|\psi_1\rangle$, through 50% $|\psi_1\rangle$ and 50% $|\psi_2\rangle$, and finally to 100% $|\psi_2\rangle$ as the phase angle ϕ between $|\psi_1\rangle$ and $|\psi_2\rangle$ varies from 0 to π.

In entangled superposition, neither the state of A nor the state of B varies as ϕ_A or ϕ_B varies;

both A and B remain in 50-50 mixtures throughout.

What does vary is the correlation between A and B.


A non-local "correlation detector" (i.e., an RTO-type of experiment!)

would find the relation between the two subsystems varies from 100% positively correlated (either pair state 11 or 22, pictured by solid and dashed paths in figure),

to 50% positively correlated and 50% anti-correlated, and finally to 100% anti-correlated (12 or 21), as the nonlocal phase difference $\phi_B - \phi_A$ varies from 0 to π.

This is a superposition of correlations,

not a superposition of composite states or of non-composite (single-system) states.



At least in idealized case of a minimally-disturbing von Neumann measurement, the initial stage of the measurement process (through formation of the measurement state (5)) can be **described** as follows:

A quantum in a simple superposition such as (3) entangles with a macroscopic which-path detector.

At the instant of entanglement, the local states of both the quantum and the detector undergo a radical change, **a quantum jump**.

Locally, the detector and the quantum jump into mixtures (10) and (11).

Simultaneously, the global state (5) continues evolving smoothly according to the Schrodinger equation.

Entanglement causes the superposed single quantum to be **instantly transformed** into superposed correlations between the quantum and the detector.

This stage of measurement process is **entirely describable** in terms of pure global states following Schrodinger equation.

“Collapse” from the local superposition to the local mixtures occurs because of the formation of the entangled state (5) **and** the resulting formation of subsystems whose local states (Eqs. (10) and (11)) have definite outcomes.

Note that the phenomenon of nonlocality is essential to preserving the pure-state nature (the unity) of the composite system.

To put this more intuitively, a reorganization throughout the entire extent of the composite entangled system is required in order to preserve unity of the (now entangled) quantum.

According to Table 1, when

two systems entangle to form the state (5),

both “collapse” into phase-independent

local mixtures.

Simple superposition:		**Entangled superposition of two sub-systems:**		
φ	State of photon	$\varphi_B - \varphi_A$	State of each photon	Correlation between the two photons
0	100% “1”, 0% “2”	0	50-50 “1” or “2”	100% corr, 0% anti
$\pi/4$	71% “1”, 29% “2”	$\pi/4$	50-50 “1” or “2”	71% corr, 29% anti
$\pi/2$	50% “1”, 50% “2”	$\pi/2$	50-50 “1” or “2”	50% corr, 50% anti
$3\pi/4$	29% “1”, 71% “2”	$3\pi/4$	50-50 “1” or “2”	29% corr, 71% anti
π	0% “1”, 100% “2”	π	50-50 “1” or “2”	0% corr, 100% anti

Table 1. In a simple superposition, the photon's state varies with phase angle. In an entangled superposition, the relationship between states of the two photons varies, while individual states of both photons are phase-independent (or "mixed").

Relativity **requires** this phase independence:

If any phase-dependent aspect of entangled state were locally observable, instant
information-containing messages could be sent, violating special relativity.

Local states of entangled subsystems must be invariant to phase changes.

Thus, only the relationship - the correlations -
between A and B, but not A or B themselves, can vary with phase angle.

Since local observers cannot detect these correlations,
the entangled state cannot be used to send superluminal signals.

This is, ultimately, the reason Schrodinger's cat
must be either alive or dead rather than a superposition of both.

A phase-dependent superposition involving both local states
would permit nonlocal signaling, violating relativity.

This conclusion implies that the standard physical description of composite non-entangled (i.e. factorable) product state such as $|\psi 1\rangle|1\rangle$ **has, for a long time, been mistaken**.

Usually we regard $|\psi 1\rangle|1\rangle$ as state of the composite system AB, where subsystem A is in state $|\psi_1\rangle$ and subsystem B is in state |1>.

But this leads us into the **paradox** of Schrodinger's cat, where (5)

$$(|\psi_1\rangle\,|1\rangle + |\psi_2\rangle\,|2\rangle)/\sqrt{2}$$

represents a state in which two macroscopically different composite states exist **simultaneously** as a superposition.

According to the present discussion, quantum theory and quantum experiments **imply** this entangled state to be a **superposition** of correlations between states rather than a superposition of composite states.

Thus $|\psi_1\rangle|1\rangle$ is **not** a state of composite system, but **instead** a correlation between two subsystems.

That is, $|\psi_1\rangle|1\rangle$ means **"subsystem A is in state $|\psi_1\rangle$ if and only if subsystem B is in state $|1\rangle$,"** an important departure from usual description.

Even if one of two subsystems happens to be macroscopic detector, entangled state (5) is simply a non-paradoxical superposition of correlations.

It says merely that state $|\psi_1\rangle$ of A is correlated with state $|1\rangle$ of B, and state $|\psi_2\rangle$ of A is correlated with the state $|2\rangle$ of B, with **non-local phase angle** $\phi_B - \phi_A$ determining degree of each correlation.

Regardless of the value of the phase angle, neither subsystem is in a superposition.

The entangled measurement state (5) is best described as a "macroscopic correlation":

a pair of superposed (i.e. phase-dependent) quantum correlations in which one subsystem happens to be macroscopic.

It is technically very difficult to create a macroscopic superposition, but macroscopic which-path detectors routinely achieve state (5).

It’s not paradoxical, even though many physicists have puzzled over it.

In entanglement, nature employs an ingenious tactic.

She must not violate relativistic causality, yet she must be nonlocal in order to maintain the pure-state nature of the original single-quantum superposition over composite objects such as bi-photons.

Thus she accomplishes nonlocality entirely via the superposition of correlations, because correlations cannot be locally detected and thus their superposition cannot violate relativity.

This tactic lies behind the nonlocal spread of phase-dependence over large spatial distances.

By means of the superposition of correlations - entanglement - nature creates a phase-dependent pure-state quantum structure across extended quantum systems such as bi-photons.

I’ve frequently used the term "local" as contrasted with "global."

For composite systems, and especially the entangled measurement state, it’s a crucial distinction.

Entangled states such as (5) have distinct local and global (nonlocal) aspects.

The local description corresponds to two observers, each observing only one subsystem.

In the case of (5), this "local description" is fully captured by the reduced density operators (10) and (11) - **each local observer detects a mixture, not a superposition, of one subsystem.**

The "global description" means the evolving pure state of the entire composite system, in our case Eq. (5).

It is a superposition of nonlocal correlations that can only be detected by observing both subsystems and, via an ensemble of trials that individually record corresponding outcomes at both subsystems, determining the state of the correlations between them.

Although the global state implies the local description, the local description cannot hint at the global correlations because any such hint would violate Einstein causality.

Thus, when an electron shows up in your lab, neither an examination of the electron nor an examination of an ensemble of identically-created electrons can give you the least hint of whether or how this electron is entangled with other quanta elsewhere in the universe.

This clarification of entanglement resolves the problem of definite outcomes, aka the Schrodinger's cat.

An ideal measurement of a superposed microscopic system A by a macroscopic detector B establishes the measurement state (5) at 100% positive correlation.

This state is equivalent to the logical conjunction "A is in local state $|\psi_1\rangle$ if and only if B is in local state $|1\rangle$, AND A is in local state $|\psi_2\rangle$ if and only if B is in local state $|2\rangle$," where AND indicates the superposition.

This conjunction is precisely what we want following a measurement.

Schrodinger's cat is not in the least paradoxical.

Still, this analysis **does not** entirely resolve the quantum measurement problem.

It resolves the problem of definite outcomes associated with the measurement state (5), but this state continues to obey Schrodinger's equation and is hence reversible.

In fact, the entangled state between a quantum and its which-path detector can actually be reversed in the Stern-Gerlach experiment.

In my view, a quantum measurement must result in a macroscopic indication such as a recorded mark, and a mark is **irreversible**.

The above analysis shows the entangled state (5)

describes a mixture of definite, not superposed, outcomes of measurements,

but these outcomes remain indeterminate and the global state remains reversible.

The irreversibility problem is the question of

how this nonlocal superposition of correlations then further “collapses” irreversibly to just one of its possible outcomes, a “collapse” that occurs in the RTO experiment only when one photon impacts a detector.

Present analysis does not seem to resolve this problem.

I think Gambler’s Ruin might do it!

In the case of the RTO experiment, however,

it seems fairly clear that the non-local superposition described by Eq. (5)

must irreversibly decohere when either of its subsystems A or B interacts with a detector.

The RTO experiment furnishes a particularly good setting for this question,

because the two photons remain in the reversible entangled state (5)

throughout their flights from the source to detectors,

and thus the two key questions of the measurement problem

(the problem of definite outcomes and the problem of irreversibility)

can be analyzed individually.

Let us now **attempt** to resolve the seeming existence of an "**irreversiblity**" problem.

The Environment as Monitor

I hope you are by now convinced that quantum physics describes the microscopic world entirely, consistently and with unparalleled experimental accuracy.

But still, there's one slight problem:

Quantum physics seems to fail utterly in describing the world around us!

We never see tables or teapots, not to mention ice cream, as wavy, space-filling fields that are possibly here and possibly there and possibly both here and there.

Tables and teapots don't quantum jump, nor does ice cream.

Classical physics may be inaccurate at the microscopic level, but it does seem to explain how such ordinary objects move in response to forces.

If quantum physics describes the microscopic world correctly, and if the macroscopic world is made of microscopic objects, then the quantum principles should lead ultimately to tables and teapots.

Let us now explain, at least in part, **one way that might work**.

As we have seen, quantum physics began with Max Planck's hypothesis that eventually led to the quantum.

The crucial quantum principle is the universe is made of these highly unified, extended **bundles** of field energy.

To the extent that any given phenomenon depends on the spatially extended field nature of quanta, the phenomenon should be considered as "quantum".

To the extent that quanta can be represented by “pointlike” objects, the phenomenon can usually be considered "classical".

But the existence of a classical-quantum boundary is a useful metaphor, not a law of nature.

Most physicists, me included, regard the world as fully quantum(as you will see later).

The clearest expression of the extended field nature of quanta is the principle of superposition.

Because electrons, photons, atoms, and so on are simply disturbances in fields, these objects superpose just as disturbances in the surface of a tub of water superpose.

Several states can all be present at the same time.

We will describe how the process called **decoherence** converts these superpositions into classical-like mixtures later.

The quantum measurement problem must be part and parcel of any discussion of how quantum physics explains our normal world.

After all, quantum measurements - quantum phenomena that cause macroscopic changes - are the bridge from the micro- to the macroworld.

Schrodinger's cat is a dramatic example.

The microscopic decay of a radioactive atom triggers a device that can kill a macroscopic cat.

Schrodinger launched the measurement problem when he noted that the quantum rules seem to imply something we never see: a macroscopic superposition - namely, a cat that is both alive and dead.

Earlier, we suggested a resolution of this problem of definite outcomes.

But there's more than that to the measurement problem.

Macroscopic processes are **irreversible**.

The moving finger moves on.

"Things run down".

The second law of thermodynamics demands it.

So quantum measurements must be irreversible, even though there appears to be no trace of irreversibility in the microscopic world.

As we'll see, decoherence **seems** to solve this mystery.

The Problem of Irreversibility

Every measurement involves a macroscopic change of some sort, and such changes must obey the second law of thermodynamics, so entropy increases.

Furthermore, we can observe this entropy increase as a permanent (i.e., irreversible) mark made by the measurement.

But when a quantum is measured, the quantum and its detector obey the time evolution equation as they **entangle** to form the measurement state, so it seems that entropy doesn't increase.


How can we resolve this conundrum?

A variation on the Stern-Gerlach experiment demonstrates the connection between quantum measurements and the second law.

I'll focus only on the features of this experiment relevant to the irreversibility of measurements.

Here's the experiment.

A horizontal stream of silver atoms passes between a pair of magnets (see figure).




The entering atoms, at the left, have all been prepared previously in identical states that I'll call the "zero state".

There's no need for us to concern ourselves with the precise nature of this initial state.

The magnets are shaped to create a so-called "inhomogeneous magnetic field" in the space between the magnets, as shown in the figure.

This field separates the stream in such a way that if a detection screen (dashed line) is placed "downstream" from the magnets as shown, atoms make **visible impacts** on the screen at two different spots, one above and the other below the original direction.

Individual atoms impact randomly at one or the other spot with a 50% probability - a striking example of quantum indeterminacy - because the atoms were all prepared identically.



Examination of the two beams shows that atoms striking the upper spot are no longer in the zero state but are in a different state called the "plus state", whereas the atoms striking the lower spot are in a new state called the "minus state".

So the magnet-plus-screen combination acts as a detector to determine which atoms are in the plus state and which are in the minus state. This is entirely analogous to a double-slit experiment with a which-slit detector putting quanta into the "slit 1 state" or the "slit 2 state" before the quanta strike a viewing screen.

There's more, however.

If one removes the detection screen and instead installs some appropriately chosen magnets (not shown in figure) at certain points along both paths, one can bend each stream back onto its original horizontal path, as shown.

Removing screen —> **No** measurements made now.

On studying the atoms in the converged stream, a perhaps surprising result emerges; none of these atoms are in the plus state and none are in the minus state.

Instead, every one is in the zero state from which it started!

We created a **"do-nothing" box**!

So this experiment, without the detection screen(without the measurement), **is** reversible.

But with the screen, the experiment is obviously **irreversible** because the atoms make an **irreversible** mark when they strike the screen.


It follows that any irreversibility and corresponding entropy increase in this experiment are **entirely due** to the impact on the screen, because we have seen that, without the screen, the experiment is reversible.

Thus the macroscopic detection makes all the difference, and it is here that we must search for all the irreversible effects of measurement.

Specifically, mere entanglement, such as occurs when a which-slit detector operates in the double-slit experiment, is **not responsible** for irreversibility.

The detection at the screen works as follows:

The detection screen interacts with each atom to form the entangled measurement state: atom is in the plus state, flash appears at the upper spot entangled (superposed) with atom is in the minus state, flash appears at the lower spot



This is the controversial Schrodinger's cat state that we analyzed earlier (the radioactive nucleus is now the silver atom, and the cat is now the screen), where we argued that this state is only a superposition of correlations.

This experiment demonstrates that the macroscopic recording of a measurement is no small detail.

No measurement is complete until it makes a mark on the macroscopic world, and making such a mark must, because of the second law, involve an entropy increase, implying irreversibility.

As John Wheeler repeatedly stressed,

> No elementary quantum phenomenon is a phenomenon until it is brought to a close by an irreversible act of **amplification(so we can see it!!)**

If the measurement is recorded by the audible click of a detector, for example, the irreversible mark is a sound wave spreading out in all directions into the air around the detector.

This sound wave warms the air a little and eventually disperses (vanishes, **for all practical purposes**) into a large volume of air.

There is no way nature is going to spontaneously gather up every last bit of this wave's dispersed energy (while necessarily cooling the air), reverse the entire dispersal process, and use this energy to restore the detector and the air to their states before the click.

In fact, the reversed process is prohibited by the second law because it would reduce the total entropy of the universe!

We conclude that microscopic quantum processes, including entanglement, remain reversible as long as they remain microscopic, and that the irreversibility of measurements must be rooted in the macroscopic recording process.

We now look at a macroscopic recording process that goes on all over the universe all the time.

How Environmental Decoherence Collapses Superpositions

A pebble lies on a sunny beach, immersed in an environment that includes atmospheric molecules, photons from the sun, cosmic rays from stars, and even photons from the Big Bang.

During every second, many such quanta interact with the pebble, reflecting or otherwise scattering off the pebble in every possible direction.

The scattered photons must carry away data about the pebble's orientation, structure, and color, because otherwise the pebble could not be seen.

Such natural "measurement" processes occur all the time, regardless of the presence or absence of humans to consciously observe the gathered data.

To study the quantum features of such a natural measurement, instead of a pebble let's consider an atmospheric atom that is in a highly nonclassical state of being superposed at two or more macroscopically separated locations.

We know that such a superposition can be created in the laboratory, for instance by passing the atom through a double-slit apparatus, but it could also occur naturally, for example if the atom passes through an opening that is sufficiently narrow to cause the atom to diffract widely.

What happens when such an atom interacts with quanta in the surrounding environment?

The superposed atom interacts with, say, an environmental photon in a manner analogous to the way a superposed electron coming through double slits interacts with a which-path detector.

If the interaction is significantly different at the atom's two superposed locations, a photon can "measure" the atom by entangling with it in a measurement-like state.

Such an entangling interaction is just like a which-path measurement as discussed earlier!

The scattered photon carries away which-path(location) data about the atom, just as a which-path detector carries away data about an electron coming through the slits.

As discussed earlier(via reduced density operator), this measurement collapses or decoheres (removes, via a series of small environmental interactions, the interference pattern from) the superposed atom, converting its state from a superposition of being at both locations to a local mixture of being either at the first location or at the second location.

In the natural environment, a single scattered photon is not likely to entirely measure (entirely decohere) the atom, because environmental quanta are not specifically organized, the way a laboratory detector is organized, to measure superpositions.

A which-path detector in a laboratory is carefully constructed to respond in detectably different ways at different locations of the measured quantum, quite unlike environmental quanta, which interact randomly in all kinds of ways.

Although a good which-path detector requires only one interaction to reliably distinguish between - reliably decohere - the superposed states of the detected quantum, a large number of environmental quanta must scatter from a typical superposed atom to completely decohere it and turn it into a local mixture.

Careful analysis shows that decoherence of typical naturally occurring superpositions requires many environmental interactions.

So decoherence by the environment **involves** a series of partial measurements and partial collapses, **each** of them instantaneous, nonlocal, and similar to the single-step measurement that occurs at the slits in a double-slit experiment with a which-path detector.

It's through this environmental decoherence process that everyday objects such as pebbles and this slide lose their extended quantum field nature and behave classically, with no obvious trace of superposition, interference, or nonlocality.

Small objects, of atomic dimensions, are less susceptible to environmental decoherence simply because fewer environmental quanta scatter from them as a result of their smaller size, and because each scattering event can cause only a tiny amount of decoherence because the entanglement involves nearly indistinguishable locations.

Thus, a superposed photon from the Big Bang might travel the universe for 13.8 billion years without decohering, while a grain of sand on Earth, should it happen to show any signs of superposition, is decohered environmentally and essentially instantly because of the myriad environmental interactions it experiences.

Widely extended superpositions of small objects, such as a fine dust grain superposed in two locations separated by a millimeter, are also decohered nearly instantly because the branches of such a superposition are so distinct that a single photon reflecting from one branch but not the other can turn the entire superposition into a mixture.

In a similar way, a single detector at only one of two parallel slits is sufficient to turn a quantum that's initially in a superposition of coming through both slits into a mixture that comes through either one or the other slit.

For mesoscopic objects such as dust grains, environmental decoherence turns superpositions quickly into mixtures.

How quick?

This has been calculated theoretically and measured in a few experiments.

A fine dust grain is some 10^{-5} m across.

If it's in a superposition of being in two places with its two superposed branches separated by this same distance so that the two branches are right next to each other, as though the grain had come through two closely adjacent slits, it would be decohered entirely by normal air on Earth in only 10^{-31} second.

Even the best laboratory vacuum (which still contains plenty of air molecules) would decohere it within 10^{-14} second - a hundredth of a trillionth of a second.

If this grain were in deep outer space, cosmic background radiation from the Big Bang would decohere it in 1 second.

For a smaller object, such as a large molecule with a diameter of 10^{-8} m, these decoherence times are longer (still assuming it's in a superposition with two branches that are separated by the molecule's diameter): 10^{-19} second if the molecule is in normal air, 0.01 second if it's in the best laboratory vacuum, but much longer than the age of the universe if it's in deep space.

The message is that sufficiently small superpositions can survive awhile, but meso- or macroscopic superpositions are fragile and are decohered quickly by tiny environmental interactions.

This is why the "quantum world" is usually identified with the microworld.

When typical quantum features become meso- or macroscopic, they generally vanish quickly.

So the quantum universe appears classical at the macroscopic level because the enveloping quantum environment "monitors" every object constantly, and macroscopic objects are especially susceptible to this decoherence process.

Nature is full of which-branch detectors! Note that humans aren’t required in any of this - no physicists, no laboratories needed.

Nature has been collapsing superpositions, and quanta have been losing their quantum field nature, at least since the Big Bang.

The role of the environment as an ever-present which-path monitor that turns mesoscopic and macroscopic superpositions into mixtures was first clearly recognized by Zurek during the early 1980s.

The work of Zurek, his colleagues, and others has, in large part, explained how the quantum world leads(via reduced density operators) to the classical world of our experience.

A wide variety of experiments have demonstrated environmental decoherence and convinced physicists that decoherence really is the mechanism that converts the quantum world into tables and teapots.

One beautiful example is an experiment by the University of Vienna group under Zeilinger and Arndt.


They use the Talbot-Lau interferometer technique (see next slide) that demonstrated, in 2002, interference in large molecules such as C70 and certain biological molecules.


The interference showed each of these large molecules to be in a superposition of following more than one path on its journey through the interferometer.

Using the same technique in 2004, this group was able to demonstrate convincingly the "environmental" decoherence of superpositions of C70 molecules.

I put environmental in quotation marks because in this experiment the environment came from within the molecules themselves rather than from an external environment.

Individual C70 molecules passed through the interferometer, as in the 2002 experiment.


detector
ionisation laser
3. grating
2. grating
1. grating
lateral limiter
height limiter
fullerene source
gold gratings
period 1 µm
5000
counts in 5 s
0
0
1
2
3
shift [µm]


Analyzer
Detector
few µm
Beam-splitter
Source grating
X-ray source

But there was a new twist; the experimenters heated the molecules just before sending them through the interferometer.

They expected that, with sufficient heating, the molecules would themselves emit thermal radiation in the form of visible and infrared "thermal photons", just as an electric hot plate emits thermal radiation (you can feel its warmth at a distance, and it might glow red) when heated.

The radiation comes from the random thermal motion of the many atoms and other quanta within the molecule.

According to decoherence theory, each of these radiated photons should act as a partial which-path detector, collapsing the molecule’s superposition state partially by carrying a certain amount of which-path data from the molecule into its surroundings.

If enough such data are transferred, this which-path measurement should decohere the molecule, causing the interference pattern to dim or vanish - the signature of a superposition evolving into a mixture.

Their results demonstrated decoherence in action.

With the molecules only slightly heated, the emitted photons had low energies and thus low frequencies and long wavelengths - too long to distinguish between the possible paths, which for such a massive molecule are separated by extremely small distances.

But when the molecules were heated to a few thousand kelvins, the emitted photons' wavelengths became short enough to distinguish between the possible paths, so the which-path data transmitted to the surroundings was sufficient to partially decohere the superposed molecules, causing the interference pattern to partially vanish.

At sufficiently high temperatures, theory and experiment showed that a mere two or three emitted photons sufficed to decohere each molecule.

There was quantitative agreement between the theory predictions and observations.

The interference pattern began decohering at just the predicted temperature, and the degree of visibility of the remaining interference was just as predicted.

The experiment revealed the step-by-step action of decoherence as data leaked, photon by photon, into the environment.

It also demonstrated the extreme sensitivity of superpositions to decoherence.

Just a few high-temperature photons turn a massive superposed molecule into an incoherent mixture.

Superposed quanta are very delicate.

Decoherence and the Measurement Problem

Decoherence provides a solution of the irreversibility problem for **natural** environmental measurements.

Decoherence was first introduced during the early 1980s to explain how our apparently classical surroundings arise from measurement-like interactions with the environment.

The local state argument, presented earlier, shows that such measurements turn superpositions into entangled nonlocal measurement states with definite outcomes, resolving the problem of definite outcomes for environmental measurements.

But just as for laboratory measurements, we must ask if this also resolves the irreversibility problem associated with **natural** measurements.

The answer is yes.

Here's why.

As we've seen, when a small superposed dust grain is decohered by the surrounding environment, the object undergoes a series of small entanglement-caused partial collapses.

These environmental measurements are "recorded" by the many environmental photons and air molecules that interact with the grain and then disperse widely.

Data about the superposition state of the grain before it decohered are now scattered randomly far and wide.

Just as one cannot unscramble an egg, one cannot reversibly gather these pieces back together and reconstruct the global state of the air plus the superposed grain.

This argument is especially compelling in the case of environmental decoherence, because of the environment's enormous size.

This is an obvious example of the second law in action; **for all practical purposes**, the process is irreversible and entropy has increased.

Because there is famous opposition to introducing **for-all-practical-purposes** arguments into physics, it needs to be noted that the second law itself is inherently a **for-all-practical-purposes** principle.

We know that a box full of gas could evolve spontaneously into separate regions of hot gas and cold gas simply by chance, with no external assistance.

The chances of this are ridiculously small, but they are not zero, and if we consider boxes containing smaller and smaller amounts of gas, these odds increase.

Such **for-all-practical-purposes** arguments that trace back to the second law are entirely in keeping with the principles of physics.

So the local state resolution plus decoherence combine to resolve the measurement problem entirely for the case of natural environmental measurements.

The environment measures a superposed object (molecule, dust grain, and so on) when myriad environmental quanta convert the superposition into a measurement state that now takes the form of a nonlocal entanglement between the object and a very dispersed state of the many environmental quanta.

The local state argument implies this measurement state represents a mixture of definite properties, whereas the highly dispersed nature of the environmental quanta guarantees the process is irreversible.

There has, for years, been a question about what is the exact relationship between decoherence and the solution of the measurement problem.

Some accounts appear to imply that decoherence alone resolves the measurement problem, but this is not true.

The connection of decoherence to the measurement problem is that it resolves the irreversibility problem in the case of natural environmental measurements, and it shows how environmental interactions transform a superposed quantum into the measurement state, but it does not solve the problem of definite outcomes associated with this measurement state.

The problem of definite outcomes is, however, resolved by the local state analysis, as explained earlier.

So the local state solution combines with environmental decoherence to solve the measurement problem for the case of environmental measurements.

What about the case of laboratory measurements?

Here, the preceding discussion resolved the definite outcomes problem, but not the irreversibility problem.

Irreversibility poses slightly different problems in the two cases, because lab measurements are sufficiently controlled that the natural environment has little effect(by design).


In fact, much of an experimentalist's efforts go precisely into ensuring that random environmental interactions have no significant effect on an experiment's outcome.

Our discussion of the Stern-Gerlach experiment suggested that the answer lies in the irreversible nature of the macroscopic detection process.

This suggestion resembles the resolution of the irreversibility problem for environmental measurements:

Environmental measurements are "recorded" by innumerable environmental quanta whereas lab measurements are recorded by detection screens, electronic clicks, or, perhaps, cats.

Let's focus on a single impact made on a detection screen by one electron in a double-slit experiment - one of the small spots in the figure below:



(a–e) Buildup of an interference pattern in the double-slit experiment using electrons.

As in a regular interference experiment, an interference pattern builds up from individual impacts over longer-time exposures.

How was it recorded?

This mark was initiated by a single electron, but to observe it macroscopically the original impact had to be detected and then **amplified** sufficiently for humans to see it.

This is the purpose of every laboratory detector of microscopic events, and it's why an **"irreversible act of amplification"** (as Wheeler puts it) is essential to quantum measurements.

In the 1989 experiment that produced the patterns of dots on a screen, each electron was emitted in a single coherent state, the same state for each electron.

EM fields accelerated these electrons to high speeds, about 180,000 kilometers per second, or 60% of light speed - much faster than the electron in a normal hydrogen atom, which orbits at an average speed of "only" 2000 kilometers per second.

High energies were needed to make a detectable impact.

Each electron went through a pair of parallel slits, then spread out into an interference state that interacted indeterminately with the screen.

The interaction entangled the electron with the screen, decohering the electron and collapsing it instantaneously from its earlier superposition state over the entire screen to a locally observed mixture(a flash).

In other words, the electron collapsed randomly into one or the other of many small atom-size regions in the screen.

For the fluorescent screen used in the experiment, this interaction created some 500 photons that marked and amplified the location of the impact - already an irreversible process because the photons are emitted randomly in all directions.

In a process called the photoelectric effect, each photon then struck a metal plate and dislodged an electron from it, converting each of the 500 photons into a low-energy electron.

These 500 low-energy electrons were then focused into a point image that could be displayed on a TV monitor in much the same way that old (before flat screens) TV tubes operated.

Every step of this process - creation of 500 photons, conversion to photoelectrons, and amplification to create the final display - creates entropy and is irreversible thermodynamically.

It's the second law of thermodynamics in action.

This illustrates the general case:

Laboratory detection and amplification of a quantum event necessarily involves irreversibility and entropy production.

The laboratory detector, a macroscopic device made of many microscopic quanta, plays the decohering role that the environment plays in natural measurements.

To summarize:

The quantum measurement problem, in both its laboratory and environmental senses, seems to be resolved entirely by the local state solution of the problem of definite outcomes and by the irreversibility of the decoherence process.

For more than a century, much has been made of the odd and supposedly paradoxical nature of the quantum.

This presumed quantum spookiness has led to an excess of attempted fixes and interpretations.

Many experts have even declared the theory to be not a description of reality at all, but only a mathematical recipe that helps humans predict the results of experiments.

As Niels Bohr put it, **“There is no quantum world. There is only an abstract quantum description.”**

According to this hypothesis, quantum theory doesn’t describe anything real at all, so there’s no cause for concern about collapse of the quantum state and other odd quantum behaviors.

We now collect everything we have proposed together, eliminate all that is unneeded and finally present a straightforward picture of the solution to the quantum measurement process

We will see that the final understanding arises because we looked at so many different approaches, went down so many dead-ends and along the way uncovered many new features of the quantum world.

So, now we will present only what we need with a more detailed analysis and a deeper and, what I believe, is a correct interpretation.

1 - INTRODUCTION

Physicists agree that Schrodinger's equation describes the unitary evolution of non-relativistic quantum states between measurements, but there is no agreement on how states change during measurements.

In fact, an apparent problem arises when one applies standard quantum theory (minus the collapse postulate) to measurements.

John von Neumann, in 1932, analyzed the problem and we follow his argument here.

> "Measurement" is the **experimental determination and macroscopic recording** of the value of a physical observable associated with a quantum system.
>
> von Neumann showed that, **unless** the system happens to be in an eigenstate of the measured observable, measurement leads to a **"measurement state" (MS)** whose mathematical representation is an **entangled state** that **seems to predict** the detector to be in a **macroscopic superposition** of exhibiting all of the possible outcomes, a paradox known as the **"problem of definite outcomes."**

We will now show, based on standard quantum theory without a collapse postulate, that this is a **pseudo-problem** and that, far from predicting superposed outcomes, von Neumann's MS **predicts an instantaneous collapse** to a single eigenstate.

Specifically, we will demonstrate, with no assumptions other than standard quantum physics (minus the collapse postulate)

(i) The MS has been **misinterpreted** (for 90 years) and **does not** in fact predict paradoxical multiple macroscopic *outcomes;* it instead correctly predicts non-paradoxical multiple ***statistical correlations between*** system and detector outcomes.

(ii) Exactly **one** outcome actually occurs.

(iii) The entanglement says that, when one outcome occurs, the **other outcomes** simultaneously and nonlocally remain "dark" (i.e. **do not occur**).

This **resolves** an objection to quantum physics first raised by Einstein in 1927.

That is, we will show von Neumann's enigmatic MS **to be in fact** the collapsed state expected upon measurement.

The collapse is derived (i.e. demonstrated) with **no assumptions** beyond the other standard principles of quantum physics(i.e., without a collapse postulate!).

We will show the problem of outcomes arises from a **mistaken understanding** of entangled superpositions, **not only** in measurements **but also** in purely microscopic processes.

The MS is obviously a **superposition** of subsystem product states.

But the MS is poorly understood because no previous work has analyzed its complete **phase dependence**. Remember our earlier discussion of state vectors for boxes of electrons!

Do the states of individual subsystems vary with phase, as they do in simple superpositions?

If not, then what does vary with phase, i.e., precisely which entities occupy indefinite states when two or more subsystems are entangled?

Such questions show that we do not fully understand the MS's phase dependence, i.e., we don't fully understand the MS.

We investigate these questions by studying earlier Bell-test quantum-optics experiments that measure momentum-entangled two-photon states.

These experiments study a **purely microscopic** entangled superposition (mathematically **identical** with the MS) **across all phases**.

The results show that entangled superpositions **differ sharply** from simple superpositions, i.e., the implication of the "**plus**" sign differs surprisingly.

The MS **does not** represent multiple detector readings, but instead represents multiple ***statistical correlations between*** system states and detector readings.

These correlations are **not paradoxical**: a macroscopic detector that simultaneously exhibited two states **would be paradoxical**, but a detector that simultaneously participates in two correlations is **not paradoxical**.

Furthermore, the MS **directly implies** that **exactly one** of these correlations is **realized** as the measurement **outcome**.

This **resolves** the so-called "Schrodinger's cat paradox," which in turn **resolves** the quantum measurement problem.

Our discussion follows from quantum theory and from experimental evidence(already discussed) that an instantaneous nonlocal collapse takes place, **resulting in one outcome occurring while the other outcomes simultaneously do not occur**.

Thus, while the present discussion **does not *postulate* collapse**, it ***derives an instantaneous collapse*** as a **consequence** of **entanglement**.

Many papers, and our earlier discussions, express the effect of measurement as a **trace** on the composite-system density operator representing the MS.

This trace operation, which **predicts definite outcomes** at the subsystems, seems to yield just what we want, namely that measurement **transforms** the subsystem states into **mixtures** over the possible eigenstates.

However, **several** objections are commonly raised against this proposal, so we now go into more detail to understand what is the **correct** interpretation that overcomes objections.

This discussion, in contrast to earlier discussions, argues that the MS **directly** represents the collapsed state of the composite system, that the collapse is a **consequence** of entanglement, and that the entanglement is **required** in order to ensure a **simultaneous** (hence instantaneous) collapse over the separated branches of the experiment.

We will **present** von Neumann's derivation of the MS, **pose** the measurement problem, **review** eight presumed measurement problem insolubility proofs, and **explain** why their conclusion cannot be correct.

We then present a **crucial** clue — In 1927, **Einstein** noted that quantum theory **implies** that measurements entail **instantaneous** collapse and suggested this would **violate** Special Relativity.

However, today (unlike 1927) we **know** that **instantaneous** nonlocal changes of **correlations** violating Bell's inequalities really **occur**, that they **do not violate** special relativity, and that they occur when disparate systems are **entangled**.

They have been observed experimentally!

Thus, from today's perspective, Einstein's **argument** implies that **nonlocal correlations** are **required** during measurements and also saying that **entanglement** is also **required**.

Is von Neumann's MS in fact precisely what we **want**?

The answer can only come from **fully understanding** the MS, particularly its **full phase dependence**.

To this end, we will review two 1990 quantum optics experiments exploring an entangled two-photon microscopic state that is **mathematically identical** to the MS, that we discussed earlier.

The **correct** understanding of the microscopic version of the MS will then be worked out.

We **find** that both entangled sub-systems are in **definite** (i.e. non-superposed) states, but that the ***degree of correlation between these states is in an indefinite state***.

We will then apply this new **insight** to the MS.

We will find the MS is **not** a macroscopic superposition of different detector *states*.

It instead **represents** *different correlations between* detector states and system states.

This is **not paradoxical**.

Furthermore, quantum theory directly implies that precisely **one outcome is realized**.

This **resolves** the Schrodinger's cat paradox.

We will then conclude the analysis by studying a particularly **simple measurement example** that also typifies the **essentials** of quantum measurements.

Finally we will summarize the results and consider **why it has taken so long to overcome this simple misunderstanding of von Neumann's MS**.

2 - ENTANGLEMENT AND MEASUREMENT

The superposition postulate entails that, if $|A1\rangle$ and $|A2\rangle$ are Hilbert space vectors ("kets") representing possible states of a quantum system *A*, then all normalized linear superpositions of $|A1\rangle$ and $|A2\rangle$ also represent possible states of *A*. For example,

$$|\Psi_A\rangle = \frac{1}{\sqrt{2}}(|A1\rangle + |A2\rangle) \qquad (1)$$

You should make a copy of all the numbered equations for reference as I lecture

($|A1\rangle$ and $|A2\rangle$ are orthonormal) represents a possible state of *A*.

The superposition postulate is prerequisite to the Hilbert-space representation of quantum states, and the basis for conceptualizing quantum states as physically real waves in a quantum field that fills the universe.

$|\Psi_A\rangle$ represents a situation in which *A* is represented neither by $|A1\rangle$ nor by $|A2\rangle$ but incorporates aspects of both, including "overlap" effects such as interference.

As **Dirac** put it, *A* goes "partly into each of the two components" and "then interferes only with itself."

If quantum system *A* interacts with another quantum system *B*, it frequently happens that the situation of *A* and *B* are then represented by an entangled superposition state vector such as

$$|\Psi_{AB}\rangle = \frac{1}{\sqrt{2}}(|A1\rangle\,|B1\rangle + |A2\rangle\,|B2\rangle) \qquad (2)$$

where $|Ai\rangle$ and $|Bi\rangle$ (*i = 1, 2*) are orthonormal kets representing the "subsystems" *A* and *B*, respectively.

Although the physical interpretation of simple superpositions such as $|\Psi_A\rangle$ is clear, the physical interpretation of entangled superpositions such as $|\Psi_{AB}\rangle$ is not comparably clear.

$|\Psi_{AB}\rangle$ is a superposition of two products $|A_i\rangle\,|B_i\rangle$ (*i =1, 2*). $|A_1\rangle\,|B_1\rangle$ is commonly interpreted to represent a state of the composite system *AB* in which *A* has the properties associated with $|A1\rangle$ and *B* has the properties associated with $|B1\rangle$, and similarly for $|A2\rangle\,|B2\rangle$.

But if this is the case, then the physical interpretation of the state $|\Psi_{AB}\rangle$ would seem to be that *AB* simultaneously exhibits properties associated with $|A1\rangle$ and $|B1\rangle$ AND properties associated with $|A2\rangle$ and $|B2\rangle$, where "AND" represents the superposition.

In the case of Schrodinger's iconic cat, this would imply that the nucleus is both decayed and undecayed and, more disturbingly, the cat is both alive and dead.

Here we will demonstrate that both quantum experiment and quantum theory show that this is actually *not* the case.

Instead, the state $|\Psi_{AB}\rangle$ entails merely that $|A1\rangle$ and $|B1\rangle$ are *coherently* (in a phase-dependent manner) *correlated* with each other AND $|A2\rangle$ and $|B2\rangle$ are coherently correlated with each other (see later discussion).

This is not paradoxical in any way.

Quantum measurements are important examples of entanglement.

As first discussed by John von Neumann, a "measurement" is the determination of the value of an observable associated with a quantum system *A*.

If *A* happens to be represented by an eigenvector of the measured observable, then a good measurement will detect the associated eigenvalue.

But what if *A* is represented by a superposition of eigenvectors of the measured observable?

It will suffice for this discussion's purpose to assume that *A*'s Hilbert space has only two dimensions, and that *A* is represented by the superposition Equation (1).

The kets $|Ai\rangle$ *(i =1, 2)* define the eigenvectors of the measured observable.

We assume the existence of a **detector** *B* designed to **distinguish** between the $|Ai\rangle$.

For example, $|A1\rangle$ and $|A2\rangle$ could represent the paths of an electron passing through the slits of a double-slit apparatus, and *B* could be an electron detector for the "which-slit" observable whose eigenvectors are the $|Ai\rangle$.

To make a which-slit measurement, *B* must distinguish between the states represented by $|A1\rangle$ and $|A2\rangle$, so *B* must contain a specific quantum detection component with quantum states represented by kets $|Bi\rangle$ such that, if *A* is in the state represented by $|Ai\rangle$, then detection yields the state represented by $|Bi\rangle$*(i = 1 or 2)*.

Assuming a minimally-disturbing measurement that leaves eigenstates unaltered, and letting $|B_{ready}\rangle$ represent the state of *B*'s quantum component prior to measurement, the process

$$|Ai\rangle\,|B_{ready}\rangle \Rightarrow |Ai\rangle\,|Bi\rangle \quad (i=1,2) \tag{3}$$

describes a measurement of the which-slit observable when *A*'s state is represented by an eigenstate.

When *A* is in the state represented by $|\Psi_A\rangle$ and *B* measures the which-slit observable, simple linearity of the time evolution implies

$$\frac{1}{\sqrt{2}}(|A1\rangle + |A2\rangle)\,|B_{ready}\rangle \Rightarrow \frac{1}{\sqrt{2}}(|A1\rangle\,|B1\rangle + |A2\rangle\,|B2\rangle) = |\Psi_{AB}\rangle \tag{4}$$

Thus von Neumann's straightforward argument shows the measurement creates the entangled superposition $|\Psi_{AB}\rangle$ of Equation (2), where "*B*" now refers to the quantum detection component of the detector.

But does von Neumann's measurement postulate imply that, when the which-slit observable is measured, *A* collapses into one of its eigenstates while *B* collapses into the corresponding detector state.

It is by no means clear that $|\Psi_{AB}\rangle$ (Equation (4)) represents such a measurement outcome.

As many physicists put it, "The problem of what to make of this is called the 'measurement problem'."

This discussion will show that $|\Psi_{AB}\rangle$ does in fact represent the collapsed state and the single definite outcome expected from von Neumann's measurement postulate.

We have used the same notation, $|\Psi_{AB}\rangle$, for the arbitrary entangled state represented by Equation (2) (where *A* and *B* are arbitrary quantum systems) and for the specific case of the entangled state that develops when a detector measures a quantum system, represented by Equation (4) (where *B* is now a detector).

We will refer to this state in the context of Equation (4) as the "measurement state" (MS).

We will also, however, need to refer to the arbitrary entangled state Equation (2), especially later where we analyze an experiment involving two microscopically entangled photons.

Thus the question of how to interpret entangled states looms large in the foundations of quantum physics.

As noted above, the interpretation of general entangled states such as the one represented by Equation (2) is already murky as compared with the interpretation of simple superposition states such as the one represented by Equation (1).

The problem of interpreting the MS is especially important, because **macroscopically distinct** states now lie on each side of the "plus" sign on the right-hand side of Equation (4).

As already discussed, the superposition $|\Psi_A\rangle$ can be interpreted to represent a situation in which *A* incorporates properties represented by both $|A1\rangle$ and $|A2\rangle$.

And a product state such as $|A1\rangle\,|B1\rangle$ represents a state of the composite system *AB* in which *A* is represented by $|A1\rangle$ and *B* is represented by $|B1\rangle$.

Thus $|\Psi_{AB}\rangle$ appears to describe a detector that simultaneously "points" to two macroscopically different outcomes $|B1\rangle$ and $|B2\rangle$!

The detector seems to display no definite outcome, a conundrum known as the "**problem of outcomes**"

Such a superposition state would be paradoxical.

Schrodinger, who imagined a cat attached to the detector in such a way that $|B1\rangle$ included a live cat and $|B2\rangle$ included a dead cat, described $|\Psi_{AB}\rangle$ as representing a “living and dead cat ...smeared out in equal parts.

As one quantum foundations expert writes,

The crucial difficulty is now that it is not at all obvious how one is to regard the dynamical evolution described by [Equation (4)] as representing measurement in the usual sense.

This is so because [Equation (4)] is ...not sufficient to directly conclude that the measurement has actually been completed.

In fact, while measurement should lead to a specific eigenstate of the measured observable, Equation (4) appears to entail that the system has been **sucked into a vortex of entanglement** and **no longer** has its own quantum state.

On top of that, the entangled state **fails to indicate** any particular measurement outcome.

As noted, it seems paradoxical that quantum measurements lead to a state represented by the MS.

Measurement should lead to a situation in which *A* is represented by one of its eigenvectors $|Ai\rangle$ and *B* is represented by the corresponding $|Bi\rangle$.

Since quantum uncertainty typically implies **unpredictable** measurement outcomes, it is reasonable to conclude that measurement should lead to a state represented by an **ignorance-interpretable mixture** of the products $|A1\rangle|B1\rangle$ and $|A2\rangle|B2\rangle$.

Assuming the initial state is represented by $|\Psi_A\rangle$, such a post-measurement mixture would be represented by the density operator

$$\rho_{mixed} = \frac{1}{2}(|A1\rangle|B1\rangle\langle B1|\langle A1| + |A2\rangle|B2\rangle\langle B2|\langle A2|) \tag{5}$$

This mixture can be interpreted(as we did earlier) as **"the system is represented by a single component** $|Ai\rangle|Bi\rangle$ **, but we cannot know whether *i = 1* or *2* until we look at the outcome."**

Beginning with von Neumann's analysis, at least eight published **"measurement problem insolubility proofs"** have assumed that, in order to obtain definite outcomes, the measurement process should transform the composite system *AB* into a mixture analogous to Equation (5) **(so have we so far!!)**

The initial state of *A* is assumed to be pure and to be represented by a superposition such as Equation (1).

The analysis then investigates whether a suitable composite-system post-measurement mixture can be reached via a unitary process.

To achieve this, the detector must be represented by a mixture initially, because unitary processes **cannot turn a pure state into a mixture**.

Since *B* is macroscopic, such an initial mixture **seems appropriate**.

Thus von Neuman and seven succeeding analysts asked:

> Is there an initial mixed-state density operator ρ_{ready} of *B* and a unitary process *U* acting on *AB* such that *U* transforms the initial composite density operator $|\Psi_A\rangle\langle\psi_A| \otimes \rho_{ready}$ into the desired composite mixture?

The eight insolubility proofs, with varying assumptions, say the answer is "no," presumably demonstrating the measurement problem to be insoluble(according to them!).

We will show later, however, that the premise of these insolubility proofs, namely that Equation (5) represents the appropriate post-measurement state, was doomed from the start, precisely because it is *not* entangled and thus cannot have the properties required if quantum theory is to describe the measurement process.

That statement represents the crucial point!!

To put this another way, there are reasons why the post-measurement state *must* be an entangled state, which implies that it *cannot* be a mixture such as Equation (5), which just represents what we measure!

We will then show that the MS does have the desired properties.

3 - A CRUCIAL CLUE FROM EINSTEIN

At the 1927 Solvay Conference, five years prior to von Neumann's analysis of quantum measurement, Einstein asked the audience to consider an experiment in which electrons pass through a tiny hole in an opaque screen and then impact a large hemispherical detection screen centered at the hole (Fig. 1).




Figure 1. Einstein's thought experiment. Each electron diffracts widely, reaching the entire screen simultaneously, yet only one point shows an impact. How do the other points *instantaneously* remain dark? Does this violate Special Relativity?

According to the Schrodinger equation, each electron's state diffracts widely, spreading and reaching the entire screen simultaneously.

Yet each electron registers at only a **single point**.

How, Einstein asked, do the other points ***instantaneously*** **remain "dark," i.e., not show an impact**?

As Einstein put it in his notes, this "**entirely peculiar mechanism of action-at-a-distance, which prevents the wave continuously distributed in space from producing an effect in *two* places on the screen,**" presents a fundamental problem.

It appears to imply instant signaling, **violating** special relativity.

Einstein's argument shows that, under a **realistic and objective** (**independent of humans**) interpretation of quantum physics, the Schrodinger equation is **at odds** with experimental facts **unless** the electron's state collapses instantaneously and nonlocally upon measurement.

Thus, realistic quantum physics **implies** instantaneously-established nonlocal correlations are **essential** to the *measurement process*.

Indeed, modern experiments have **verified the nonlocal character** of the measurement transition.

Since nonlocality is essential to measurement, the presumed post-measurement mixed state(5) was doomed from the start precisely because it **does *not* exhibit** the required nonlocality.

But entangled superpositions **do exhibit** the required nonlocality.

So from our **modern point of view**, Einstein's argument shows that **entanglement, far from being an unwelcome paradox, is *required* in measurements.**

This is a **crucial clue** and good news for quantum foundations, because von Neumann's predicted MS is **just such an entangled superposition**!

One particular argument, at least, should be mentioned. Its resolution of the measurement problem "diagonalizes the density operator" as we discussed earlier on one of our adventures.

They form the exact density operator $\rho = |\Psi_{AB}\rangle \langle \Psi_{AB}|$, which can be written

$$|\Psi_{AB}\rangle\langle\Psi_{AB}| = \rho_{\text{diagonal}} + \rho_{\text{off-diagonal}} \tag{6}$$

where $\rho_{diagonal} = \rho_{mixed}$ (Equation (5)) and

$$\rho_{off-diagonal} = \frac{(|A1\rangle|B1\rangle\langle B2|\langle A2| + |A2\rangle|B2\rangle\langle B1|\langle A1|)}{2}. \tag{7}$$

Recall that the exact expectation value of any observable F is

$$\langle F\rangle = \mathrm{Tr}(\rho F) = \sum_j \sum_k \rho_{jk} F_{kj} \tag{8}$$

where ρ_{ijk} and F_{ijk} are matrix elements of ρ and F .

This idea argues that off-diagonal terms in Equation (8) can be ignored because they involve matrix elements such as $\langle B1|\langle A1|F|A2\rangle|B2\rangle$ that are non-zero only for a **"fantastic"** observable F because $|B1\rangle$ and $|B2\rangle$ represent the states of **widely separated detectors**.

It says that matrix elements for such **fantastic** observables can, **for all practical purposes**, be neglected so that we can replace ρ by ρ_{mixed}.

But we have seen that this premise is doomed because ρ_{mixed} lacks the required nonlocal properties, so this type of proposal fails (it leaves our original discussions in limbo for the moment also!!).

4 - EXPERIMENTAL STUDIES OF STATES HAVING ENTANGLED SPATIAL PATHS

Earlier we presented the measurement problem and some previous thoughts on the problem.

We now will present a suggested resolution of all difficulties.

We now review interferometry experiments and theory that investigate the *microscopic* entangled superposition $|\Psi_{AB}\rangle$ Equation (2) over its full 0 - to - π range of phases, for a system of two momentum-entangled (i.e. path-entangled) photons.

The results provide a key insight into solving the measurement puzzle.

As preparation, we first study the simple superposition Equation (1).

Consider the interferometer experiment of Figure 2.

On each experimental trial, a photon enters a 50-50 beam splitter *BS1* which transforms the photon's state into the superposition Equation (1) where $|A1\rangle$ and $|A2\rangle$ respectively represent paths 1 and 2.

A series of single-photon trials probes this state using mirrors *M* that bring the two branches together, phase shifters ϕ_1 and ϕ_2 that lengthen the two paths by phases ϕ_1 and ϕ_2 , and a second beam splitter *BS2* that mixes the branches together.

Measurement occurs at photon detectors *B1* and *B2*.




Figure 2. Mach-Zehnder interferometer experiment. A photon traverses a beam splitter, travels on two phase-shifted paths to another beam splitter, and is detected.

Figure 3 shows the results(discussed earlier).




Figure 3. Single photon interference, pointing to Dirac's conclusion that "each photon ...interferes only with itself," i.e. each photon follows both paths.

Varying ϕ_1 through 180 degrees causes the photon's state to shift from 100% probability of detection at *B1*, through diminishing probabilities at *B1* and increasing probabilities at *B2*, finally reaching 100% probability of detection at *B2*.

The photon exhibits similar interference upon varying ϕ_2 .

Note that *A*'s state depends only on the phase *difference* $\phi_2 - \phi_1$.

Since single-trial results vary regardless of which phase shifter varies, it is hard to avoid the conclusion that each photon follows both paths.

In fact, let us assume the contrary, namely that each photon follows only one path.

Suppose the phase shifters are set to ensure 100% probability of detection at *B1*.

Under our assumption, this setting guarantees that every photon following path 1, and every photon following path 2, is detected at *B1*.

Suppose path 2 is now blocked without changing the phase settings, so that (still under our one-path assumption) every detected photon must now follow path 1 and be detected at *B1*.

But the experiment shows that, to the contrary, 50% of the detected photons now go to *B2*.

Conclusion: each photon follows both paths.

For a full discussion, see earlier part of this lecture.

As Paul Dirac put it,

"The new theory, which connects the wave function with probabilities for one photon, gets over , the difficulty by making each photon go partly into each of the two components.

Each photon then interferes only with itself."

This illustrates why the "plus" sign in a superposition such as Equation (1) is interpreted by the word “AND".

We turn now to the entangled state Equation (2).

Many Bell inequality tests used polarization-entangled photon pairs to study the full phase dependence of this state.

More useful for this discussion are two interferometer experiments conducted in 1990.

Both of these experiments used *momentum*-entangled photon pairs to conduct Bell inequality tests of the entangled superposition Equation (2) (see earlier discussions).

Figure 4 shows the layout.




Figure 4. The RTO experiments. In each trial, each of two entangled photons travels two phase-shifted paths (one solid, the other dashed) to beam splitters and coincidence detectors. Think of one biphoton spreading from the source along both the solid and the dashed paths.

The source creates entangled pairs of photons *A* (moving leftward) and *B* (moving rightward) by laser down-conversion in a non-linear crystal.

The down-converted photons are prepared in the entirely microscopic state represented by $|\Psi_{AB}\rangle$ (Equation (2)) by selecting four single-photon beams, each a plane wave having a distinct momentum (i.e. wave vector), from the output of the crystal.

Figure 4 resembles two back-to-back Mach- Zehnder interferometer experiments (Figure 2) with BS1 located effectively inside the source.

For simplicity and clarity, Figure 4 differs from the layout shown in the published papers.

In Figure 4, paired photons are directed oppositely.




Figure 4. The RTO experiments. In each trial, each of two entangled photons travels two phase-shifted paths (one solid, the other dashed) to beam splitters and coincidence detectors. Think of one biphoton spreading from the source along both the solid and the dashed paths.

This arrangement would result if the entanglement were prepared by the cascade decay of an atom.

In the actual experiments, however, down-converted photon pairs are emitted into two angular cones, resulting in photons that are not oppositely directed.

Figure 4's simpler geometry is pedagogically useful and has no effect on our arguments.

Although Figure 4 represents each photon as a wave packet spreading along two paths directed leftward or rightward, the composite system *AB* should be regarded as a *single* object, a **"biphoton".**

In each trial, a biphoton spreads outward from the source along two superposed branches.

One branch, represented by the first term $|A1\rangle\,|B1\rangle$ in Equation (2), spreads along the solid path and the other branch, $|A2\rangle\,|B2\rangle$, spreads along the dashed path.

As the biphoton *AB* moves outward along the solid path, *A* encounters a mirror *M,* then a beam splitter *BS* where it transmits and reflects to detectors *A1* and *A2*; photon *B* encounters a mirror *M,* a phase shifter ϕ_B , and a beam splitter *BS* where it transmits/reflects to detectors *B1*/*B2*.

The other half of the entanglement, namely the dashed path, has a similar description.

The experiments record outcomes at four photon detectors equipped with coincidence timers.

Published papers predict the experimental results theoretically and we **follow** their optical-path analysis here(slightly different but equivalent to our earlier arguments).

They begin by calculating the two-point nonlocal quantum field amplitudes $\Psi(Ai, Bj)$ at the four coincidence detectors *(Ai, Bj)*, and from these results they predict single-photon results.

For example, $\Psi(A1, B2)$ has two contributions, one from phase shifts in the beam following the solid path (the first term in $|\Psi_{AB}\rangle$) and the other from the dashed path (the second term).

From Equation (2), assuming distinct plane waves $\exp(i\mathbf{k}\cdot\mathbf{x})$ for each single-photon beam,

$$\Psi(A1, B2) = \frac{1}{2\sqrt{2}}\{\exp(i\phi_w)\exp[i(\phi_x + \phi_B)] + \exp[i(\phi_y + \phi^A)]\exp(i\phi_z)\} \tag{9}$$

where $\phi_w, \phi_x, \phi_y, \phi_z$ are fixed phase-shifts resulting from mirrors and beam splitters, and the additional factor of 1/2 comes from the superpositions created at the two beam splitters.

Using the Born rule, Equation (9) implies the coincidence probability

$$P(A1, B2) = |\Psi(A1, B2)|^2 = \frac{1}{4}[1 + \cos(\phi_B - \phi_A + \phi_v)] \tag{10}$$

where ϕ_u is a fixed phase arising from $\phi_w, \phi_x, \phi_y, \phi_z$.

Similarly,

$$P(A1, B1) = \frac{1}{4}[1 + \cos(\phi_B - \phi_A + \phi_u)] \tag{11}$$

where ϕ_u is another fixed phase.

Remarkably, the sinusoidal terms predict coherent (phase-dependent) nonlocal interference between *A* and *B*, **regardless of their separation**.

There are similar expressions for *P(A2, B1)* and *P(A2, B2)*.

Single-photon predictions then follow. For example, from simple probability theory

$$P(A1) = P(A1, B1) + P(A1, B2)$$
$$= [1+\cos(\phi_B-\phi_A + \phi_u)]/4 + [1+\cos(\phi_B-\phi_A + \phi_v)]/4. \quad (12)$$

The theory papers then show the two fixed phase factors ϕ_u and ϕ_v differ by π:

$$\phi_v = \phi_u + \pi \ (mod\ 2\pi). \quad (13)$$

Thus the sinusoidal terms in Equation (12) *interfere destructively*, and *P(A1) = 1/2 regardless of phase*.

Equations (12) and (13) show this remarkable result to arise from destructive interference of two phase-dependent nonlocal contributions from the distant ***other*** photon *B*!

The result at all four single-photon detectors is the same:

$$P(A1) = P(A2) = P(B1) = P(B2) = 1/2. \quad (14)$$

Unlike the non-entangled single-photon superposition $|\Psi_A\rangle$, where the superposed photon is coherent (phase-dependent) as shown by Figure 3, each entangled photon is **"decohered"** and cannot interfere with itself.

Instead the two photons interfere with each other despite being separated by an arbitrary distance.

More accurately, ***each biphoton interferes with itself***.

Thus no single-photon interference fringes are associated with the state represented by $|\Psi_{AB}\rangle$.

Special relativity requires that this must be the case:

Since single-photon phase dependence could be used to establish an instantaneous communication channel between *A* and *B*, entanglement ***must*** deprive individual photons of their phase.

The result is **nonlocal coherence of the biphoton, AND decoherence of individual photons**.

The decoherence is required by special relativity.

Equation (14) can also be derived by tracing the pure state density operator $|\Psi_{AB}\rangle\langle\Psi_{AB}|$ over one subsystem to obtain the density operator for the other subsystem, which is what we did earlier.

This yields two density operators that appear to be mixtures but are not really "ignorance mixtures" as the word "mixture" is usually understood because the biphoton is in fact not in a mixed state but rather in a pure state represented by $|\Psi_{AB}\rangle$.

That is why our earlier idea was incomplete!

The optical path analysis, above, derives Equation (14) while avoiding these controversial subtleties

A few definitions can put these predictions into more comprehensible form:

If one photon is detected in state 1 and the other in state 2, the two outcomes are said to be "different."

Otherwise, the outcomes are the "same."

Then from Equations (10) and (11), and similar expressions for *P(A2, B1)* and *P(A2, B2)*,

$$P(same) = P(A1,B1) + P(A2,B2) = 1/2[1 + \cos(\phi_B - \phi_A)] \qquad (15)$$

$$P(different) = P(A1,B2) + P(A2,B1) = 1/2[1 - \cos(\phi_B - \phi_A)]. \qquad (16)$$

Their difference, graphed in Figure 5, is called the **"degree of correlation"**:

$$C = P(same) - P(different) = \cos(\phi_B - \phi_A). \qquad (17)$$




Figure 5. Results of the RTO experiment, demonstrating nonlocality (violation of Bell's inequality). The two photons interfere with each other across an arbitrary distance, i.e. each biphoton interferes with itself.

We now explore its physical significance.

5 - INTERPRETATION OF ENTANGLED MICROSCOPIC STATES

The original purpose of RTO's experiments was to demonstrate violations of Bell's inequality by comparing theoretical predictions, Figure 5, with experimental measurements.

The experimental results agreed with Figure 5 and violated Bell's inequality by 10 standard deviations, confirming the nonlocal nature of $|\Psi_{AB}\rangle$.

What does Figure 5 mean conceptually?




Figure 5. Results of the RTO experiment, demonstrating nonlocality (violation of Bell's inequality). The two photons interfere with each other across an arbitrary distance, i.e. each biphoton interferes with itself.

At zero phase difference, where the two phase shifters are set to equal phases, P(same) = 1 and C = P(same) - P(different) = +1.

Thus both stations always agree, despite the presence of beam splitters that randomize each photon prior to detection (see Figure 4).

It is as though coins were flipped at each station and they always came out either both heads or both tails!

Zero is the "measurement" phase angle where B's state is perfectly (and instantaneously) correlated with A's state.

The nonlocality is intuitively obvious:

Each photon acts like a detector of the state of the other photon regardless of separation!

Simply based on this conclusion, we can make an important observation about the entangled MS Equation (4):

***Nonlocality is a central feature of quantum measurements*.**

For small non-zero phase differences, there is a small probability P(different) that results at the two stations will differ, i.e. observation of *B* no longer provides reliable information about *A*.

With increasing phase difference, this unreliability increases until, at $\pi/2$, the two detector pairs are entirely uncorrelated, and C = 0.

As the phase further increases from $\pi/2$ to π, P(different) increases while P(same) decreases, making C more and more negative.

Finally, C = -1 at phase difference π, implying perfect anti-correlation.

Thus *C* is aptly called the "**degree of correlation**".

This description gives us a clear sense of the physical meaning of the fully entangled state Equation (2), indicating precisely which entities are superposed.

The biphoton's phase controls the degree to which the fixed phase-independent 50-50 states of its two spatially separated subsystems are ***statistically correlated***.

Compare this with the phase of the simple superposition $|\Psi_A\rangle$ (Equation (1)), which controls the degree to which the single system *A* is represented by one or the other *state*.

The entities before and after the plus signs in Equations (1) and (2) are conceptually quite different:

Equation (1) sums two *states* while Equation (2) sums two *correlations between states*.

This **distinction is crucial**.

A ***state*** is a situation (or configuration or path) of a single quantum object, but a ***correlation*** is a statistical relationship between two (or more) quantum objects.

A ***superposition*** is the simultaneous existence of two or more states of a single quantum object.

An ***entanglement*** is the simultaneous existence of two or more relationships (specifically, correlations) between the states of two or more quantum objects.

Creating an entanglement is **quite a different matter** from creating a superposition.

Simple superposition of 1 photon.		Entangled superposition of 2 photons		
$\phi_2-\phi_1$	State of photon	$\phi_B-\phi_A$	State of each photon	Correlation between photons
0	100% 1, 0% 2	0	50-50 1 or 2	100% corr, 0% anticorr
$\pi/4$	71% 1, 29% 2	$\pi/4$	50-50 1 or 2	71% corr, 29% anticorr
$\pi/2$	50% 1, 50%2	$\pi/2$	50-50 1 or 2	50% corr, 50% anticorr
$3\pi/4$	29% 1, 71% 2	$3\pi/4$	50-50 1 or 2	29% corr, 71% anticorr
π	0% 1, 100% 2	π	50-50 1 or 2	0% corr, 100% anticorr

Table 1. Comparison between a simple superposition (Fig, 2) and an entangled superposition (Fig. 4). In Fig. 2, single-photon states vary with phase. In Fig. 4, only the *correlation between single-photon states* varies with phase while single-photon states have no phase. Thus *each biphoton is coherent* but its subsystems are incoherent. That is, entanglement *decoheres* each photon while transferring coherence to the biphoton.

To elaborate, Table 1 compares the superposition represented by $|\Psi_A\rangle$ (columns 1-2) with the entanglement represented by $|\Psi_{AB}\rangle$ (columns 3-5) at five different phases.

Column 2 demonstrates interference between the states represented by $|A1\rangle$ and $|A2\rangle$, implying the photon is in a superposition of following both paths and that the *state of A* varies with phase.

The phase dependence in column 2 shows that *A* interferes with itself.

Simple superposition of 1 photon.		Entangled superposition of 2 photons		
$\phi_2-\phi_1$	State of photon	$\phi_B-\phi_A$	State of each photon	Correlation between photons
0	100% 1, 0% 2	0	50-50 1 or 2	100% corr, 0% anticorr
$\pi/4$	71% 1, 29% 2	$\pi/4$	50-50 1 or 2	71% corr, 29% anticorr
$\pi/2$	50% 1, 50%2	$\pi/2$	50-50 1 or 2	50% corr, 50% anticorr
$3\pi/4$	29% 1, 71% 2	$3\pi/4$	50-50 1 or 2	29% corr, 71% anticorr
π	0% 1, 100% 2	π	50-50 1 or 2	0% corr, 100% anticorr

Table 1. Comparison between a simple superposition (Fig, 2) and an entangled superposition (Fig. 4). In Fig. 2, single-photon states vary with phase. In Fig. 4, only the *correlation between single-photon states* varies with phase while single-photon states have no phase. Thus *each biphoton is coherent* but its subsystems are incoherent. That is, entanglement *decoheres* each photon while transferring coherence to the biphoton.

In contrast, column 4 shows that, when the subsystems are represented by the pure state $|\Psi_{AB}\rangle$, neither photon has a phase.

Thus neither photon can interfere with itself, so **neither photon** can be represented by a superposition state.

They are decohered.

Both photons are **represented** by fixed, phase-independent, 50-50 states at all phase angles, **just as though they were in ignorance mixtures (which, however, they are not)**.

But phase dependence has not vanished, it has only been **transferred** to the composite system.

As column 5 reveals, the *degree of correlation between the fixed states of A and B* now varies with phase.

A photon represented by $|\Psi_A\rangle$ is in a coherent (phase-dependent) superposition of being in two states (i.e. of following two *paths)*.

$|\Psi_{AB}\rangle$, on the other hand, represents the coherent superposition of two *correlations between fixed states*.

Instead of two coherent *states* existing simultaneously, two coherent *relationships between states* exist simultaneously.

Neither subsystem is "*smeared*" (as Schrodinger apparently believed);

instead, *only the relationship between subsystems is smeared*.

Briefly, $|\Psi_A\rangle$ is a superposition of *states* and $|\Psi_{AB}\rangle$ is a superposition of *correlations*.

Thus $|\Psi_{AB}\rangle$ is qualitatively different from $|\Psi_A\rangle$.

$|\Psi_A\rangle$ exhibits properties of $|A1\rangle$ AND $|A2\rangle$, where "AND" indicates the superposition.

If you amplify *A* to macroscopic dimensions, you will get a macroscopic superposition.

$|\Psi_{AB}\rangle$ exhibits properties of correlations between $|A1\rangle$ and $|B1\rangle$ AND correlations between $|A2\rangle$ and $|B2\rangle$.

If you amplify *A* and *B* to macroscopic dimensions, you will *not* get a macroscopic superposition, you will simply get **correlations between macroscopic objects**.

The entanglement process transfers the coherence (phase dependence) of each photon to correlations between the two photons, leaving individual photons in mixtures that are incoherent but that are **not** ignorance mixtures.

$|\Psi_{AB}\rangle$ is a "superposition of correlations between properties," in contrast to $|\Psi_A\rangle$ which is a "superposition of properties."

There is a even better way to think about all this:

Regard *AB* as a single object, a biphoton.

Then Equation (2) describes a superposition of this object.

In the experiment (Figure 4), the two superposed states are represented by the solid line and the dashed line.

In this context, it makes no sense to speak of the superposition of a single subsystem, but it does make sense to speak of the superposition of the biphoton.

It is the biphoton that goes through the phases graphed in Figure 5 and indicated in Table 1 column 5.

Both branches (solid and dashed) of the biphoton exist simultaneously.

6 - INTERPRETATION OF THE MEASUREMENT STATE

Earlier we analyzed the microscopic state represented by Equation (2) mathematically, and then interpreted this state physically.

In order for *B* to be a reliable detector, its states must be perfectly correlated with *A*'s states--it must exhibit $|Bi\rangle$ when and only when *A* is represented by $|Ai\rangle$ *(i = 1, 2)*.

Thus Figure 5 implies the MS must be established at zero non-local phase: $\phi_B - \phi_A = 0$.

At this phase, two nonlocal perfect statistical correlations between a phase-independent 50-50 state of *A* and the corresponding phase-independent 50-50 state of *B* exist simultaneously.

As shown earlier, contrary to Schrodinger's description, *neither subsystem state can be "smeared out" (superposed) because neither subsystem has a phase*.

Instead, *correlations between* fixed states of *A* and *B* are smeared as shown in Table 1, while the detector indicates a single definite outcome.

Applying Table 1 to Schrodinger's example, the cat is predicted to be alive in 50% of trials, dead in the other 50%, and never in both states simultaneously.

Phase alterations would not smear the cat, they would smear only the correlations between the cat and the nucleus leading not to a smeared cat but **only** to imperfect detection.

There is no paradoxical macroscopic superposition.

But if neither *A* nor *B* is superposed, what *is* superposed?

What does the MS's "plus" sign really mean?

The answer, from Table 1 at zero phase:

Simple superposition of 1 photon.		Entangled superposition of 2 photons		
$\phi_2-\phi_1$	State of photon	$\phi_B-\phi_A$	State of each photon	Correlation between photons
0	100% 1, 0% 2	0	50-50 1 or 2	100% corr, 0% anticorr
$\pi/4$	71% 1, 29% 2	$\pi/4$	50-50 1 or 2	71% corr, 29% anticorr
$\pi/2$	50% 1, 50%2	$\pi/2$	50-50 1 or 2	50% corr, 50% anticorr
$3\pi/4$	29% 1, 71% 2	$3\pi/4$	50-50 1 or 2	29% corr, 71% anticorr
π	0% 1, 100% 2	π	50-50 1 or 2	0% corr, 100% anticorr

Table 1. Comparison between a simple superposition (Fig, 2) and an entangled superposition (Fig. 4). In Fig. 2, single-photon states vary with phase. In Fig. 4, only the *correlation between single-photon states* varies with phase while single-photon states have no phase. Thus *each biphoton is coherent* but its subsystems are incoherent. That is, entanglement *decoheres* each photon while transferring coherence to the biphoton.

$|A1\rangle$ is perfectly correlated with $|B1\rangle$ AND $|A2\rangle$ is perfectly correlated with $|B2\rangle$, where "AND" represents the superposition.

This simply says *both correlations exist simultaneously:*

$|A1\rangle$ if and only if $|B1\rangle$ AND $|A2\rangle$ if and only if $|B2\rangle$.

Again, there is no paradoxical macroscopic superposition. It's only the *correlations (relationships) between states,* not the states themselves, that are superposed.

Entanglement transforms a superposition of 2 *states* into a superposition of *two correlations between states*.

This makes quantum measurements possible because subsystem states can then be amplified to macroscopic dimensions **without requiring the creation of a macroscopic superposition**.

Neither subsystem is in a macroscopic superposition.

Since neither subsystem is superposed, only a single outcome occurs--a conclusion that also follows from Equation (14).

This single definite outcome occurs instantly upon entanglement, as facilitated by the nonlocal properties of the entangled MS.

Thus we have derived the collapse as an inevitable consequence of entanglement, and have **no need to postulate** such a process.

The MS **is** the collapsed state.

Our conclusion follows merely from standard principles of quantum theory with no other assumptions.

So von Neumann's enigmatic measurement state, Equation (4), is just what we want.

This entangled pure state provides the desired correlations, a single outcome, and the nonlocality required by Einstein's argument.


Note that the collapse is established at the microscopic level, prior to macroscopic amplification.

Next, we provide an example of the sequence of events.

7 - EXAMPLE

The following simple example typifies quantum measurements and illustrates the preceding insights in terms of a specific measurement process.

Consider the set-up in Figure 6.



A single photon traverses a beam splitter, creating the superposition represented by Equation (1) whose branches correspond to separate paths toward widely separated photon detectors.

Analogously to Figure 1, we assume the two detectors are equidistant from the beam splitter.

Each detector contains a photo-sensitive plate that, upon absorbing a photon, **releases** an electron.

Von Neumann's argument implies that, as the two branches of the superposition “approach” the detectors, at some point the branches overlap the detectors sufficiently that the entanglement process represented by Equation (4) occurs, where $|B_{ready}\rangle$ denotes the microscopic state of the detectors prior to entanglement while $|B1\rangle$ and $|B2\rangle$ denote their states following entanglement but prior to amplification and macroscopic recording.

At the instant of interaction, the state jumps from a superposition of two paths of *A* (Equation (1)) to a superposition of two correlations between *A* and *B* (right-hand side of the process in Equation (4)).

This entangled state is not paradoxical.

The right-hand side of Equation (4) entails precisely the proper correlations:

$$|A1\rangle \text{ if and only if } |B1\rangle \quad \text{AND} \quad |A2\rangle \text{ if and only if } |B2\rangle \ .$$

The excitation is transferred to only *one* detector while the other detector remains unexcited.

More correctly, either the solid branch or the dashed branch of the superposed biphoton (Figure 4) is randomly selected.

In fact, it has been shown experimentally and theoretically that the set-up shown in Figure 6 leads to entanglement and that the predicted nonlocal collapse occurs.

The nonlocality of the collapse is verified quantitatively by the experimental violation of a mathematical inequality derived based on QM.

Thus the collapse, a non-linear and irreversible process, occurs at the microscopic level.

Once one photoelectron is released, the process is thermodynamically irreversible because the electron is released into a vast number--a continuum--of free electron states and cannot feasibly be reversed.

This electron triggers an avalanche of other electrons leading to a macroscopic mark at one detector.

Other measurement set-ups follow the same general principles.

For example, in the measurement described by Einstein (discussed earlier), each small region of the detection screen acts as a single detector, and the diffracted electron's quantum state entangles with all these many regions.

Thus the argument above, involving only two detectors, must be extended to N detectors.

8 - SUMMARY AND CONCLUSION

Using only the standard principles of quantum physics, but minus the collapse postulate, we have shown that quantum state collapse occurs as a consequence of the entanglement that occurs upon measurement as described in 1932 by the von Neumann measurement state MS (Equation (4)).

The entangled "measurement state" of a quantum system and its detector is the collapsed state:

It incorporates the required perfect correlations between the system and its detector, it predicts precisely one definite outcome, and it incorporates the nonlocal properties--the instantaneous collapse across all branches of the superposition--that Einstein showed to be required in quantum measurements.

The measurement state Equation (4) does not describe a detector in a paradoxical superposition of displaying multiple outcomes, as had been supposed by Schrodinger and others.

Instead, quantum theory concludes that this state entails just what we expect following a measurement:

The states represented by $|A1\rangle$ and $|B1\rangle$ are perfectly correlated, AND the states represented by $|A2\rangle$ and $|B2\rangle$ are perfectly correlated, where "AND" represents the plus sign in the mathematical representation of the state.

Entanglement entails merely the simultaneous occurrence of two correlations between subsystems, not the simultaneous occurrence of two individual states of either subsystem.

There can be no paradoxical superposition of different detector states or of different system states, because the entanglement has shorn both the detector and the quantum system of their quantum phases - just as Feynmann supposed in the quote I gave earlier!

The phase has been transferred from the individual subsystems to the degree of correlation between subsystems.

To put all of this more directly, the single quantum object *AB* (the biphoton) collapses from a superposition to one of its members.

The measurement state's entanglement and its nonlocal properties, far from being paradoxical, are required in order to guarantee that the collapse occurs simultaneously across all branches of the superposition.

Eight previous insolubility proofs failed because they did not incorporate this required nonlocality.

Nonlocality is a central feature of quantum measurement.

There is no need for a special collapse postulate because the entangled state is the collapsed state.

Collapse occurs instantly upon entanglement.

This analysis should not be regarded as one more interpretation of quantum physics.

It is instead a **correction** of the previous misunderstanding of von Neumann's entangled measurement state.

It is not surprising that this misunderstanding has persisted for nearly 90 years.

After all, entanglement and nonlocality are deeply involved in the measurement problem's proper resolution but they only began to be understood in 1964, leading to a long period of gradual acceptance with confirmation only in 2015.

The delay in understanding measurement stemmed from this delay in understanding nonlocality.

For more than a century, much has been made of the odd and supposedly paradoxical nature of the quantum.

This presumed quantum spookiness has led to an excess of attempted fixes and interpretations.

Many experts have even declared the theory to be not a description of reality at all, but only a mathematical recipe that helps humans predict the results of experiments.

As Niels Bohr put it, **“There is no quantum world. There is only an abstract quantum description.”**

According to this hypothesis, quantum theory doesn’t describe anything real at all, so there’s no cause for concern about collapse of the quantum state and other odd quantum behaviors.

Such an easy resolution of the quantum quandaries amounts to giving up on science’s project of understanding the realities of the natural world.

It’s an extraordinary claim, requiring extraordinary proof. But there is no such proof, and there are no grounds for regarding quanta as any less real than rocks.

Indeed, rocks are made of quanta.

Although it has long been a conceit of humankind to imagine the universe to be centered around us, there is no reason to think that reality comprises only the kinds of things we experience in our own daily lives.

The real world does not fade from existence, nor does it become incomprehensible, at distances that happen to be several powers of ten smaller than teapots.

Atomic and subatomic processes are just as real as teapots and, with the help of technology, accessible to human experimentation and understanding.

From the viewpoint of the macroscopic and classical world that we are pleased to call “normal", there is certainly oddness in wave-particle duality, indeterminacy, quantum states, superposition, nonlocality, measurement, and quantum jumps.

But there are no logical contradictions here, no disagreements with experiment, and nothing that should persuade us that quantum physics is about anything other than the real world.

Quantum physics is either charmingly counterintuitive or maddeningly puzzling, depending on your taste, but it is entirely self-consistent and experimentally accurate.

It’s time to accept it with all its charms and puzzles, and stop trying to repair or reinterpret it.

It’s time, in other words, to relax and admit the world is not as we had thought.

Nature is far more creative than we could have conceived.

Our most fundamental theory is in better shape than its detractors suppose.

Quantum physics is a remarkable treasure trove of far-reaching phenomena and ideas whose surface we have probably only begun to scratch.

It’s time to fully embrace these ideas, incorporating them into our ways of thinking about the universe, about our planet, and about ourselves.

This is a process that will engage our minds and stretch our imaginations far into the future, for quantum physics is, indeed, not what anybody could have imagined.

Remembering one statement from Richard Feynman……

“Since we are not quantum objects we may never be able to see the details of this process, but it clearly happens. “

So let me just say.....

Don’t believe all the hype.

There is no need for many-worlds, multiverse and hidden variables, and so on.

There are no problems with standard quantum mechanics!

Now you have some idea of what I am thinking about and working on in the area of Foundations of QM!

Hope you had fun……..

If you want more details I could continue……..(but it gets more technical)